

OPEN ACCESS

Full open access to this and thousands of other papers at <http://www.la-press.com>.

Correlation Between Peroxisome Proliferation and Up-Regulation of Cytochrome P450 CYP4A and Peroxisomal Beta-Oxidation Fatty Acyl CoA Oxidases (AOX) in the Koala (*Phascolarctos Cinereus*)

SNT Ngo¹, I Stupans² and RA McKinnon²

¹School of Animal and Veterinary Sciences, The University of Adelaide, Roseworthy Campus, Roseworthy, SA 5371, Australia. ²School of Pharmacy and Medical Sciences, University of South Australia, Adelaide, SA 5000, Australia. Email: suong.ngo@adelaide.edu.au

Abstract: Peroxisomes are membrane bound cytoplasmic organelles that are involved in lipid metabolism and other biological functions. In rat and mouse, profound xenobiotic-induced peroxisome proliferation has been reported, with a marked increase in number and size of peroxisomes in liver parenchymal cells and induction of lipid metabolising enzymes, in particular cytochrome P450 CYP4A and peroxisomal β -oxidation palmitoyl CoA oxidases (AOX1). The present study investigates whether the previously observed higher hepatic CYP4A and AOX1 expression in the koala (*Phascolarctos cinereus*), a unique Australian marsupial, compared with rat and human is associated with peroxisome proliferation. Visualisation and quantification of peroxisomes were performed on liver samples from three koalas utilising transmission electron microscopy, with rat and bandicoot livers being used for comparative purposes. Numerous catalase positive peroxisomes, which clearly stand out by their black single membrane globular structures, were detected in all test ultra-thin sections from koala livers. A higher average number of peroxisomes per hepatocyte was observed for the koala, an obligate eucalyptus feeder, compared with non-eucalyptus feeders rat and bandicoot. No species differences in the average size of peroxisomes were detected. This is the first morphological study examining hepatic peroxisomes in an Australian marsupial. The results suggested that dietary eucalyptus constituents might possess peroxisome proliferating activities.

Keywords: marsupials, eucalyptus terpenes, cytochrome P450 CYP4A, peroxisomal beta-oxidation, fatty palmitoyl CoA oxidases AOX1

Gene Expression to Genetical Genomics 2010:3 1–6

This article is available from <http://www.la-press.com>.

© the author(s), publisher and licensee Libertas Academica Ltd.

This is an open access article. Unrestricted non-commercial use is permitted provided the original work is properly cited.



Introduction

Peroxisomes are membrane bound cytoplasmic organelles that are involved in lipid metabolism and many other biological functions including respiration, gluconeogenesis, thermogenesis, and purine catabolism. In most animal cells, peroxisomes are widely distributed and their number, volume, and density remain relatively constant under various physiological conditions.^{1,2} Peroxisomes were first identified in 1954 as single-membrane limited cytoplasmic organelle that contains a finely granular matrix and were named “microbody”.³ In 1956, microbodies were found in the parenchymal cells of rat liver.⁴ Subsequent studies in the 1960s revealed the presence of microbodies with some variation in size and sharp in various vertebrate liver and kidney cells. These microbodies were also found to have different kinds of nucleoid or crystalline core.^{5–7} The enzymes urate oxidase, D-amino acid oxidase, and catalase were then characterised in rat liver microbodies, with the term “peroxisomes” being introduced in 1965 to emphasise the biochemical property of the microbody enzymes to generate hydrogen peroxide.^{2,8,9} In 1968, a histological assay was developed to identify catalase, the marker enzyme of peroxisomes, and initiated the identification of certain peroxisomes in various cell types.¹⁰ Important enzymes found in peroxisomes include enzymes of the fatty acid β -oxidation system, acyl transferase, catalase, urate oxidase, and D-amino acid oxidase.^{2,8–10}

In susceptible species such as rat and mouse, xenobiotic-induced peroxisome proliferation has been reported, which is characterised by a profound increase in number and size of peroxisomes in liver parenchymal cells and an increase in liver weight.¹¹ Xenobiotic-induced peroxisome proliferation has also been found to be related to fatty acid metabolism due to the presence of fatty acid β -oxidation enzymes in peroxisomes. A class of structurally diverse compounds, which are collectively classified as peroxisome proliferators (PPs), have been identified to cause peroxisome proliferation. This occurs via activation of peroxisome proliferator activated receptor-alpha (PPAR- α).^{12,13} Exposure to PPs has been found to induce expression of genes encoding peroxisomal fatty acid β -oxidation enzymes such as palmitoyl-CoA oxidase (AOX1) and cytochrome P450 CYP4As.^{12–14} Rat liver peroxisomes contain three

acyl-CoA oxidases, namely palmitoyl-CoA oxidase, pristanoyl-CoA oxidase, and trihydroxycoprostanoyl-CoA oxidase. Among the three, only palmitoyl-CoA oxidase (AOX1) is inducible by peroxisome proliferators.¹⁵ While rodents are highly responsive to PPs, human and marmosets are weakly or non-responsive. This might be due to having less abundance or lack of hepatic PPAR- α . PPAR- α was first isolated from mouse liver,¹⁶ subsequently cloned from rat,¹⁷ guinea-pig,¹⁴ human,^{18,19} also Amphibia,^{20,21} Reptilian,²² different teleostei species,^{23–29} and *S. purpuratus*.³⁰ *Xenopus* PPAR- α has been suggested to be activated by naturally occurring fatty acids.^{20,21} In fish, the transcription of PPAR- α is also activated in response to a wide spectrum of mammalian PPAR ligands.³¹

Unlike most other species, the koala (*Phascolarctos cinereus*) diet is restricted to eucalyptus leaves, which contain a high proportion of monoterpenes. Previous studies conducted in our laboratories have suggested a possible dietary induction of hepatic CYP4A enzymes in the koala due to eucalyptus terpenes which may act as potent activators of the nuclear receptor PPAR- α .^{32–34} Our subsequent work identified the presence of functional koala liver *AOX1* and *AOX2* isoforms and suggested higher expression of AOX mRNA in the koala liver as compared to rat and human.^{34,35} In humans, rat and other species studied, both *CYP4A* and *AOX* are PPAR- α target genes.^{36–38} In a more current study, PPAR- α has been isolated from the koala liver and its expression was characterised across species.³⁹ The aim of the present study was to examine whether our previously observed higher hepatic CYP4A and AOX expression in the koala, a unique specialist eucalyptus feeder, compared with rat and human, non-eucalyptus feeders, is associated with peroxisome proliferation. This is the first morphological study examining hepatic peroxisomes in an Australian marsupial.

Materials and Methods

Materials

Glutaraldehyde, hydrogen peroxide, and sodium cacodylate trihydrate were obtained from Sigma (St. Louis, MO, USA). Durcupan ACM mixtures 1 and 2 were purchased from Fluka AG, Chemische Fabrik (Bucks, England, UK). DAB (3,3'-diaminobenzidine) was obtained from Biorad (Sydney, NSW, Australia). All solvents and other chemicals were of analytical grade.



Hepatic tissues

Livers were scavenged from road injured male koalas and a bandicoot, ultimately euthanised using phenobarbital. Male Hooded Wistar rats (*Rattus norvegicus*) were used as the source of rat livers. All animals were sexually mature and the Ethic Committees of the appropriate institutions have approved the use of these tissues.

Methods

Visualisation of koala peroxisomes was performed on liver specimens from three koalas. Two rats and a bandicoot were used for comparative purposes. For each animal tissue, several tissue blocks were dissected, randomly from different areas of the liver, and subjected to the EM procedure as described shortly. Blocks without staining with DAB from each animal were used as negative controls.

Specimen preparation

Morphological examination of liver peroxisomes by electron microscopy (EM) in the koala, a eucalyptus feeder, was accomplished using livers of rat and bandicoot, non-eucalyptus feeders, as control samples. All livers sections for light and electron microscopy were dissected within five minutes after sacrifice of the animals, and then immersed immediately in glutaraldehyde solution.

Fixation and DAB staining

Pre-fixation: Fresh tissues were cut out into small blocks with a razor blade and fixed by immersion in at least 10 × volumes of cold 2.5% (v/v) glutaraldehyde in 0.1 M sodium cacodylate buffer (pH 7.4) for 24 hours at 4 °C. The fixed tissues were rinsed twice in cacodylate buffer, each of 10 minutes, with constant rotation, and then stored in the same buffer overnight until required at 4 °C.

DAB staining and post-fixation: The glutaraldehyde pre-fixed blocks were stained for catalase by incubating in at least 10 × volumes of a freshly prepared solution, which contains 0.05% (w/v) DAB in 0.1 M Tris-HCl buffer (pH 8.5) and 0.02% (v/v) hydrogen peroxide, for 60 minutes at 37 °C with gentle agitation. Blocks were then rinsed in cacodylate buffer several times, each of 10 minutes, with constant rotation. DAB-stained blocks were post-fixed for 1 hour at room temperature in 1% (w/v) osmium tetroxide in 0.05 M sodium cacodylate buffer (pH 7.4).

Dehydration

Prior to dehydration, osmium-fixed blocks were rinsed twice in cacodylate buffer, each of 10 minutes, and then washed several times with distilled water, each of 10 minutes. Dehydration of the sections were performed in a graded series of ethanol concentrations (v/v) in distilled water on a rotor as follows: 50% ethanol for 10 minutes, 70% ethanol for 10 minutes then stored overnight, two changes of 70% ethanol each of 10 minutes, two changes of 90% ethanol each of 10 minutes, absolute ethanol for 10 minutes, and then absolute ethanol for 1 hour.

Infiltration and embedding using epoxy resin Durcupan ACM mixtures

Durcupan mixtures number 1 and 2 were prepared as described by the manufacturer (Fluka AG). Infiltration was accomplished on a rotor by gradual and continuous replacement of the dehydrating agent with an embedding medium as follows: 50% (v/v) Durcupan mixture 1 in absolute ethanol for 1 hour at room temperature, 75% (v/v) Durcupan mixture 1 in absolute ethanol for 1 hour at room temperature, 100% (v/v) Durcupan mixture 1 at 50 °C for 1 hour overnight, and then 50% (v/v) Durcupan mixture 2 in Durcupan mixture 1 at 50 °C for 1 hour. Tissue blocks were then placed in small embedding capsules in Durcupan mixture 2 and polymerised at 60 °C for 48 hours, allowed to cool for 4 hours, and the plastic capsule was then removed.

Sectioning and visualisation

Prior to sectioning, semi-thin sections of 0.5 to 1 µm were cut, stained with 10% (w/v) toluidine blue in 5% (w/v) aqueous borax solution, and examined by light microscopy. Morphology studies were then performed using a transmission electron microscope. Ultra-thin sections of 70 to 100 nm, with pale gold interference colour, were obtained to enable adequate penetration of the electron beam for reasonable resolution and contrast.

Quantification and determination of size

Many ultra thin sections were cut from several tissue blocks for quantification and measurement of peroxisomes. Tissue blocks were chosen randomly from different areas of the liver and the dissecting was repeated using the same procedure for each liver sample, to ensure consistency in counting and size measurement. For each individual animal, counting was performed on at least 50 hepatocytes.

The measurement of the size of peroxisomes was accomplished on at least 40 peroxisomes from many different hepatocytes. Due to the small number of animals in the control group, it was not feasible to perform statistical analysis.

Results and Discussion

Visualisation of koala peroxisomes by EM

This is the first EM morphological study examining hepatic peroxisomes in an Australian marsupial. Representative electron micrographs of koala peroxisomes are shown in Figure 1 and 2A. Representative electron micrograph of rat peroxisomes is shown in Figure 2B.




Figure 1. Upper panel (A) DAB-cytochemical visualisation of peroxisomes in koala liver by electron microscopy (original magnification $\times 10500$). Test ultra-thin section of koala liver stained with diaminobenzidine (DAB) showing peroxisomes, indicated by an arrow, that clearly stand out by their single-membrane bound globular structures and a regular striation within peroxisomes. Lower panel (B) DAB-cytochemical visualisation of peroxisomes in koala liver by electron microscopy (original magnification $\times 4 \times 10500$). Test ultra-thin section of koala liver stained with diaminobenzidine (DAB) that shows peroxisomes, indicated by a symbol P, in Figure 1A being enlarged 4 folds featuring striation, indicated by an arrow, within peroxisomes.




Figure 2. DAB-cytochemical visualisation of peroxisomes in koala and rat livers by electron microscopy (original magnification $\times 10500$). Test ultra-thin section of liver stained with diaminobenzidine (DAB) showing peroxisomes, indicated by an arrow, featured by their single-membrane bound globular structures and regular dark striations within peroxisomes. Upper panel (A) Koala liver. Lower panel (B) Rat liver, being used as control animal.

Numerous catalase positive peroxisomes were detected in all test ultra-thin sections from koala livers. As shown in Figure 1, the koala peroxisomes were observed to clearly stand out as dark globular structures that are surrounded by a single membrane and characterised by a very regular striation within peroxisomes. This was consistent with that described in other species.^{6,7} The koala peroxisomes were distinguished from stained lipid droplets and other globular structures of similar sizes by their single-membrane bound structures and their core-striation characteristics. This dense core or nucleoid has been described as crystalline, semi-crystalline, crystalloid, or multi-lamellated and is the most striking structural component of liver peroxisomes.² A representative image of such structures is shown in Figure 1B, where four

peroxisomes in Figure 1A have been enlarged 4 folds, with the striation being clearly visible in the peroxisome located in the lower centre, the middle one in the group of three peroxisomes below the mitochondrion.

It was observed that peroxisomes did not stand out in the EM micrographs of negative ultra-thin sections that were not incubated with DAB. This is because there was formation of precipitated DAB polymer, catalysed by peroxisome marker enzyme catalase, resulting in increased electron density and improved visibility of peroxisomes in ultra-thin sections stained with DAB.⁴⁰

Quantification and measurement of peroxisomes

The average number of peroxisomes per hepatocyte and the average size of peroxisome from each species are shown in Table 1.

As shown in Table 1, a higher average number of peroxisomes per hepatocyte was observed for the specialist eucalyptus feeder koala compared to non-eucalyptus feeders rat and bandicoot. Although the density of peroxisome often varies with the zone distribution of the hepatocytes, it is unlikely that this zone variation has a significant implication in the interpretation of the results of this study. This was because several tissue blocks from each species were cut randomly from different parts of the liver and many ultra-thin sections from each block were selected. The observed higher number of peroxisomes per hepatocyte in the koala might be one of the indications of peroxisome proliferation, suggesting that dietary eucalyptus terpenes may act as peroxisome proliferators. In rat livers, eucalyptus terpenes such as citral have been found to induce CYP4A and peroxisomal β -oxidation enzymes and caused a marked increase in liver weight.⁴¹ No species differences in

the average size of peroxisomes were detected. The average size values of peroxisomes for koala, rat, and bandicoot livers are within the range of values reported in most species studies (0.3 to 1 μ m).⁴²

In summary, the current study described the first morphological study that examines hepatic peroxisomes in an Australian marsupial. The results shown the presence of more peroxisomes in hepatocytes of koala, a specialist eucalyptus feeder, compared to those of rat and bandicoot, non-eucalyptus feeding species, suggesting that dietary eucalyptus terpenes may possess peroxisome proliferating activities. While rat and human have been reported to be highly responsive and non-responsive to peroxisome proliferators, respectively, this study suggested that the marsupial koala may be a responsive species, given that the numbers of peroxisomes are the sole important criterion.

Disclosures

This manuscript has been read and approved by all authors. This paper is unique and is not under consideration by any other publication and has not been published elsewhere. The authors report no conflicts of interest.

References

1. Tolbert NE. Metabolic pathways in peroxisomes and glyoxysomes. *Ann Rev Biochem.* 1981;50:133.
2. De-Duve C, Baudhuin P. Peroxisomes (microbodies and related particles). *Physiol Rev.* 1966;46:323.
3. Rhodin J. Correlation and ultrastructural organisation and functions in normal and experimentally changed peroxisomal convoluted tubule cells of the mouse kidney. Sweden: Institute Stockholm; 1954.
4. Rouiller C, Bernhard W. "Microbodies" and the problem of mitochondrial regeneration in liver cells. *J Biophys Biochem Cytol.* 1956;Suppl 2:355.
5. Allen JM, Beard M. α -hydroxy acid oxidase: localisation in the renal microbodies. *Science.* 1965;149:1507.
6. Shnitka TK. Comparative ultrastructure of hepatic microbodies in some mammals and birds in relation to species difference in uricase activity, pyridine nucleotide levels in the liver of rats fed clofibrate- or pyrazinamide-containing diets. *J Ultrastruct Res.* 1966;16:598.
7. Hruban Z, Jr, Recheigl M. Microbodies and related particles. Morphology, biochemistry and physiology. *Int Rev Cytol.* 1969;Suppl 1:20.
8. De-Duve C, Baudhuin P. Exploring cells with a centrifuge. *Science.* 1975; 189:186.
9. Leighton F, Poole B, Beautay H, et al. Cytochrome P450x. Structure, Function and Metabolism. The larger scale separation of peroxisomes, mitochondria, and lysosomes from the livers of rats injected with Triton WR-1339. *J Cell Biol.* 1968;37:482.
10. Novikoff AB, Goldfischer S. Visualisation of peroxisomes (microbodies) and peroxisomes and mitochondria with diaminobenzidine. *J Histochem Cytochem.* 1969;17:675.
11. Lake BG, Evans JG, Gray TJB, Korosi SA, North CJ. Comparative studies on nafenopin induced hepatic peroxisome proliferation in rat, Syrian hamster, guinea pig and marmoset. *Toxicol Appl Pharmacol.* 1989; 99:148–60.
12. Osumi T, Wen J, Hashimoto T. Two cis-acting regulatory sequences in the peroxisome proliferator-responsive enhancer region of rat acyl-CoA oxidase gene. *Biochem Biophys Res Commun.* 1991;175:866–71.

Table 1. Quantification and measurement of peroxisomes in koala, rat, and bandicoot livers.

Samples	Number of peroxisomes per hepatocyte	Size (μ m)
Koala (n = 3)	20 \pm 2	0.403 \pm 0.032
Rat (n = 2)	11	0.504
Bandicoot (n = 1)	9	0.375

For koala, values were expressed as mean SD of three animals. Results for rat were expressed as a mean of two animals, and those for bandicoot were from a single animal.



13. Sher T, Yi HF, McBride OW, Gonzalez FJ. cDNA cloning, chromosome mapping, and functional characterisation of the human peroxisome proliferator activated receptor. *Biochemistry*. 1993;32:5598–604.
14. Tugwood JD, Holden PR, James NH, Prince RA, Roberts RA. A peroxisome proliferator-activated receptor- α (PPAR- α) cDNA cloned from guinea-pig liver encodes a protein with similar properties to the mouse PPAR- α : implication for species differences in responses to peroxisome proliferators. *Mol Toxicol*. 1998;72:169–77.
15. Schepers L, Van Veldhoven PP, Casteels M, Eyssen J, Mannaerts GP. Presence of three acyl-CoA oxidases in rat liver peroxisomes. An inducible fatty acyl-CoA oxidase, a noninducible fatty acyl-CoA oxidase, and a noninducible trihydroxycoprostanoyl-CoA oxidase. *J Biol Chem*. 1990; 265:5242–6.
16. Issemann I, Green S. Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature*. 1990;347:645–50.
17. Göttlicher M, Widmark E, Li Q, Gustafson JÅ. Fatty acids activate a chimera of the clofibrilic acid-activated receptor and the glucocorticoid receptor. *Proc Natl Acad Sci U S A*. 1992;89:4653–7.
18. Qi C, Zhu Y, Pan JN, et al. Peroxisomal β -oxidation and peroxisome proliferator-activated receptor alpha: an adaptive metabolic system. *Annu Rev Nutr*. 2001;21:193–230.
19. Jow L, Mukherjee R. The human peroxisome proliferator-activated receptor (PPAR) subtype NUC1 represses the activation of hPPAR alpha and thyroid hormone receptors. *J Biol Chem*. 1995;270:3836–40.
20. Dreyer C, Krey G, Keller H, Givel F, Helftenbein G, Wahli W. Control of peroxisome β -oxidation pathway by a novel family of nuclear hormone receptors. *Cell*. 1992;68:879–87.
21. Krey G, Keller H, Mahfoudi A, et al. Xenopus peroxisome proliferator activated receptors: genomic organization, response element recognition, heterodimer formation with retinoid X receptor and activation by fatty acids. *J Steroid Biochem Mol Biol*. 1993;47:65–73.
22. Takada I, Yu RT, Xu HE, et al. Alteration of a single amino acid in peroxisome proliferator-activated receptor- α (PPAR- α) generates a PPAR delta phenotype. *Mol Endocrinol*. 2000;14:733–40.
23. Ruyter B, Andersen O, Dehli A, et al. Peroxisome proliferator activated receptors in Atlantic salmon (*Salmo salar*): effects on PPAR transcription and acyl-CoA oxidase activity in hepatocytes by peroxisome proliferators and fatty acids. *Biochim Biophys Acta*. 1997;1348:331–8.
24. Maglich JM, Caravella JA, Lambert MH, Willson TM, Moore JT, Ramamurthy L. The first completed genome sequence from a teleost fish (*Fugu rubripes*) adds significant diversity to the nuclear receptor superfamily. *Nucleic Acids Res*. 2003;31:4051–8.
25. Maglich JM, Sluder AE, Willson TM, Moore JT. Beyond the human genome: examples of nuclear receptor analysis in model organisms and potential for drug discovery. *Am J Pharmacogenomics*. 2003;3:345–53.
26. Boukouvala E, Antonopoulou E, Favre-Krey L, et al. Molecular characterization of three peroxisome proliferator-activated receptors from the sea bass (*Dicentrarchus labrax*). *Lipids*. 2004;39:1085–92.
27. Raingeard D, Cancio I, Cajaraville MP. Cloning and expression pattern of peroxisome proliferator-activated receptor alpha in the thicklip grey mullet *Chelon labrosus*. *Mar Environ Res*. 2006;62:S113–7.
28. Kondo H, Misaki R, Gelman L, Watabe S. Ligand-dependent transcriptional activities of four torafugu pufferfish *Takifugu rubripes* peroxisome proliferator-activated receptors. *Gen Comp Endocrinol*. 2007; 154:120–7.
29. Tabe Y, Konopleva M, Kondo Y, et al. PPAR γ -active triterpenoid CDDO enhances ATRA-induced differentiation in APL. *Cancer Biol Ther*. 2007;6:1967–77.
30. Goldstone JV, Hamdoun A, Cole BJ, et al. The chemical defense: environmental sensing and response genes in the Strongylocentrotus purpuratus genome. *Dev Biol*. 2006;300:366–84.
31. Leaver MJ, Boukouvala E, Antonopoulou E, et al. Three peroxisome proliferator-activated receptor isoforms from each of two species of marine fish. *Endocrinology*. 2005;146:3150–62.
32. Ngo SNT, Kong S, Kirlich A, McKinnon RA, Stupans I. Cytochrome P450 4A, peroxisomal enzymes and nicotinamide cofactors in koala liver. *Comp Biochem Physiol C*. 2000;127:327–34.
33. Ngo SNT, McKinnon RA, Stupans I. The effects of eucalyptus terpenes on microsomal cytochrome P450 CYP4A, peroxisomal acyl CoA oxidase (AOX) and peroxisome proliferator activated receptor alpha (PPAR α) in the common brush tail possum liver, *Trichosurus vulpecular*. *Comp Biochem Physiol C*. 2003;127:327–34.
34. Ngo SNT, McKinnon RA, Stupans I. Cloning and characterisation of liver CYP4A15 from the koala (*Phascolarctos cinereus*). *Gene*. 2006;376:123–312.
35. Ngo SNT, McKinnon RA, Stupans I. Identification and cloning of two forms of liver peroxisomal fatty Acyl CoA Oxidase from the koala (*Phascolarctos cinereus*). *Gene*. 2003;309:91–9.
36. Johnson EF, Palmer CN, Hsu MH. The peroxisome proliferator-activated receptor: transcriptional activation of the CYP4A6 gene. *Ann N Y Acad Sci*. 1996;804:373–86.
37. Hsu MH, Palmer CM, Griffin KJ, Johnson EF. A single amino acid change in the mouse peroxisome proliferator activated receptor alpha alters transcriptional responses to peroxisome proliferators. *Mol Pharmacol*. 1995;48:559–67.
38. Varanasi U, Chu R, Huang Q, Castellon R, Yeldandi A. Identification of a peroxisome proliferator-responsive element upstream of the human peroxisomal fatty acid acyl-coenzyme A oxidase gene. *J Biol Chem*. 1996;271:2147–55.
39. Ngo SNT, McKinnon RA, Stupans I. Hepatic nuclear receptor PPAR α in the koala (*Phascolarctos cinereus*): Cloning and molecular characterisation. *Comp Biochem Physiol C*. 2007;146:375–82.
40. Roels F, Prest BD, Prest GD. Liver and chorion cytochemistry. *J Inher Metab Dis*. 1995;1(Suppl 18):155–71.
41. Roffey SJ, Walker R, Gibson GG. Hepatic peroxisomal and microsomal enzyme induction by citral and linalool in rats. *Food Chem Toxicol*. 1990;28:403–8.
42. Reddy JK, Lalwani ND. Carcinogenesis by hepatic peroxisome proliferators. Evaluation of the risk of hepatocarcinogenic drugs and industrial plasticizers to humans. *Cirt Rev Toxicol*. 1983;12:1–58.

Publish with Libertas Academica and every scientist working in your field can read your article

“I would like to say that this is the most author-friendly editing process I have experienced in over 150 publications. Thank you most sincerely.”

“The communication between your staff and me has been terrific. Whenever progress is made with the manuscript, I receive notice. Quite honestly, I’ve never had such complete communication with a journal.”

“LA is different, and hopefully represents a kind of scientific publication machinery that removes the hurdles from free flow of scientific thought.”

Your paper will be:

- Available to your entire community free of charge
- Fairly and quickly peer reviewed
- Yours! You retain copyright

<http://www.la-press.com>