

CHEMISTRY OF INDOLES CARRYING A BASIC FUNCTION. PART

VI.<sup>1</sup> SYNTHESIS OF A NEW RING SYSTEM WITH INDOLE NUCLEUS


István Moldvai, Mihály Balázs, Eszter Gács-Baitz, Tünde Platthy, Eszter Temesvári-Major, and Csaba Szántay\*

Institute of Chemistry, Chemical Research Center, Hungarian Academy of Sciences, H-1525 Budapest, POB. 17, Hungary

E-mail: szantay@chem.bme.hu imoldvai@cric.chemres.hu


**Abstract** — Stobbe condensed product half ester (**2E**; *E*-1,3,4,5-tetrahydrobenz[*c,d*]indole-5-ylidene-3'-ethoxycarbonylpropionic acid) afforded a monomer tetracyclic indole derivative (**4**; 5-ethoxycarbonyl-7-acetoxy-1*H*,9*cH*-3,4-dihydronaphto[*c,d,e*]indole) and a dimer (**5**; 5-ethoxycarbonyl-1*H*,9*cH*-3,4-dihydronaphto[*c,d,e*]indole-7-(*O-E*-1,3,4,5-tetrahydrobenz[*c,d*]indole-5-ylidene-3'-ethoxycarbonylethylpropionate) in an unexpected cyclisation.

Stobbe condensation is widely used to prepare  $\alpha,\beta$ -unsaturated acids from ketones.<sup>2</sup> In the hope of constructing ergoline skeleton by *intramolecular* version of the Stobbe condensation, a few succinic diester derivatives were prepared by Uhle<sup>3</sup> and Stoll<sup>4,5</sup> early in the fifties but the expected cyclizations have never been published. (Recently we described the formation of a few tetra- and pentacyclic products starting from similar type of compounds).<sup>6</sup> On the other hand a successful *intermolecular* reaction of Uhle's ketone (**1**), leading to the usual half ester (**2**) in a good yield, was also described by Stoll's group. To elaborate the half ester side chain at C-5, the ester group was hydrolysed to the dicarboxylic acid and then the latter was attempted to transform into a conjugated monocarboxylic acid. However, the end product proved to be an acid anhydride (**3**), and this approach was finished at this level.<sup>4</sup>




Nevertheless the half ester (**2**) seemed to be a valuable intermediate to build up the ring D of the ergoline skeleton. One of our conceptions was to perform an allylic bromination with NBS at C-4, followed by bromine-amine exchange. The starting material (**2**) was prepared by a modified Stobbe condensation (NaH, benzene + EtOH)<sup>7</sup> in 92 % yield. The crude product was formed as an isomeric mixture (**2E:2Z**  $\approx$  8:2; established by NMR examinations) and the dominant **2E**-isomer (where the carboxyl group is oriented toward the aromatic ring) was obtained in pure form by column chromatography. (After a few days the purified isomer (**2E**) was isomerised in solution to an equilibrium mixture of the two isomers, where the ratio of **2E:2Z** turned out to be similar to that in the crude product).


Despite our expectation, the bromination of **2** with NBS gave an unseparable, and unstable mixture which was unsuitable for any further transformation. In order to enhance the chance of the bromination process, *N*-acetylation of Stobbe product (**2**) was attempted. Adopting the Bowman's procedure<sup>8</sup> for protection, **2E** was treated with acetyl chloride in the presence of K<sub>2</sub>CO<sub>3</sub> in methyl ethyl ketone (4 h, rt). After working-up the reaction mixture (extraction, column chromatography), two unexpected products (**4** yield: 20 %; and **5** yield: 15 %) were obtained as crystals. Not a trace of *N*-acetyl derivative of **2** could be isolated. To the best of our knowledge compounds (**4**) and (**5**) represent a new ring system with indole nucleus.



Concerning the reaction sequence, at the outset we can assume the formation of mixed anhydride (**6**) which attacks the aromatic ring at carbon-6 forming keto ester (**7**). This latter compound isomerises and affords phenoxide (**8**), subsequently acylated with acetyl chloride to afford monomer (**4**), or with mixed anhydride (**6**) yielding dimer (**5**).



To avoid the above undesired cyclisation during the *N*-protection of **2**, the carboxyl group was esterified with diazomethane (2 h, 0 °C) affording diester (**9**) (yield: 93 %) as an isomer mixture in about the same ratio as the starting half ester. (Nevertheless, the pure **9E**-isomer could be isolated by crystallization). Diester (**9**) was allowed to react with acetyl chloride in the presence of  $\text{K}_2\text{CO}_3$  but after 2 days only the starting material could be detected. For the *N*-acylation, diester (**9**) was treated with pivaloyl chloride applying the procedure of Goto's group<sup>9</sup> for protection of 3-indolepropionic acid (*n*-BuLi, - 78 °C, pivaloyl chloride, 2 h, 0 °C → rt). In this case *N*-pivaloyl diester (**10**) was obtained as a single *E*-isomer in 37.5 % yield. (The stereochemical arrangement of **10** was verified by NOE measurements).



## EXPERIMENTAL

Mps are uncorrected. MS spectra were run on an AEI-MS-902 (70 eV; direct insertion) and on a Kratos MS-902 mass spectrometers. IR spectra were taken on a Nicolet 205 and Nicolet 7795 FT-IR spectrophotometers. NMR measurements were carried out on a Varian Unity Inova (400 MHz) instrument. Chemical shifts are given relative to TMS = 0.00 ppm.

Glassware was flame dried before use. THF and benzene were distilled from sodium/benzophenone; CH<sub>2</sub>Cl<sub>2</sub> was distilled from P<sub>2</sub>O<sub>5</sub> and EtOH was distilled from Mg.

### *E/Z*-1,3,4,5-Tetrahydrobenz[*c,d*]indole-5-ylidene-3'-ethoxycarbonylpropionic acid (**2**)

Uhle's ketone (**1**, 1.7 g; 10 mmol) was suspended in diethyl succinate (5.0 mL; 30 mmol) at rt. Sodium hydride (1.2 g; 30 mmol; 60% suspension in oil) was washed with dry benzene and added to the above suspension in benzene (5 mL). The reaction mixture was stirred for 0.5 h, then a mixture of benzene and ethanol (10 mL + 0.1 mL) was dropped while stirring the mixture. After 2 h the red crystals (sodium salt of **2**) were filtered off, washed with hexane to remove the excess of the reagent. The crystals were dissolved in a mixture of chloroform (100 mL) and aqueous HCl solution (0.5 N; 30 mL). After extraction, the organic phase was washed with water (3x10 mL), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and the filtrate evaporated to dryness under reduced pressure. The residue was chromatographed (eluent: chloroform + methanol, 10:1). As a result of this purification isomer-**2E** (2.0 g; 66.9 %) and the mixture of isomers (0.75 g; 25.0 %) were obtained.

Compound (**2E**): mp 158-162 °C (ether). IR (KBr): 3381 (νNH indole), 3300-2500 (νOH carboxylic acid), 1711 (νC=O carboxylic acid), 1688 (νC=O conjugated ester), 1618 (νC=C), 1445 (β<sub>s</sub>CH<sub>2</sub>), 1193 (ν<sub>as</sub>COC ester), 1087 (ν<sub>s</sub>COC ester), 755 (γC<sub>Ar</sub>H) cm<sup>-1</sup>. MS (*m/z*, %): 299 (M<sup>+</sup>, 67). <sup>1</sup>H NMR (CDCl<sub>3</sub>), δ (ppm): 1.37 (3H, t, *J*=7.1 Hz, -CH<sub>2</sub>CH<sub>3</sub>), 3.02 (2H, t, *J*=6.1 Hz, H-3), 3.22 (2H, t, *J*=6.1 Hz, H-4), 3.63 (2H, s, H-2'), 4.32 (2H, q, *J*=7.1 Hz, -CH<sub>2</sub>CH<sub>3</sub>), 6.92 (1H, br s, H-2), 7.12 (1H, dd, *J*=7.2 + 1.2 Hz, H-8), 7.18 (1H, dd, *J*=7.2 + 7.5 Hz, H-7), 7.32 (1H, dd, *J*=7.5 + 1.2 Hz, H-6), 8.02 (1H, br s, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>), δ (ppm): 14.13 (-CH<sub>3</sub>), 23.49 (C-4), 31.43 (C-3), 37.70 (C-2'), 61.03 (-OCH<sub>2</sub>), 111.62 (C-8), 113.44 (C-2a), 117.66 (C-6), 118.21 (C-7), 121.37 (C-3'), 122.40 (C-2), 129.00 (C-8b), 134.11 (C-8a), 148.31 (C-5), 169.30 (-CO<sub>2</sub>Et), 177.18 (C-1'). Anal. Calcd. for C<sub>17</sub>H<sub>17</sub>NO<sub>4</sub>, C, 68.21, H, 5.72, N, 4.68. Found C, 68.13, H, 5.68, N, 4.72.

The NMR data of minor isomer (**2Z**) was determined in the equilibrium of the solution. <sup>1</sup>H NMR(CDCl<sub>3</sub>), δ (ppm): 1.15 (3H, t, *J*=7.1 Hz, -CH<sub>2</sub>CH<sub>3</sub>), 2.80 (2H, t, *J*=6.0 Hz, H-3), 3.10 (2H, t, *J*=6.0 Hz, H-4), 3.90 (2H, s, H-2'), 4.20 (2H, q, *J*=7.1 Hz, -CH<sub>2</sub>CH<sub>3</sub>), 6.90 (1H, br s, H-2), 7.08-7.25 (3H, m, H-8, H-7, H-6), 8.0 (1H, br s, NH).

#### **5-Ethoxycarbonyl-7-acetoxy-1*H*,9*cH*-3,4-dihydronaphto[*c,d,e*]indole (4)**

#### **5-Ethoxycarbonyl-1*H*,9*cH*-3,4-dihydronaphto[*c,d,e*]indole-7-(*O-E*-1,3,4,5-tetrahydrobenz[*c,d*]indole-5-ylidene-3'-ethoxycarbonyl ethylpropionate) (5)**

To a solution of **2E** (782 mg; 2.6 mmol) in methyl ethyl ketone (35 mL) at rt, dry potassium carbonate (2.0 g; 14.5 mmol) and acetyl chloride (0.5 mL; 7.0 mmol) was added. The reaction mixture was stirred at the above temperature for 4 h. (After 2 h, a further portion of acetyl chloride (0.2 mL) was added). The reaction mixture was evaporated to dryness under reduced pressure and the residue was dissolved in a mixture of chloroform and water (50 mL + 10 mL). The phases were separated and the organic layer was washed with water (2x5 mL), and dried (Na<sub>2</sub>SO<sub>4</sub>). The filtrate was evaporated and the residue (730 mg) was chromatographed (eluent: hexane + ethyl acetate, 1/1).

The first fraction gave **4** (160 mg; 20 %). From the second fraction, **5** was obtained (211 mg; 15 %).

Compound (**4**): mp 186-192 °C (from hexane + ethyl acetate, 1/1). IR (KBr): 3480, 1785, 1710, 1480, 1400, 1390 cm<sup>-1</sup>. MS (*m/z*, %): 323 (M<sup>+</sup>, 42.5), 313 (6.4), 281 (100), 252(17). <sup>1</sup>H NMR (CDCl<sub>3</sub>), δ (ppm): 1.42 (3H, t, *J*=7.0 Hz, -CH<sub>2</sub>CH<sub>3</sub>), 2.48 (3H, s, COCH<sub>3</sub>), 3.18 (2H, t, *J*=6.1 Hz, H-3), 3.80 (2H, t, *J*=6.1 Hz, H-4), 4.40 (2H, q, *J*=7.0 Hz, -CH<sub>2</sub>-CH<sub>3</sub>), 6.94 (1H, br s, H-2), 7.43 (1H, d, *J*=8.1 Hz, H-8), 7.54 (1H, d, *J*=8.1 Hz, H-9), 7.68 (1H, s, H-6), 8.32 (1H, br s, NH). <sup>13</sup>C NMR (CDCl<sub>3</sub>), δ (ppm): 14.37 (CH<sub>3</sub>), 20.04 (C-3), 20.99 (COCH<sub>3</sub>), 26.46 (C-4), 60.84 (-OCH<sub>2</sub>), 113.20 (C-2a), 114.61 (C-8), 115.21 (C-9), 116.86 (C-6), 117.58 (C-2), 121.66 (C-7a), 124.01 (C-9b), 124.64 (C-5), 127.13 (C-9c), 129.94 (C-9a), 134.81 (C-4a), 144.56 (C-7), 167.14 (-CO<sub>2</sub>Et), 169.86 (-OCOCH<sub>3</sub>). Anal. Calcd. for C<sub>19</sub>H<sub>17</sub>NO<sub>4</sub>, C, 70.57, H, 5.29, N, 4.33. Found C, 70.55, H, 5.32, N, 4.32.

Compound (**5**): mp 156-162 °C (from hexane + ethyl acetate, 1/1). IR (KBr): 3410, 1790, 1700, 1600 cm<sup>-1</sup>. FAB-MS (DMSO+NOBA): 563 (M+H). <sup>1</sup>H NMR (CDCl<sub>3</sub>), δ (ppm): part A: 1.38 (3H, t, *J*=7.0 Hz, -CH<sub>2</sub>CH<sub>3</sub>), 3.15 (2H, t, *J*=6.0 Hz, H-3), 3.78 (2H, t, *J*=6.0 Hz, H-4), 4.39 (2H, q, *J*=7.0 Hz, -CH<sub>2</sub>CH<sub>3</sub>), 6.88 (1H, br s, H-2), 7.38 (1H, d, *J*=8.0 Hz, H-9), 7.45 (1H, d, *J*=8.0 Hz, H-8), 7.72 (1H, s, H-6), 8.35

(1H, br s, NH); part B: 1.44 (3H, t,  $J=7.0$  Hz,  $-\text{CH}_2\text{CH}_3$ ), 3.03 (2H, t,  $J=6.0$  Hz, H-3), 3.28 (2H, t,  $J=6.0$  Hz, H-4), 4.28 (2H, s,  $-\text{COCH}_2$ ), 4.41 (2H, q,  $J=7.0$  Hz,  $-\text{CH}_2\text{CH}_3$ ), 6.85 (1H, br s, H-2), 7.21 (1H, t,  $J=7.4 + 7.1$  Hz, H-7), 7.28 (1H, dd,  $J=7.1 + 1.1$  Hz, H-8), 7.32 (1H, dd,  $J=7.4 + 1.1$  Hz, H-6), 8.04 (1H, br s, NH).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  (ppm): part A: 14.39 ( $-\text{CH}_3$ ), 20.03 (C-3), 26.46 (C-4), 60.84 ( $-\text{OCH}_2$ ), 113.28 (C-2a), 114.65 (C-8), 115.23 (C-9), 116.76 (C-6), 117.58 (C-2), 121.69 (C-7a), 123.88 (C-9b), 124.51 (C-5), 127.09 (C-9c), 129.93 (C-9a), 134.81 (C-4a), 144.62 (C-7), 169.10 ( $-\text{CO}_2\text{Et}$ ); part B: 14.27 ( $-\text{CH}_3$ ), 23.51 (C-4), 31.50 (C-3), 61.03 ( $-\text{OCH}_2$ ), 111.73 (C-8), 113.04 (C-2a), 117.55 (C-6), 118.39 (C-7), 121.57 (C-3'), 122.28 (C-2), 128.81 (C-8b), 129.07 (C-5a), 134.16 (C-8a), 148.54 (C-5), 167.19 + 170.96 ( $2\times\text{CO}_2$ ). Anal. Calcd. for  $\text{C}_{34}\text{H}_{30}\text{N}_2\text{O}_6$ , C, 72.58, H, 5.37, N, 4.98. Found C, 72.55, H, 5.48, N, 4.95.

***E/Z*-Methyl-1,3,4,5-tetrahydrobenz[*c,d*]indolylidene(5)-3'-ethoxycarbonylpropionate (9):**

Half ester (**2**) (450 mg; 1.5 mmol) was dissolved in  $\text{CH}_2\text{Cl}_2$  (50 mL) at  $0^\circ\text{C}$  and diazomethane (10 mmol in 10 mL of  $\text{CH}_2\text{Cl}_2$ ) was dropped. The mixture was stirred for 2 h. The reaction mixture was diluted with methanol (10 mL) and the mixture was evaporated to dryness under reduced pressure. The residue was purified with column chromatography (silica-Merck 9385, eluent: chloroform + methanol, 20/1) to yield **9** (439 mg, 93.0 %). Pure **9E**-isomer could be obtained by crystallization from ether.

Compound (**9E**): mp  $111\text{--}112^\circ\text{C}$ . IR (KBr): 3401, 2981, 2951, 2845, 1736, 1715, 1604,  $1437\text{ cm}^{-1}$ . MS ( $m/z$ , %): 312 ( $\text{M}^+$ , 56).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 1.31 (3H, t,  $J=7.0$  Hz,  $-\text{CH}_2\text{CH}_3$ ), 3.01 (2H, t,  $J=6.1$  Hz, H-3), 3.19 (2H, t,  $J=6.1$  Hz, H-4), 3.78 (3H, s,  $-\text{OCH}_3$ ), 3.84 (2H, s, H-2'), 4.30 (2H, q,  $J=7.0$  Hz,  $-\text{CH}_2\text{CH}_3$ ), 6.91 (1H, br s, H-2), 7.09 (1H, dd,  $J=7.2 + 1.1$  Hz, H-8), 7.17 (1H, dd,  $J=7.5 + 7.1$  Hz, H-7), 7.29 (1H, dd,  $J=7.5 + 1.2$  Hz, H-6), 8.0 (1H, br s, NH).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 14.18 ( $-\text{CH}_3$ ), 23.46 (C-4), 31.33 (C-3), 37.73 (C-2'), 52.09 ( $-\text{OCH}_3$ ), 60.74 ( $-\text{OCH}_2$ ), 111.45 (C-8), 113.40 (C-2a), 117.56 (C-6), 118.16 (C-7), 122.27 (C-2), 122.14 (C-3'), 128.67 (C-8b), 129.04 (C-5a), 134.04 (C-8a), 147.25 (C-5), 169.07 ( $-\text{CO}_2\text{Et}$ ), 172.44 (C-1'). Anal. Calcd. for  $\text{C}_{18}\text{H}_{18}\text{NO}_4$ , C, 69.22, H, 5.80, N, 4.48. Found C, 69.17, H, 5.77, N, 4.51.

The NMR data of the minor **9Z**-isomer was determined from the isomer mixture.

Compound (**9Z**):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 1.16 (3H, t,  $J=7.0$  Hz,  $-\text{CH}_2\text{CH}_3$ ), 2.79 (2H, t,  $J=6.2$  Hz, H-3), 3.03 (2H, t,  $J=6.2$  Hz, H-4), 3.77 (2H, s, H-2'), 3.81 (3H, s,  $-\text{OCH}_3$ ), 4.18 (2H, q,  $J=7.0$  Hz,  $-\text{CH}_2\text{CH}_3$ ),

6.89 (1H, br s, H-2), 6.99 (1H, dd,  $J=7.2 + 1.1$  Hz, H-8), 7.08 (1H, dd,  $J=7.4 + 7.2$  Hz, H-7), 7.24 (1H, dd,  $J=7.4 + 1.1$  Hz, H-6), 7.91 (1H, br s, NH).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 13.71 (-CH<sub>3</sub>), 22.68 (C-4), 29.98 (C-3), 36.70 (C-2'), 51.85 (-OCH<sub>3</sub>), 61.04 (-OCH<sub>2</sub>), 110.84 (C-8), 113.00 (C-2a), 116.46 (C-6), 118.07 (C-7), 122.10 (C-2), 128.05 (C-8b), 128.46 (C-5a), 134.00 (C-8a), 149.60 (C-5), 167.60 (-CO<sub>2</sub>Et), 171.70 (C-1').

***E*-Methyl-*N*-pivaloyl-1,3,4,5-tetrahydrobenz[*c,d*]indolylidene(5)-3'-ethoxycarbonylpropionate (10):**

Diester (**9**) (624 mg; 2.0 mmol) was dissolved in THF (40 mL) and the solution was cooled at -78 °C. To the solution of **9**, *n*-BuLi solution (2.0 mL; 3.2 mmol, 1.6 mol/L in hexane) was added through siringe and the reaction mixture was stirred for 10 min. Then a solution of pivaloyl chloride (0.4 mL; 3.2 mmol) in THF (10 mL) was added and the mixture was stirred for further 3 h, while the temperature of the reaction mixture was allowed to warm up to rt. The mixture was diluted with aqueous 10 % NH<sub>4</sub>Cl solution (50 mL) and ethyl acetate (30 mL). After extraction the phases were separated, the organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated under reduced pressure. The residue was chromatographed (Merck 9385, eluent: hexane + ethyl acetate, 8/2) to yield **10** (297 mg, 37.5 %) as an oil.

IR (KBr): 2971, 2934, 2873, 1736, 1740, 1714, 1693, 1611, 1588, 1478, 1434, 1404, 1343, 1371 cm<sup>-1</sup>. MS ( $m/z$ , %): 396 (M<sup>+</sup>, 82).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 1.35 (3H, t,  $J=7.0$  Hz, -CH<sub>2</sub>CH<sub>3</sub>), 1.51 (9H, s, -C/CH<sub>3</sub>/<sub>3</sub>), 2.95 (2H, t,  $J=6.2$  Hz, H-3), 3.16 (2H, t,  $J=6.2$  Hz, H-4), 3.76 (3H, s, -OCH<sub>3</sub>), 3.80 (2H, s, H-2'), 4.32 (2H, q,  $J=7.1$  Hz, -CH<sub>2</sub>CH<sub>3</sub>), 7.25 (1H, dd,  $J=7.3 + 1.2$  Hz, H-6), 7.32 (1H, dd,  $J=7.3 + 7.2$  Hz, H-7), 7.42 (1H, br s, H-2), 8.32 (1H, dd,  $J=7.2 + 1.2$  Hz, H-8).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 14.18 (-CH<sub>3</sub>), 23.30 (C-4), 28.61 (-C/CH<sub>3</sub>/<sub>3</sub>), 30.16 (C-3), 37.58 (C-2'), 40.97 (-C/CH<sub>3</sub>/<sub>3</sub>), 52.12 (-OCH<sub>3</sub>), 60.88 (-OCH<sub>2</sub>), 118.36 (C-8), 118.81 (C-2a), 119.34 (C-7), 121.74 (C-6), 123.08 (C-3'), 125.49 (C-2), 129.07 (C-5a), 129.92 (C-8b), 135.51 (C-8a), 145.91 (C-5), 168.73 (-CO<sub>2</sub>Et), 172.11 (C-1'), 177.05 (NCO). Anal. Calcd. for C<sub>23</sub>H<sub>27</sub>NO<sub>5</sub>, C, 69.50, H, 6.84, N, 3.52. Found C, 69.48, H, 6.79, N, 3.53.

## ACKNOWLEDGEMENTS

The authors wish to thank *Dr. G. Czira* for MS and *Dr. O. Egyed* for IR spectra. Support for this research under grant No. T 031753 from the *National Scientific Research Foundation (OTKA)* is gratefully acknowledged.

## REFERENCES

1. For part V see: I. Moldvai, E. Temesvári-Major, M. Balázs, E. Gács-Baitz, O. Egyed, and Cs. Szántay, *J. Chem. Res.(S)*, 1999, 687; *J. Chem. Res.(M)*, 1999, 3018.
2. W. S. Johnson and G. H. Daub, *Org. Reactions*, 1951, **6**, 1.
3. F. C. Uhle, *J. Am. Chem. Soc.*, 1951, **73**, 2402.
4. A. Stoll, J. Rutschmann, and T. Petrzilka, *Helv. Chim. Acta*, 1950, **33**, 2257.
5. A. Stoll, and J. Rutschmann, *Helv. Chim. Acta*, 1952, **35**, 141.
6. I. Moldvai, E. Temesvári-Major, E. Gács-Baitz, O. Egyed, Á. Gömör, L. Nyulászi, and Cs. Szántay, *Heterocycles*, 1999, **51**, 2321.
7. G. H. Daub and W. S. Johnson, *J. Am. Chem. Soc.*, 1948, **70**, 418.
8. a) R. E. Bowman, D. D. Evans, J. Guyett, H. Nagy, J. Weale, D. J. Weyell, and A. C. White, *J. C. S. Perkin I*, 1972, 1926; b) R. E. Bowman, D. D. Evans, J. Guyett, J. Weale, and A. C. White, *J. C. S. Perkin I*, 1973, 760.
9. K. Teranishi, S. Hayashi, S. Nakatsuka, and T. Goto, *Tetrahedron Lett.*, 1994, **35**, 8173.