

HETEROCYCLES, Vol. 68, No. 2, 2006, pp. 237 - 245. © The Japan Institute of Heterocyclic Chemistry
Received, 29th September, 2005, Accepted, 11th January, 2006, Published online, 17th January, 2006. COM-05-10581

SYNTHESIS OF SOME BENZO-14-CROWN-4 ETHERS SUBSTITUTED TO 7,8-DIHYDROXY-3-PHENYLCOUMARIN DERIVATIVES

Ümit Salan^a and Mustafa Bulut^{a,*}

^aUniversity of Marmara, Department of Chemistry, Kadıköy, 81040, İstanbul, Turkey, usalan@marmara.edu.tr, mbulut@marmara.edu.tr

Abstract – 7,8-Dihydroxy-3-phenylcoumarin derivatives reacted with 1,2-bis-(3-tosyloxypropoxy)benzene in CH₃CN/alkali carbonate to furnish 3-phenylchromenone-14-crown-4 ether derivatives, which were identified with elemental analysis, IR, ¹H NMR, ¹³C NMR and MS spectroscopy.

INTRODUCTION

Macrocyclic molecules have attracted much attention because of their potential use in a variety of chemical processes, e.g., selective complexing agents for alkaline-earth metal ions, photo-induced electron transfer bio-mimetic studies, etc.¹⁻⁴ The coumarin nucleus is a very interesting chromophore due to its photochemical and photophysical properties and has been used to convert crown ethers and cryptands into fluorescent probes for alkaline and alkaline-earth metal ions.⁵ In addition, it has been shown that 3-phenylcoumarins exhibit strong fluorescence emission intensities due to their high quantum yields.⁶ We have recently synthesised various crown ethers with different chromophore moieties. The crown ether derivatives of the 4-*H* and 4-methyl-6,7-dihydroxy- and 7,8-dihydroxycoumarins displayed the binding effect of alkaline cations on the fluorescence emission spectra.⁷

This paper deals with the synthesis and structure elucidation of novel 3-phenyl-2*H*-chromenone derivatives of 14-crown-4 macrocycles.

RESULTS AND DISCUSSION


Benzo-14-crown-4 ethers (**6a-6g**) substituted to 7,8-dihydroxy-3-phenylcoumarin derivatives were synthesized from the 1,2-bis-(3-tosyloxypropoxy)benzene (**5**) and the corresponding 7,8-dihydroxy-3-phenylcoumarin derivatives (**3a-3g**) which were prepared from 2,3,4-trihydroxybenzaldehyde and corresponding phenylacetic acid in NaOAc/Ac₂O mixture.

Compound (**5**) was reacted with **3a-3g** to give the crown ethers (**6a-6g**) respectively in the presence of Na_2CO_3 in CH_3CN . The residue was chromatographed over a silica gel column eluting with CHCl_3 . The fractions were further purified by preparative methods and the 3-phenylchromenone-14-crown-4 ethers (**6a-6g**) were obtained in 13-21% yield. The synthetic approach of benzo-14-crown-4 ethers (**6a-6g**) is shown in **Scheme 1**. The structures of the compounds (**6a-6g**) were characterized by elemental analysis, ^1H and ^{13}C NMR, MS and IR spectroscopies.



The IR spectra of **6a** showed two absorption at $2876\text{-}2927\text{ cm}^{-1}$ for their C-H stretching frequency. The characteristic absorptions of the carbonyl group (C=O) and benzene ring were appeared at 1720 cm^{-1} and 1625 cm^{-1} respectively. The C-O-C ether chain of the crown ether was characterized by an absorption at $1310\text{-}1090\text{ cm}^{-1}$. A crown ether skeleton could also be deduced from analysis of the ^1H NMR spectral data. The ^1H NMR spectrum of compound (**6a**), which showed two triplets for the methylene protons $[-\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}-]$ at δ 4.31 ($J = 6\text{ Hz}$) and δ 4.48 ($J = 6\text{ Hz}$) and one quintet for the another methylene protons $[-\text{OCH}_2\text{CH}_2\text{CH}_2\text{O}-]$ at δ 2.33 ppm ($J = 6\text{ Hz}$), implied the presence of 14-crown-4 ether. In addition the chemical shifts of the aromatic protons were observed in the region of δ 6.88-7.72 ppm. The structure of newly synthesized **6a** was also checked by MS spectrometry. In the MS spectrum of **6a**, we observed a molecular ion at m/z 444 $[\text{M}]^+$ (calcd for $\text{C}_{27}\text{H}_{24}\text{O}_6$, $M=444$).

The IR spectrum of **6b-6g** showed similar absorptions compared to the spectra of compound (**6a**). The ^1H NMR spectra of compound (**6b**) showed a pair of doublet at δ 7.07 (br d, $J = 8\text{ Hz}$, H-3' and H-5') and δ 7.58 ppm (br d, $J = 8\text{ Hz}$, H-2' and H-6') for aromatic protons and a singlet at δ 2.40 ppm for methyl protons. As expected, the absorptions of the C-O-C ether chain, benzene ring and carbonyl group of compound (**6b**) were appeared at $1300\text{-}1180\text{ cm}^{-1}$, 1625 cm^{-1} and 1720 cm^{-1} respectively.

The IR, ^1H and ^{13}C NMR spectral data of **6c** and **6d** were very similar to those of compound (**6b**). In the ^1H NMR spectra of **6c**, signals were easily assigned to aromatic protons at δ 6.96 (br d, $J = 8\text{ Hz}$, H-3' and H-5') and δ 7.64 (br d, $J = 8\text{ Hz}$, H-2' and H-6') ppm. The methoxy protons showed a singlet at δ 3.85 ppm. In addition, the chemical shifts of the aromatic and ethereal carbons were observed around δ 114.09-160.76 ppm and δ 42.07-69.67 ppm respectively. Spectroscopic data of compound (**6d**) suggested the presence of *o*-methoxy substituted crown ethers. The structure of compound (**6e**) was elucidated by elemental analyses (C, H and N) and other spectroscopic data. The results of spectroscopy supported the structure of the new *p*-nitro substituted 14-crown-4 ether. The ^1H NMR spectrum of **6f** and **6g** showed the expected resonances and integrals due to the protons of these two crown ether derivatives. Also MS spectra and elemental analysis data confirmed the formation of compounds (**6f**) and (**6g**).



	2a	2b	2c	2d	2e	2f	2g
R₁	H	H	H	OCH ₃	H	H	H
R₂	H	H	H	H	H	OCH ₃	OCH ₃
R₃	H	CH ₃	OCH ₃	H	NO ₂	OCH ₃	OCH ₃
R₄	H	H	H	H	H	H	OCH ₃



	6a	6b	6c	6d	6e	6f	6g
R₁	H	H	H	OCH ₃	H	H	H
R₂	H	H	H	H	H	OCH ₃	OCH ₃
R₃	H	CH ₃	OCH ₃	H	NO ₂	OCH ₃	OCH ₃
R₄	H	H	H	H	H	H	OCH ₃

Scheme 1. Synthesis of benzo-14-crown-4 ethers

EXPERIMENTAL

Melting points were measured on a Gallenkamp apparatus uncorrected. IR spectra were recorded by using potassium bromide pellets on a Shimadzu FTIR-8300 spectrophotometer as KBr pellets. Elemental

analyses were performed by the Instrumental Analysis Laboratory of Tübitak-Ankara. ^1H NMR and ^{13}C NMR spectra were recorded in deuteriochloroform or methanol with an instrument Mercury-VX 400 MHz. EI MS spectra were recorded on a Fission UG-ZABSPEC.

The starting chemicals (**1**, **2a-2g**) were purchased from Aldrich or Merck. Initial compounds (**3a-3g**) and 3-phenylchromenone-14-crown-4 ether derivatives (**6a-6g**) have been prepared according to our early studies.^{8,9}

General procedure for the synthesis of 7,8-dihydroxy-3-phenylchromenones (**3a-3g**)

A mixture of (4.62 g, 30 mmol) of 2,3,4-trihydroxybenzaldehyde (**1**), (4.08 g, 30 mmol) of phenylacetic acid and (150 mmol) of sodium acetate in 25 mL of acetic anhydride was heated with stirring at 160 °C under N_2 for 6 h. After removal of acetic acid by distillation, the resulting mixture was treated with $\text{CH}_3\text{OH}/10\%$ HCl, and the precipitates were collected by filtration. The dried product was purified by recrystallisation from ethanol.

7,8-Dihydroxy-3-phenyl-2H-chromen-2-one (3a)

A mixture of compound (**1**) (4.62 g, 30 mmol), phenylacetic acid (**2a**) (4.08 g, 30 mmol), dry sodium acetate (12.30 g 150 mmol) in acetic anhydride (25 mL) was treated as described above to afford **3a**, 6.45 g (85%), mp 254 °C (ethanol); IR (KBr) 3361-3463, 3055, 1727, 1600-1625, 1200 cm^{-1} ; ^1H -NMR (MeOD, 400 MHz): δ 6.52 (d, $J = 8.5$ Hz, 1H, H-6), 7.13 (d, $J = 8.5$ Hz, 1H, H-5), 7.18 (br d, $J = 8$ Hz, 2H, H-2' and 6'), 7.25 (t, $J = 8$ Hz, 1H, H-4'), 7.38 (t, $J = 8$ Hz, 2H, H-3' and 5'), 7.88 (s, 1H, H-4). Anal. Calcd for $\text{C}_{15}\text{H}_{10}\text{O}_4$: C 70.86; H 3.96. Found C 70.63; H 3.77.

7,8-Dihydroxy-3-(4'-methylphenyl)-2H-chromen-2-one (3b)

Compound (**1**) (4.62 g, 30 mmol), 4-methylphenylacetic acid (**2b**) (4.50 g, 30 mmol), dry sodium acetate (12.30 g 150 mmol) in acetic anhydride (25 mL) was treated as described above to afford **3b**, 4.58 g (57%), mp 220 °C (ethanol); IR (KBr) 3400, 3120, 2850-2953, 1710, 1590-1625, 1200 cm^{-1} ; ^1H -NMR (MeOD, 400 MHz): δ 2.41 (s, 3H, CH_3), 6.70 (d, $J = 8.5$ Hz, 1H, H-6), 6.83 (d, $J = 8.5$ Hz, 1H, H-5), 6.95 (d, $J = 8$ Hz, 2H, H-3' and H-5'), 7.18 (d, $J = 8$ Hz, 2H, H-2' and H-6'), 7.85 (s, 1H, H-4). Anal. Calcd for $\text{C}_{16}\text{H}_{12}\text{O}_4$: C 71.64; H 4.47. Found C 71.03; H 4.32.

7,8-Dihydroxy-3-(4'-methoxyphenyl)-2H-chromen-2-one (3c)

Compound (**1**) (4.62 g, 30 mmol), 4-methoxyphenylacetic acid (**2c**) (4.98 g, 30 mmol), dry sodium acetate (12.30 g 150 mmol) in acetic anhydride (25 mL) was worked up as described above to give **3c**, 8.05 g (94%), mp 244 °C (ethanol); IR (KBr) 3400, 3100, 2870-2930, 1725, 1600-1630, 1150 cm^{-1} ; ^1H -NMR (MeOD, 400 MHz): δ 3.85 (s, 3H, OCH_3), 6.70 (d, $J = 8.5$ Hz, 1H, H-6), 6.83 (d, $J = 8.5$ Hz, 1H,

H-5), 6.95 (d, $J = 8$ Hz, 2H, H-3' and H-5'), 7.18 (d, $J = 8$ Hz, 2H, H-2' and H-6'), 7.85 (s, 1H, H-4). Anal. Calcd for $C_{16}H_{12}O_5$: C 67.60; H 4.25. Found C 67.45; H 4.02.

7,8-Dihydroxy-3-(2'-methoxyphenyl)-2H-chromen-2-one (3d)

Compound **(1)** (4.62 g, 30 mmol), 2-methoxyphenylacetic acid (**2d**) (4.98 g, 30 mmol) dry sodium acetate (12.30 g, 150 mmol) in acetic anhydride (25 mL) was reacted as described above to give **3d**, 7.10 g (83%), mp 209 °C (ethanol); IR (KBr) 3400, 3100, 2850-2978, 1720, 1600-1620, 1180 cm^{-1} ; 1H -NMR (MeOD, 400 MHz): δ 3.70 (s, 3H, OCH₃), 6.70 (d, $J = 8.5$ Hz, 1H, H-6), 6.83 (d, $J = 8.5$ Hz, 1H, H-5), 6.93 (d, $J = 8$ Hz, 1H, H-3'), 7.18 (dd, $J = 8$ Hz, $J = 2$ Hz, 1H, H-6'), 7.25 (t, $J = 8$ Hz, 2H, H-5' and H-4'), 7.63 (s, 1H, H-4). Anal. Calcd for $C_{16}H_{12}O_5$: C 67.60; H 4.23. Found C 66.20; H 3.95.

7,8-Dihydroxy-3-(4'-nitrophenyl)-2H-chromen-2-one (3e)

Compound **(1)** (4.62 g, 30 mmol), 4-nitrophenylacetic acid (**2e**) (5.43 g, 30 mmol) dry sodium acetate (12.30 g, 150 mmol) in acetic anhydride (25 mL) was treated as described above to afford **3e**, 6.10 g (68%), mp 270 °C (ethanol); IR (KBr) 3400, 3050, 1700, 1600-1620, 1100 cm^{-1} ; 1H -NMR (MeOD, 400 MHz): δ 6.70 (d, $J = 8.5$ Hz, 1H, H-6), 6.83 (d, $J = 8.5$ Hz, 1H, H-5), 6.85 (d, $J = 8$ Hz, 2H, H-2' and H-6'), 7.46 (d, $J = 8$ Hz, 2H, H-3' and H-5'), 7.90 (s, 1H, H-4). Anal. Calcd for $C_{15}H_9NO_6$: C 60.20; H 3.01. Found C 60.45; H 3.45.

7,8-Dihydroxy-3-(3',4'-dimethoxyphenyl)-2H-chromen-2-one (3f)

A mixture of compound **(1)** (4.62 g, 30 mmol), 3,4-dimethoxyphenylacetic acid (**2f**) (5.88 g, 30 mmol), dry sodium acetate (12.30 g, 150 mmol) in acetic anhydride (25 mL) was treated as described above to give **3f**, 7.72 g (82%), mp 220 °C (ethanol); IR (KBr) 3400, 3100, 2853-2950, 1700, 1580-1620, 1190 cm^{-1} ; 1H -NMR (CD₃OD, 400 MHz): δ 3.76 (s, 3H, OCH₃), 3.78 (s, 3H, OCH₃), 6.71 (d, $J = 8.5$ Hz, 1H, H-6), 6.88 (d, $J = 8.5$ Hz, 1H, H-5), 6.94 (d, $J = 8.5$ Hz, 1H, H-5'), 7.15 (dd, $J = 8.5$ and 2 Hz, 1H, H-6') 7.22 (d, $J = 2$ Hz, 1H, H-2'), 7.80 (s, 1H, H-4). Anal. Calcd for $C_{17}H_{14}O_6$: C, 64.97; H, 4.49. Found: C 64.82; H 4.12.

7,8-Dihydroxy-3-(3',4',5'-trimethoxyphenyl)-2H-chromen-2-one (3g)

Compound **(1)** (4.62 g, 30 mmol), 3,4,5-trimethoxyphenylacetic acid (**2g**) (6.78 g, 30 mmol), dry sodium acetate (12.30 g, 150 mmol) in acetic anhydride (25 mL) was worked up as described above to give **3g**, 8.87g (86%), mp 238-239 °C (ethanol); IR (KBr) 3361-3463, 3055, 2851-2953, 1727, 1625, 1242 cm^{-1} ; 1H NMR (CD₃OD, 400 MHz): δ 3.69 (s, 3H, OCH₃), 3.78 (s, 6H, OCH₃), 6.72 (d, $J = 8.5$ Hz, 1H, H-6), 6.90 (br s, 2H, H-2' and H-6'), 6.96 (d, $J = 8.5$ Hz, 1H, H-5), 7.88 (s, 1H, H-4). Anal. Calcd. for $C_{18}H_{16}O_7$: C, 62.79; H, 4.68. Found: C, 61.60; H, 4.50.

Synthesis of 1,2-bis(3-tosyloxypropoxy)benzene (5)

Catachol (24.25 g, 0.22 mol) was dissolved in 220 mL of C₂H₅OH and the mixture was stirred at 50 °C under N₂. Finely ground NaOH (21.2 g, 0.53 mol) was added. After dissolving, 1-chloro-3-hydroxypropane (47 g, 0.50 mol) was added intervals over an ½ h period, the whole was stirred for 20 h at reflux temperature. The reaction mixture was worked up by filtering the undissolved salt then evaporated. The residue was dissolved in 750 mL of CHCl₃. The CHCl₃ solution was washed with 1M 100 mL of NaOH and 2 x 200 mL of H₂O. The CHCl₃ solution was dried over sodium sulfate and solvent stripped to yield 1,2-bis(3-hydroxypropoxy)benzene (4). 32 g (64%), mp 60 °C; IR (KBr) 3388, 3085, 2871-2900, 1600-1640, 1257-1053 cm⁻¹; ¹H-NMR (CDCl₃, 400 MHz): δ 2.05 (quintet, *J* = 5 Hz, 4H, -CH₂CH₂CH₂-), 3.85 (t, *J* = 5 Hz, 4H, -CH₂OH-), 4.16 (t, *J* = 5 Hz, 4H, -OCH₂CH₂CH₂OH), 6.87-6.93 (m, 4H). Anal. Calcd for C₁₂H₁₈O₄: C 63.70; H 7.96. Found C 63.40; H 7.80.

Compound (4) (5 g, 22 mmol) was dissolved in minimum amount of pyridine and cooled to -5 °C. Tosyl chloride (10.48 g, 55 mmol) in 10 mL of pyridine was added intervals over an ½ h. The reaction mixture was stirred for 2 days at 0-5 °C and then poured into the ice bath. The precipitate was filtered to yield 1,2-bis-(3-tosyloxypropoxy)benzene (5) 2 g (17%), mp 25 °C (ethanol); IR (KBr) 3056, 2873-2900, 1465-1590, 1386-1035 cm⁻¹; ¹H-NMR (CDCl₃, 400 MHz): δ 2.25 (s, 6H), 2.26 (quintet, *J* = 6 Hz, 4H, -CH₂CH₂CH₂-), 3.77 (t, *J* = 6 Hz, 4H, -CH₂OH-), 4.20 (t, *J* = 6 Hz, 4H, -OCH₂CH₂CH₂-), 6.91-6.95 (m, 12H). Anal. Calcd for C₂₆H₃₀O₈S₂: C 58.43; H 5.62; S 12.00. Found C 58.30; H 5.25; S 12.44.

General procedure for the synthesis of 3-phenylchromenone-crown ethers (6a-6g)

1,2-Bis-(3-tosyloxypropoxy)benzen (5) (1 g, 1.8 x 10⁻³ mol), 7,8-dihydroxy-3-phenylcoumarin derivatives (3a-3g) (1.8 x 10⁻³ mol) and sodium carbonate (0.38 g, 3.6 x 10⁻³ mol) in 200 mL of acetonitrile were stirred for 5 days at 70-80 °C under N₂ atmosphere. The reaction mixture was acidified using 10% HCl and extracted with CHCl₃. The crude product was chromatographed over a silica gel column eluting with chloroform, a gradient of chloroform-methanol up to 100% methanol.

7,8-[6,7-Benzo-1,5,8,12-tetraoxadecylene]-3-phenyl-2H-1-benzopyran-2-one (6a)

Compound (5) (1 g, 1.8 x 10⁻³ mol), 3a (0.457 g, 1.8 x 10⁻³ mol) and sodium carbonate (0.38 g, 3.6 x 10⁻³ mol) in acetonitrile (200 mL) was treated as describe above to afford 6a, (0.12 g, 15%), mp 110 °C (benzene); IR (KBr) 3100, 2927-2876, 1720, 1625, 1310, 1090; ¹H NMR (400 MHz, CDCl₃): δ 2.33 (quintet, *J* = 6 Hz, 4H, H-3 and H-10), 4.31 (t, *J* = 6 Hz, 4H, H-4 and H-9), 4.48 (t, *J* = 6 Hz, 4H, H-2 and H-11), 6.88 (d, *J* = 8 Hz, 1H, coumarin H-6), 6.90-7.00 (m, 4H, benzo), 7.18 (d, *J* = 8 Hz, 1H, coumarin H-5), 7.40 (m, 3H, coumarin H-3', H-4' and H-5'), 7.68 (br d, *J* = 8 Hz, 2H, coumarin H-2' and H-6'), 7.72 (s, 1H, coumarin H-4); EIMS: *m/z* (rel. int.) 444 [M]⁺ (27%). Anal. Calcd for C₂₇H₂₄O₆: C 72.97; H 5.44. Found C 72.60; H 5.35.

7,8-[6,7-Benzo-1,5,8,12-tetraoxadecylene]-3-[4'-methylphenyl]-2H-1-benzopyran-2-one (6b)

Compound (**5**) (1 g, 1.8×10^{-3} mol), **3b** (0.482 g, 1.8×10^{-3} mol) and sodium carbonate (0.38 g, 3.6×10^{-3} mol) in acetonitrile (200 mL) was treated as describe above to afford **6b**, (0.14 g, 17%), mp 146-147 °C (benzene); IR (KBr) 3100, 2927-2876, 1720, 1625, 1300, 1180; ^1H NMR (400 MHz, CDCl_3): δ 2.32 (quintet, $J = 6$ Hz, 4H, H-3 and H-10), 2.40 (s, 3H, $-\text{CH}_3$), 4.31 (t, $J = 6$ Hz, 4H, H-4 and H-9), 4.48 (t, $J = 6$ Hz, 4H, H-2 and H-11), 6.88 (d, $J = 8$ Hz, 1H, coumarin H-6), 6.88-6.95 (m, 4H, benzo), 7.07 (br d, $J = 8$ Hz, 2H, coumarin H-3' and H-5'), 7.18 (d, $J = 8$ Hz, 1H, coumarin H-5), 7.58 (br d, $J = 8$ Hz, 2H, coumarin H-2' and H-6'), 7.70 (s, 1H, coumarin H-4); EIMS: m/z (*rel. int.*) 459 $[\text{M}+\text{H}]^+$ (13%). Anal. Calcd for $\text{C}_{28}\text{H}_{26}\text{O}_6$: C 73.35; H 5.67. Found C 73.58; H 5.88.

7,8-[6,7-Benzo-1,5,8,12-tetraoxadecylene]-3-[4'-methoxyphenyl]-2H-1-benzopyran-2-one (6c)

Compound (**5**) (1 g, 1.8×10^{-3} mol), **3c** (0.511 g, 1.8×10^{-3} mol) and sodium carbonate (0.38 g, 3.6×10^{-3} mol) in acetonitrile (200 mL) was treated as describe above to afford **6c**, (0.18 g, 21%), mp 180-181 °C (benzene); IR (KBr) 3070, 2927-2876, 1720, 1625, 1310, 1120; ^1H NMR (400 MHz, CDCl_3): δ 2.32 (quintet, $J = 6$ Hz, 4H, H-3 and H-10), 3.85 (s, 3H, OCH_3), 4.31 (t, $J = 6$ Hz, 4H, H-4 and H-9), 4.48 (t, $J = 6$ Hz, 4H, H-2 and H-11), 6.87 (d, $J = 8$ Hz, 1H, coumarin H-6), 6.90-7.00 (m, 4H, benzo), 6.96 (br d, $J = 8$ Hz, 2H, coumarin H-3' and H-5'), 7.18 (d, $J = 8$ Hz, 1H, coumarin H-5), 7.64 (br d, $J = 8$ Hz, 2H, coumarin H-2' and H-6'), 7.67 (s, 1H, coumarin H-4); ^{13}C NMR (400 MHz, CDCl_3): δ 42.07, 68.27, 69.67, 55.71, 112.94, 114.09, 122.25, 123.56, 123.97, 124.95, 139.02, 139.63, 147.22, 147.73, 148.30, 150.15, 159.94, 159.98, 160.76.; EIMS: m/z (*rel. int.*) 474 $[\text{M}]^+$ (20 %). Anal. Calcd for $\text{C}_{28}\text{H}_{26}\text{O}_7$: C 70.87; H 5.48. Found C 70.58; H 5.38.

7,8-[6,7-Benzo-1,5,8,12-tetraoxadecylene]-3-[2'-methoxyphenyl]-2H-1-benzopyran-2-one (6d)

Compound (**5**) (1 g, 1.8×10^{-3} mol), **3d** (0.511 g, 1.8×10^{-3} mol) and sodium carbonate (0.38 g, 3.6×10^{-3} mol) in acetonitrile (200 mL) was reacted as describe above to afford **6d**, (0.11 g, 13%), mp 175 °C (benzene); IR (KBr) 3055, 2927-2876, 1720, 1625, 1290, 1100; ^1H -NMR(400 MHz, CDCl_3): 2.20 (q, $J = 5$ Hz, 4H, H-3 and H-10), 3.80 (s, 3H, OCH_3), 4.13 (t, $J = 5.5$ Hz, 4H, H-4 and H-9), 4.51 (t, $J = 5.5$ Hz, 4H, H-2 and H-11), 6.86 (d, $J = 8$ Hz, 1H, coumarin H-6), 6.90-7.00 (m, 4H, benzo), 6.94 (br d, $J = 8$ Hz, 1H, coumarin H-6'), 7.05 (d, $J = 8$ Hz, 1H, coumarin H-5), 7.10 (dd, $J = 7$ Hz, $J = 1.5$ Hz, 1H, coumarin H-3'), 7.22 (t, $J = 7$ Hz, 1H, coumarin H-5'), 7.35 (t, $J = 7$ Hz, 1H, coumarin H-4'), 7.70 (s, 1H, coumarin H-4). Anal. Calcd for $\text{C}_{28}\text{H}_{26}\text{O}_7$: C 70.87; H 5.48. Found C 71.05; H 5.35.

7,8-[6,7-Benzo-1,5,8,12-tetraoxadecylene]-3-[4'-nitrophenyl]-2H-1-benzopyran-2-one (6e)

Compound (**5**) (1 g, 1.8×10^{-3} mol), **3e** (0.538 g, 1.8×10^{-3} mol) and sodium carbonate (0.38 g, 3.6×10^{-3} mol) in acetonitrile (200 mL) was treated as describe above to afford **6e**, (0.18 g, 20%), mp 206-208 °C

(benzene); IR (KBr) cm^{-1} 3100, 2927-2876, 1720, 1625, 1270, 1120; ^1H NMR (400 MHz, CDCl_3): δ 2.33 (quintet, $J = 6$ Hz, 4H, H-3 and H-10), 4.31 (t, $J = 6$ Hz, 4H, H-4 and H-9), 4.50 (t, $J = 6$ Hz, 4H, H-2 and H-11), 6.90-7.00 (m, 4H, benzo), 6.93 (d, $J = 8$ Hz, 1H, coumarin H-6), 7.20 (d, $J = 8$ Hz, 1H, coumarin H-5), 7.81 (s, 1H, coumarin H-4), 7.83 (d, $J = 8$ Hz, 2H, coumarin H-2' and H-6'), 8.22 (br d, $J = 8$ Hz, 2H, H-3' and H-5'). Anal. Calcd for $\text{C}_{27}\text{H}_{23}\text{NO}_8$: C 66.25; H 4.70, N 2.86. Found C 65.75; H 4.38, N 2.83.

7,8-[6,7-Benzo-1,5,8,12-tetraoxadecylene]-3-[3',4'-dimethoxyphenyl]-2H-1-benzopyran-2-one (6f)

Compound (**5**) (1 g, 1.8×10^{-3} mol), **3f** (0.57 g, 1.8×10^{-3} mol) and sodium carbonate (0.38 g, 3.6×10^{-3} mol) in acetonitrile (200 mL) was treated as describe above to afford **6f**, (0.13 g, 14%), mp 182°C (benzene); IR (KBr) 3050, 2927-2876, 1720, 1625, 1350, 1150; ^1H -NMR (400 MHz, CDCl_3): 2.18 (q, $J=6$ Hz, 4H, H-3 and H-10), 3.66 (t, $J = 6$ Hz, 2H, H-9), 3.82 (s, 3H, 4'- OCH_3), 3.86 (s, 3H, 3'- OCH_3), 4.08 (t, $J = 6$ Hz, 2H, H-4), 4.26 (t, $J = 5.5$ Hz, 2H, H-11), 4.46 (t, $J = 5.5$ Hz, 2H, H-2), 6.81 (d, $J = 8$ Hz, 1H, coumarin H-6), 6.84-6.93 (m, 4H, benzo), 7.08 (d, $J = 8$ Hz, 1H, coumarin H-5), 7.56 (dd, $J = 9$ Hz, $J=1.5$ Hz, 1H, coumarin H-6'), 7.63 (s, 1H, coumarin H-4), 7.64 (d, $J = 1.5$ Hz, 1H, coumarin H-2'), 7.72 (d, $J = 8.5$ Hz, 1H, coumarin H-5'); EIMS: m/z (*rel. int.*) 505 $[\text{M}+\text{H}]^+$ (10%), 504 $[\text{M}]^+$ (25%). Anal. Calcd for $\text{C}_{29}\text{H}_{28}\text{O}_8$: C 69.04; H 5.55. Found C 69.80; H 5.38.

7,8-[6,7-Benzo-1,5,8,12-tetraoxadecylene]-3-[3',4',5'-trimethoxyphenyl]-2H-1-benzopyran-2-one (6g)

Compound (**5**) (1 g, 1.8×10^{-3} mol), **3g** (0.62 g, 1.8×10^{-3} mol) and sodium carbonate (0.38 g, 3.6×10^{-3} mol) in acetonitrile (200 mL) was treated as describe above to afford **6g**, (0.17 g, 18%), mp 185°C (benzene); IR (KBr) 3100, 2927-2876, 1720, 1625, 1300, 1110; ^1H -NMR (400 MHz, CDCl_3): 2.08 (q, $J = 6$ Hz, 4H, H-3 ve H-10), 3.82 (s, 3H, OCH_3), 3.84 (s, 6H, OCH_3), 4.14 (t, $J = 6$ Hz, 4H, H-4 ve H-9), 4.23 (t, $J = 6$ Hz, 2H, H-2), 4.40 (t, $J = 6$ Hz, 2H, H-11), 6.80-6.90 (m, 4H, benzo), 7.42 (d, $J = 6$ Hz, 1H, coumarin H-6), 7.44 (d, $J = 7$ Hz, 1H, coumarin H-5), 7.60 (s, 1H, coumarin H-4), 7.62 (d, $J = 2$ Hz, 2H, coumarin H-2' ve H-6'); EIMS: m/z (*rel. int.*) 535 $[\text{M}+\text{H}]^+$ (5 %), 534 $[\text{M}]^+$ (20 %). Anal. Calcd for $\text{C}_{30}\text{H}_{30}\text{O}_9$: C 67.41; H 5.62. Found C 67.15; H 5.48.

ACKNOWLEDGEMENTS

We are thankful to The Scientific and Technical Research Council of Turkey (TUBITAK) for financial support.

REFERENCES

1. B. Valeur and I. Leray, *Coordin. Chem. Rev.*, 2000, **3**, 205.
2. B. R. Pandya and Y. K. Agrawal, *Dyes and Pigments*, 2000, **52**, 161.
3. M. V. Alfimov and S. P. Gromov In: W. Rettig, B. Strehmel, S. Schrader, and H. Seifort, Ed.,

Applied fluorescence in chemistry, biology and medicine, Springer-Verlag, Berlin, Heidelberg, 1999, 161.

4. C. J. Pedersen, *Angew. Chem., Int. Ed. Engl.*, 1988, **100**, 1053.
5. Ç. Erk, A. Göçmen, and M. Bulut, *Supramol. Chem.*, 2000, **11**, 49; B. Raju and T. S. Vadarajan, *J. Phys. Chem.*, 1994, **98**, 8903.
6. M. Bulut and Ç. Erk, *J. Heterocycl. Chem.*, 2001, **38**, 1291.
7. Ç. Erk, A. Göçmen, and M. Bulut, *J. Incl. Phenom.*, 1998, **31**, 319; Ç. Erk, A. Göçmen, and M. Bulut, *J. Incl. Phenom.*, 2000, **37**, 441.
8. S. Abdurrahmanoğlu, C. Gündüz, Ü. Çakır, B. Çiçek, and M. Bulut, *Dyes and Pigments*, 2005, **65**, 197.
9. C. Gündüz, Ü. Salan, N. Özkul, İ. Başaran, Ü. Çakır, and M. Bulut, *Dyes and Pigments*, in press.