

NUCLEOPHILIC AROMATIC SUBSTITUTION OF A METHOXY GROUP BY AN ALKYLTHIO GROUP IN THE CASE OF 10-SUBSTITUED 2-METHOXY-9-THIOACRIDINONES

Djeridi Dhiif, Jean-Pierre Galy*, and Jacques Barbe

Groupe d'Etudes et de Recherches en Chimie Thérapeutique Organique Physique

27, Bd Jean-Moulin, 13385 Marseille Cedex 5, France


José Elguero

Instituto de Química Médica, Calle Juan de la Cierva, 3, 28006 Madrid, Spain

Abstract - The synthesis and nmr study (^1H and ^{13}C) of eleven acridine derivatives are reported. ^{13}C Chemical shifts are very sensitive to the oxygen or sulfur nature of the substituents at positions 2 (alkoxy vs thioalkoxy) and 9 (acridinones vs thioacridinones). The rather unusual replacement of an alkoxy group by a thioalkoxy group in thioacridinones, when treated by alkyl mercaptans in the presence of aluminium chloride, is described.

We became interested in the 2-hydroxy-10-methyl-9-thioacridinone as starting material for the preparation of bisacridine derivatives.¹⁻⁵ The simplest way to obtain this product would obviously be the demethylation of 2-methoxy-10-methyl-9-thioacridinone **1a**. However, use of hydrobromic acid, which gives excellent results in the case of methoxyacridinones,⁶ led to the 2-hydroxy-10-methyl-9-acridinones **2**.⁷⁻⁸ Hence to avoid the simultaneous hydrolysis of the thione group, two alternative methods were tried. The use of aluminium iodide in benzene⁹ failed and the starting material was recovered. In contrast, aluminium chloride in ethanethiol¹⁰⁻¹¹ afforded 2-ethylthio-10-methyl-9-thioacridinones **3a**. The compound so obtained was somewhat unexpected because there is usually a demethylation under these experimental conditions.¹⁰ Yet, substituted derivatives were obtained from methoxynaphthalene¹¹ but until now, this has never been mentioned in the field of heterocyclic chemistry. The same reaction was observed when the substituent on the nitrogen is benzyl **1b** or ethyl **1c** - prepared by thiation of the 9-oxo analogs **4** according to the method of Smolders¹² - or when ethanethiol is replaced by propanethiol : compounds **3b-3e**. In contrast, the reaction failed with butanethiol, with 9-(10H)-thioacridinones, and with 2-methoxyacridinones (both NH and N-methyl). According to the opinion of Node,¹¹ this could be due to the decrease in the nucleophilic reactivity.

These results are schematized in the Figure and data about the compounds prepared are listed in Table 1.



Compounds	R	2-substituent
1a	CH ₃	OCH ₃
1b	CH ₂ C ₆ H ₅	OCH ₃
1c	C ₂ H ₅	OCH ₃
3a	CH ₃	SC ₂ H ₅
3b	CH ₂ C ₆ H ₅	SC ₂ H ₅
3c	C ₂ H ₅	SC ₂ H ₅
3d	CH ₃	SC ₃ H ₇
3e	C ₂ H ₅	SC ₃ H ₇
4a	CH ₃	OCH ₃
4b	CH ₂ C ₆ H ₅	OCH ₃
4c	C ₂ H ₅	OCH ₃

The ^{13}C chemical shifts gathered in Table II, prove that the SR group occupies the same position, C-2, that the methoxy leaving group. Actually, with a reference to 9-thioacridinone, ^{13}C resonance peaks are shifted of about 33 ppm for C-2, 20 ppm for C-1, and 10 ppm for C-3 in the case of the 2-methoxy substitution, whilst resonance peaks are respectively shifted of only 8 ppm for C-2, and approximately 0.5 to 1.5 ppm for C-1 or C-3, in the case of compounds **3**. Added to this, ^1H nmr leads to the same conclusion. Indeed, protons 1 and 8 are deshielded (dd, $\delta = 9.1$ ppm) when there are no substituents branched on the

9-thioacridinone nucleus. In the case of substitution with a methoxy group in position 2, there are two signals for the protons under discussion [H (1), $\delta = 9.1$ ppm and H (8), $\delta = 8.6$ ppm], whilst in the case of substitution with a thioethyl group, chemical shifts are 9.0 ppm for H (1) and 9.1 ppm for H (8). This is due to the deshielding effect of the thioether group.

Such a structure can be also deduced from mass spectra. Indeed with reference to the data presented in Table III, attention must be drawn, on the one hand, on the parent-ion peaks at m/z 285, 351 and 299 for **3a**, **3b** and **3c**, and, on the other hand, on ion peaks at m/z 241 which correspond to the loss of all alkyl substituents for **3a**, **3b**, and **3c**, whilst ion peaks at m/z 270 is due to the loss of the benzyl group, for **3b**.

Thus, the mechanism involved corresponds to the S_NAr mechanism previously proposed by Nade in the case of aromatic hydrocarbons,¹¹ with the possibility of a mono-electronic process.

Table III. Mass Spectra Data (m/z)

3a	285 (M^+) (100), 257 (45), 256 (30), 252 (24), 241 (23)
3b	361 (M^+) (38), 333 (1), 300 (1.5), 270 (50), 242 (7), 241 (8), 91 (100)
3c	299 (M^+) (100), 284 (12), 271 (33), 270 (25), 266 (16), 242 (13), 241 (15)

Table 1. ¹H Nmr data

Compound	mp (°C)	Yield %	Analysis (%)			Calc H	N	S	¹ H-Nmr ^a (CDCl ₃ -TMS) δ (ppm), J (Hz)
			Found C	Found H	Found N				
1a (13)	175	37	70.59	5.10	5.49	12.55	70.71	5.29	12.50 9.2 (d, J = 8.5 Hz, 1H); 9.6 (s, 1H); 7.7 to 7.3 (m, 5H); 4.1 (s, 3H); 4.0 (s, 3H)
1b	184	76	76.13	5.16	4.23	9.63	76.26	5.23	4.34 9.63 9.3 (d, J = 8.5 Hz, 1H); 8.6 (s, 1H); 7.7 to 6.9 (m, 10H); 5.6 (s, 2H); 3.9 (s, 3H)
1c	149	35	71.38	5.58	5.20	11.90	71.46	5.77	5.35 11.70 9.2 (d, J = 8.5 Hz, 1H); 8.6 (s, 1H); 7.7 to 7.2 (m, 5H); 4.5 (q, J = 7 Hz, 2H); 3.9 (s, 3H); 1.5 (t, J = 7 Hz, 3H)
3a	152	92	67.37	5.26	4.91	22.45	67.39	5.20	4.69 22.60 9.0 (d, J = 7.5 Hz, 1H); 8.9 (s, 1H); 7.7 to 7.2 (m, 5H); 3.9 (s, 3H); 3.0 (q, J = 7 Hz, 2H); 1.4 (t, J = 7 Hz, 3H)
3b	163	34	73.13	5.26	3.88	17.73	73.14	5.33	3.83 17.90 9.3 (d, J = 7 Hz, 2H); 9.1 (s, 1H); 7.7 to 7.0 (m, 10H); 5.6 (s, 2H); 3.1 (q, J = 7 Hz, 2H); 1.4 (t, J = 7 Hz, 3H)
3c	129	83	68.23	5.69	4.68	21.40	68.37	5.76	4.52 21.20 9.2 (d, J = 7 Hz, 1H); 9.1 (s, 1H); 7.7 to 7.2 (m, 5H); 4.5 (q, J = 7 Hz, 2H); 3.9 (q, J = 7 Hz, 2H); 1.7 to 1.3 (m, 6H)
3d	140	38	68.23	5.69	4.68	21.40	68.35	5.78	4.73 21.50 9.1 (d, J = 8 Hz, 1H); 9.0 (s, 1H); 7.6 to 7.2 (m, 5H); 3.9 (s, 3H); 2.9 (t, J = 7 Hz, 2H); 2.5 to 1.5 (m, 2H) 1.0 (t, J = 7 Hz, 3H)
3e	143	37	69.01	6.07	4.47	20.45	69.04	5.19	4.51 20.30 9.2 (d, J = 8 Hz, 1H); 9.1 (s, 1H); 7.7 to 7.2 (m, 5H); 4.5 (q, J = 7 Hz, 2H); 3.0 (t, J = 7.2 Hz, 2H); 1.9 to 1.5 (m, 5H); 1.1 (t, J = 7 Hz, 3H)
4a (14,15)	138	91	75.31	5.44	5.86		75.47	5.58	5.77 8.5 (d, J = 8 Hz, 1H); 7.9 (s, 1H); 7.6 to 7.2 (m, 10H); 3.9 (s, 3H); 3.8 (s, 3H)
4b (14,16)	148	56	30.00	5.40	4.44		30.05	5.53	4.57 8.6 (d, J = 8.5 Hz, 1H); 7.9 (s, 1H); 7.6 to 7.2 (m, 10H); 5.5 (s, 2H); 3.9 (s, 3H)
4c (14,17,18)	212	68	75.99	5.93	5.53		75.92	6.04	5.68 8.6 (d, J = 8 Hz, 1H); 8.0 (s, 1H); 7.8 to 7.3 (m, 5H); 4.5 (q, J = 7 Hz, 2H); 3.9 (s, 3H); 1.5 (t, J = 7 Hz, 3H)


a) Recorded with a Varian A 60 spectrometer.

Table II. ¹³C Nmr chemical shifts δ (in ppm)

Carbons	COMPOUNDS													
	1a ^a	1b ^a	1c ^a	3a ^a	3b ^a	3c ^a	3d ^a	3e ^a	4a ^a	4b ^c	4c ^b	4d ^b	9-Theacridinone ^c	
1	110.53	110.86	110.89	131.56	131.91	132.26	131.77	132.29	105.9	103.76	107.12	129.9		
2	155.83	156.16	155.86	131.18	131.59	131.51	131.42	131.48	159.88	154.38	122.6	153.8	122.6	
3	125.36	125.56	125.73	133.62	134.08	134.02	133.76	134.00	124.00	123.7	134.19	124.30	132.8	
4	115.36	115.79	115.05	115.51	115.95	115.17	115.49	115.16	117.9	121.09	114.25	120.2		
5	117.14	117.62	116.87	116.17	116.64	115.86	116.12	115.81	115.8	119.17	116.20	120.2		
6	133.42	133.75	133.67	134.71	135.27	135.29	134.92	135.26	133.5	139.90	133.48	132.8	132.8	
7	122.68	123.00	122.74	122.78	123.23	122.96	122.91	122.95	120.7	126.49	120.71	122.6	122.6	
8	131.97	132.09	132.44	130.10	130.53	130.85	130.18	130.73	126.5	130.71	127.81	129.9	129.9	
9	198.12	199.43	198.01	198.89	200.41	199.20	199.24	199.23	175.9	169.46	177.21	192.9	192.9	
11	132.00	132.60	131.64	131.91	132.44	132.00	132.32	132.30	120.6	119.53	123.22	129.0	129.0	
12	132.61	132.33	132.20	135.67	135.94	134.94	135.82	134.89	137.0	141.10	136.29	138.0	138.0	
13	137.09	137.23	136.11	137.14	137.43	136.36	137.30	136.36	141.8	143.72	141.00	138.0	138.0	
14	130.81	131.04	130.91	131.18	131.65	131.54	131.36	131.57	122.3	128.32	121.74	129.0	129.0	
α	55.79	55.63	55.63	27.95	28.05	28.05	35.98	36.01	55.4	57.43	55.71	-	-	
β	-	-	-	14.20	14.26	14.24	22.47	22.48	-	-	-	-	-	
γ	-	-	-	-	-	-	13.51	13.50	-	-	-	-	-	
α'	34.82	51.86	42.16	34.65	51.76	42.07	34.72	42.05	33.6	54.91	40.90	-	-	
β'	-	135.26	13.05	-	135.09	12.88	-	12.89	-	134.57	12.65	-	-	
γ'	-	125.68	-	-	125.70	-	-	-	-	126.83	-	-	-	
δ	-	129.34	-	-	129.36	-	-	-	-	131.40	-	-	-	
ε'	-	128.03	-	-	128.05	-	-	-	-	126.32	-	-	-	

a) CDCl₃/TMS
 b) TFA-d₄/TMS
 c) DMSO-d₆/TMS

* These assignments may be reversed



EXPERIMENTAL

10-Alkyl-2-methoxy-9-acridinones, 4 ; General Procedure : A stirred mixture of 2-methoxy-9(10H)-acridinone²⁰ (0.015 mol), alkylating agent (0.0375 mol), triethylbenzylammonium chloride (0.0075 mol), 50 % potassium hydroxide (75 ml) and toluene (150 ml) is heated for 5 days at 110°C. The toluene layer is separated, washed three times with water (about 50 ml each time), dried with sodium sulfate, and evaporated in vacuo. The crude is recrystallized from ethanol.

10-Alkyl-2-methoxy-9-thioacridinones, 1 ; General Procedure : A mixture of 10-alkyl-2-methoxy-9-acridinones **4** (0.010 mol), tetraphosphorus decasulfide (0.010 mol) and hexamethylphosphoramide (30 ml) is stirred at 100°C for 24 h. The solution is then poured out into 500 ml of cold water. The precipitate obtained is filtered out, washed with cold water and recrystallized from methanol.

10-Alkyl-2-alkylthio-9-thioacridinones, 3 ; General Procedure : Aluminium chloride (0.022 mol) is added to 10-alkyl-2-methoxy-9-thioacridinones **1** (0.004 mol). The use of any solvent is not necessary. The mixture is mildly heated for 10 min at 30°C, safe from moisture. Alkyl mercaptan (0.15 mol) is then added, and the mixture is stirred for a time which depends on the nature of the mercaptan (2 h with ethyl mercaptan, 7 h with propyl mercaptan), before the solution be poured out into water (500 ml). The red precipitate obtained is filtered out, washed with cold water and recrystallized from acetonitrile.

REFERENCES

1. G. Giovannangeli, J. C. Soyfer, J. P. Galy, and J. Barbe, J. Chem. Engineer. Data, 1986, 31, 127.
2. R. Vidal, J. P. Galy, E. J. Vincent, A. M. Galy, and J. Barbe, Heterocycles, 1986, 24, 1419.
3. S. Ammor, A. M. Galy, J. P. Galy, and J. Barbe, J. Chem. Engineer. Data, 1986, 31, 374.
4. R. Vidal, J. P. Galy, E. J. Vincent, and J. Barbe, Synthesis, 1987, 148.
5. D. Dhif, J. P. Galy, E. J. Vincent, and J. Barbe, Heterocycles, 1989, 29, 1149.
6. G. K. Hugues, N. K. Matheson, A. T. Norman, and E. Ritchie, Austr. J. Sci. Res., 1952, A5, 206.
7. J. W. Schulenberg, Ger. Offen, 1972, 2, 759, 468 (Chem. Abstr., 1979, 91, 74487g).
8. Jap. Kokai, 1977, 77, 142, 077 (Chem. Abstr., 1978, 89, 43152f).
9. S. Andersson, Synthesis, 1985, 437.
10. M. Node, K. Nishide, K. Fuji, and E. Fujita, J. Org. Chem., 1980, 45, 4275.
11. M. Node, K. Nishide, K. Ohta, K. Fuji, E. Fujita, H. Hori, and S. Imayama, Chem. Pharm. Bull., 1983, 31, 545.
12. R. R. Smolders, J. Hanuise, R. Coomans, V. Proietto, N. Voglet, and A. Waefelaer, Synthesis, 1982, 493.

13. R. Faure, J. P. Galy, E. J. Vincent, A. M. Galy, J. Barbe, and J. Elguero, Spectrosc. Lett., 1983, 16, 431.
14. K. Eilen and S. Nitzche, J. Prakt. Chem., 1939, 153, 225.
15. A. Mahamoud, J. P. Galy, E. J. Vincent, A. M. Galy, and J. Barbe, J. Heterocycl. Chem., 1982, 19, 503.
16. I. Mester and D. R. Bergenthal, Z. Naturforsch., 1979, 34B, 516 (Chem. Abstr., 1979, 91,74757c).
17. B. A. Chafiaa, C. Montginoul, E. Toreilles, and L. Giral, Bull. Soc. Chim. Fr., 1980, 2, 345.
18. C. Blanchard, J. M. Fabre, C. Montginoul, B. A. Chafiaa, E. Toreilles, and L. Giral, J. Heterocycl. Chem., 1978, 15, 149.
19. J. W. Schulenberg, U.S. Patent, 1981, 4, 244, 954 (Chem. Abstr., 1981,94, 208727r).
20. K. Lehmstedt and H. Hundertmark, Ber., 1931, 64, 2386.

Received, 7th February, 1990