

**A SYNTHESIS OF 1,2,4-TRIAZOLO[1,5-f]PHENANTHRIDINES BY REARRANGEMENTS OF 1,2,5-OXADIAZOLES INVOLVING AN NCN SEQUENCE WITH THE IMINE NITROGEN IN AN AROMATIC HETEROCYCLIC RING**


Giuseppe Cusmano, Gabriella Macaluso, Michelangelo Gruttadauria, and Silvestre Buscemi

Dipartimento di Chimica Organica, Università di Palermo,  
Via Archirafi, 20, 90123, Palermo, Italy

**Abstract** — A synthetic pathway to 1,2,4-triazolo[1,5-f]phenanthridine system by a base catalyzed rearrangement of 3-(6-phenanthridine)-amino-1,2,5-oxadiazoles (5a,b) has been investigated. This ring transformation is the first example of the applicability of the mononuclear heterocyclic rearrangement involving an NCN sequence to the synthesis of bridged nitrogen systems.


In previous researches, we have reported a new method for the synthesis of nitrogen containing polycondensed heterocyclic systems, through an initial  $6\pi$  electrocyclic reaction involving a hydrazone side chain of a five membered heterocycle, followed by ring opening and ring closure processes.<sup>1-4</sup>

The present paper describes the results of our researches aimed to annelate polycondensed heterocyclic ring through an alternative synthetic route. The key step of this new method, as depicted in the Scheme 1, is the rearrangement of the type 1  $\longrightarrow$  2.



Scheme 1

This rearrangement shows a close resemblance with the previously reported rearrangement of arylformamidinoazoles into 1,2,4-triazole derivatives.<sup>5-8</sup> In the present case the nucleophilic imine nitrogen of the formamidino side-chain is incorporated in an aromatic heterocyclic ring.



a: R = Me; b: R = Ph

Scheme 2

As an approach to examine the proposed route to azapolycondensed systems attention was directed towards the study of the rearrangement reactions leading to the known 1,2,4-triazolo[1,5-f]phenanthridine system, previously synthesized through the addition - rearrangement method.<sup>1</sup>

An access to the starting 3-(N-heteroaryl)aminoazoles involved the nucleophilic replacement of halogen atom by 3-aminoazoles. Taking into consideration that such nucleophilic substitution takes place from 6-chlorophenanthridine (3) only when the reactants are heated beyond their melting point<sup>9</sup> and that the 1,2,5-oxadiazole ring has a low reactivity towards mononuclear heterocyclic rearrangements,<sup>8</sup> we thought of heating 3-amino-1,2,5-oxadiazoles (4a,b) and 3 to obtain the intermediate 3-(6-phenanthridine)amino-1,2,5-oxadiazoles (5a,b). Compounds (5a,b) should not rearrange in such thermal conditions.

In fact the amino derivatives (5a,b) were obtained in high yield from equimolar amounts of 6-chlorophenanthridine (3)<sup>10</sup> and 3-amino-4-methyl-1,2,5-oxadiazole (4a)<sup>11</sup> or 3-amino-4-phenyl-1,2,5-oxadiazole (4b)<sup>12</sup> by heating the mixture in an oil bath at 110-140 °C for 1 h and subsequently at 160-180 °C for 1 h.

The transformation of the amino derivatives (5a,b) was carried out by refluxing in anhydrous DMF with equimolar amounts of potassium t-butoxide and the products (6a,b) were obtained in 90-92% yield (see Experimental). In order to verify the structures of the rearrangement products, the oximes (6a,b) were synthesized by oximation reaction of 2-acyl-1,2,4-triazolo[1,5-f]phenanthridines (11a,b), obtained by the addition-rearrangement method outlined in Scheme 2. The rearrangement of 5a afforded only the oxime (E-6a) identical with the oxime obtained through the oximation reaction of 11a; on the other hand, the rearrangement of 5b led to a mixture of E- and Z-6b identical with the oximation mixture of 2-benzoyl-1,2,4-triazolo[1,5-f]phenanthridine (11b).<sup>13</sup> In the IR spectra of compounds 6, the O-H stretching bands at 3270 cm<sup>-1</sup> and 3180 cm<sup>-1</sup> are indicatives for E and Z isomers, respectively. However, in the <sup>1</sup>H NMR spectra in DMSO solution, the OH protons of both E and Z isomers of the oxime (6b) appeared as a broad signal at low field.

The present ring transformation is the first example of the applicability of the mononuclear heterocyclic rearrangement of type 1 → 2 to the synthesis of bridged nitrogen systems. The success of this method seems to depend on the ease of the formation of an N-N bond through an electrocyclic reaction of the conjugate anion of compounds 5.

## EXPERIMENTAL

Melting points were determined with a Kofler hot-stage apparatus; ir spectra (nujol mull) were recorded on a Perkin-Elmer infrared spectrophotometer (model 297); uv spectra (ethanol) were determined with a Varian Superscan 3 spectrophotometer;  $^1\text{H}$  nmr spectra were recorded on a Varian EM 360 spectrometer. Chemical shifts are reported as  $\delta$  values (ppm) relative to TMS as internal standard. Hplc analyses were performed with a Perkin-Elmer Series 10 instrument, by using a C-18 SIL-X-10 Perkin-Elmer column, eluting with methanol, and a Silica 50-10 Perkin-Elmer column, eluting with dichloromethane/ethyl acetate.

### General Method for the Preparation of the 3-(6-Phenanthridine)amino-1,2,5-oxadiazoles (5a,b).

A mixture of 6-chlorophenanthridine (3)<sup>10</sup> (300 mg, 1.4 mmol) and equimolar amounts of 3-amino-4-methyl- (4a)<sup>11</sup> or 3-amino-4-phenyl-1,2,5-oxadiazole (4b)<sup>12</sup> was heated in an oil bath at 110° or 140° for 1 h. The reaction mixture was then kept for 1 h at 160° or 180°. After cooling this mixture was triturated with a minimum amount of chloroform and the resultant solid filtered off.

Compound 5a (R = Me) colourless needles from DMF/H<sub>2</sub>O (yield 77%), mp 170°C; ir:  $\nu_{\text{max}}$ : 3270, 3190 cm<sup>-1</sup> (NH), 1635, 1605 cm<sup>-1</sup> (C=N);  $^1\text{H}$  nmr (DMSO-d<sub>6</sub>)  $\delta$ : 2.40 (s, 3H, CH<sub>3</sub>), 7.2-8.1 (m, 5H, H-2,3,4,8,9), 8.2-8.9 (m, 3H, H-1,7,10), 10.90 (broad s, 1H, NH). Anal. Calcd for C<sub>16</sub>H<sub>12</sub>N<sub>4</sub>O: C, 69.55; H, 4.38; N, 20.28. Found: C, 69.37; H, 4.27; N, 20.33.

Compound 5b (R = Ph) colourless needles from DMF (yield 71%), mp 245°C; ir:  $\nu_{\text{max}}$ : 3260, 3190 cm<sup>-1</sup> (NH), 1625, 1605 cm<sup>-1</sup> (C=N). Anal. Calcd for C<sub>21</sub>H<sub>14</sub>N<sub>4</sub>O: C, 74.54; H, 4.17; N, 16.56. Found: C, 74.48; H, 4.21; N, 16.61.

### Rearrangement of the Amine (5a) into the E Oxime of 2-Acetyl-1,2,4-triazolo-[1,5-f]phenanthridine (6a).

A mixture of 5a (200 mg, 0.72 mmol) and equimolar amounts of potassium t-butoxide (81 mg, 0.72 mmol) in dry DMF (10 ml) was heated at 120-140° for 5 min. After cooling, the E-oxime of 2-acetyl-1,2,4-triazolo[1,5-f]phenanthridine (6a) (180 mg, 90%) was filtered off. Colourless needles from DMSO/H<sub>2</sub>O, mp 270°C;

ir:  $\nu_{\max}$ : 3270  $\text{cm}^{-1}$  (OH), 1610, 1595  $\text{cm}^{-1}$  (C=N);  $^1\text{H}$  nmr (DMSO- $d_6$ )  $\delta$ : 2.40 (s, 3H,  $\text{CH}_3$ ), 7.5-8.1 (m, 4H, H-5,8,9,12), 8.3-8.9 (m, 4H, H-6,7,10,11), 11.80 (very broad, 1H, NOH). Anal. Calcd for  $\text{C}_{16}\text{H}_{12}\text{N}_4\text{O}$ : C, 69.55; H, 4.38; N, 20.28. Found: C, 69.43; H, 4.41; N, 20.37.

Rearrangement of the Amine (5b) into the E and Z Oximes of 2-Benzoyl-1,2,4-triazolo[1,5-f]phenanthridine (6b).

A mixture of 5b (200 mg, 0.59 mmol) and equimolar amounts of potassium *t*-butoxide (66 mg, 0.59 mmol) in dry DMF (10 ml) was heated at 120-140° for 2 h. After cooling, dilution with water (10 ml) gave a crude product which turned out (hplc) to be a mixture of the E and Z isomers of the oxime of 2-benzoyl-1,2,4-triazolo[1,5-f]phenanthridine (6b) (184 mg, 92%), mp 283-285°C; ir:  $\nu_{\max}$ : 3270, 3180  $\text{cm}^{-1}$  (E and Z OH), 1610, 1590  $\text{cm}^{-1}$  (C=N);  $^1\text{H}$  nmr (DMSO- $d_6$ )  $\delta$ : 2.40 (s, 3H,  $\text{CH}_3$ ), 7.2-8.2 (m, 9H, H-7,8,10,11 and  $\text{C}_6\text{H}_5$ ), 8.2-9.0 (m, 4H, H-5,8,9,12), 12.20 (broad s, 1H, NOH). A sample of this product was crystallized from DMF/ $\text{H}_2\text{O}$ , mp 282°C. Anal. Calcd for  $\text{C}_{21}\text{H}_{14}\text{N}_4\text{O}$ : C, 74.54; H, 4.17; N, 16.56. Found: C, 74.63; H, 4.24; N, 16.51.

Preparation of 2-Acylindazole Phenylhydrazones (9a,b).

Phenylhydrazones (9a,b) were prepared according to the procedure described previously for 9a,<sup>1</sup> by treatment of indazole with equimolar amounts of suitable hydrazidoyl chloride<sup>14</sup> and threefold excess of triethylamine.

Compound 9a (R = Me) was crystallized from ethanol (yield 93%), mp 140°C.<sup>1</sup>

Compound 9b (R = Ph) was purified by column chromatography on silica gel (eluent cyclohexane/dichloromethane 2:1) and recrystallized from ethanol (yield 75%), mp 136°C; ir:  $\nu_{\max}$ : 3180  $\text{cm}^{-1}$  (NH), 1640  $\text{cm}^{-1}$  (CO); uv:  $\lambda_{\max}$  nm (log  $\epsilon$ ): 244 (4.44), 298 (4.27), 377 (4.33);  $^1\text{H}$  nmr ( $\text{CDCl}_3$ )  $\delta$ : 6.7-8.6 (m, 10H, ArH), 7.72 (m, 2H, H-5 and H-6), 7.98 (m, 2H, H-4 and H-7), 8.92 (s, 1H, H-3), 13.20 (s, 1H, NH); (DMSO- $d_6$ )  $\delta$ : 8.70 (s, 1H, H-3), 11.73 (s, 1H, NH). Anal. Calcd for  $\text{C}_{21}\text{H}_{16}\text{N}_4\text{O}$ : C, 74.10; H, 4.74; N, 16.46. Found: C, 74.23; H, 4.78; N, 16.39.

Rearrangement of Phenylhydrazones (9a,b) into 1,2,4-Triazole Derivatives (10a,b)

Triazoles (10a,b) were prepared following the procedure previously described for 10a<sup>1</sup> by refluxing 9a,b (0.5 g) with conc. HCl (0.5 ml) in ethanol (20 ml) for two days. Evaporation of the solvent under reduced pressure, repeated additions of benzene to the residue and further evaporation under reduced pressure gave

**10a,b** as hydrochlorides. The corresponding free bases were obtained by column chromatography (eluent cyclohexane/ethyl acetate 4:1) of a chloroform solution of equimolar amounts of **10a** hydrochloride and triethylamine and by treating an ethanol solution of **10b** hydrochloride with aqueous ammonia (30%).

1-Phenyl-5-(o-aminophenyl)-3-acetyl-1,2,4-triazole (10a) (R = Me) : mp 139-141°C.<sup>1</sup>

1-Phenyl-5-(o-aminophenyl)-3-benzoyl-1,2,4-triazole (10b) (R = Ph) : mp 150°C (ethanol) (yield 69%); ir:  $\nu_{\max}$ : 3420, 3320  $\text{cm}^{-1}$  ( $\text{NH}_2$ ), 1669  $\text{cm}^{-1}$  (CO); uv:  $\lambda_{\max}$  nm (log  $\epsilon$ ): 260(4.24), 323(3.64);  $^1\text{H}$  nmr ( $\text{CDCl}_3$ )  $\delta$  : 4.75 (br s, 2H,  $\text{NH}_2$ ), 6.3-7.6 (m, 12H, ArH), 8.33 (m, 2H, ArH). Anal. Calcd for  $\text{C}_{21}\text{H}_{16}\text{N}_4\text{O}$ : C, 74.10; H, 4.74; N, 16.46. Found: C, 74.08; H, 4.71; N, 16.49.

Preparation of the 2-Acetyl- and 2-Benzoyl-1,2,4-triazolo[1,5-f]phenanthridines (11a,b)

Compounds **11a,b** were prepared by the procedure previously described for **11a**.<sup>1</sup>

2-Acetyl-1,2,4-triazolo[1,5-f]phenanthridine (11a) (R = Me) was crystallized from ethanol (yield 85%), mp 207-208°C.<sup>1</sup>

2-Benzoyl-1,2,4-triazolo[1,5-f]phenanthridine (11b) (R = Ph) was crystallized from ethanol (yield 94%), mp 225°C; ir:  $\nu_{\max}$ : 1620, 1595  $\text{cm}^{-1}$  (C=N), 1665  $\text{cm}^{-1}$  (CO); uv:  $\lambda_{\max}$  nm (log  $\epsilon$ ): 242s (4.73), 248 (4.83), 278s (4.36), 312s (3.79), 328 (3.61);  $^1\text{H}$  nmr ( $\text{CDCl}_3$ )  $\delta$  : 7.2-7.9 (m, 7H, H-6,7,10,11, meta- and para-H of  $\text{COC}_6\text{H}_5$ ), 8.1-8.8 (m, 6H, H-5,8,9,12 and ortho-H of  $\text{COC}_6\text{H}_5$ ). Anal. Calcd for  $\text{C}_{21}\text{H}_{13}\text{N}_3\text{O}$ : C, 78.00; H, 4.05; N, 13.00. Found: C, 78.18; H, 4.03; N, 12.89.

General Procedure for the Preparation of the Oximes (6a,b) from Hydroxylamine on Ketones (11a,b).

To a solution of hydroxylamine hydrochloride (70 mg, 1 mmol) and sodium acetate (82 mg, 1 mmol) in ethanol (30 ml) 1 mmol of **11a** or **11b** was added and the suspension heated under reflux for 8 h. After cooling, the solid was filtered off and washed with ethanol to give E oxime (**6a**) (ir and hplc) or a mixture of E and Z isomers of the oxime (**6b**) (ir and hplc).

## ACKNOWLEDGEMENTS

We thank Ministero della Pubblica Istruzione (Rome) for support.

## REFERENCES AND NOTES

1. M. Ruccia, N. Vivona, G. Cusmano, and A. M. Almerico, Heterocycles, 1978, 9, 1577.
2. G. Cusmano, G. Macaluso, N. Vivona, and M. Ruccia, Heterocycles, 1986, 24, 3181.
3. G. Cusmano and G. Macaluso, Heterocycles, 1989, 29, 339.
4. G. Cusmano, G. Macaluso, M. Gruttadauria, and S. Buscemi, Heterocycles, 1989, 29, 2149.
5. M. Ruccia, N. Vivona, and G. Cusmano, J. Heterocycl. Chem., 1975, 12, 985.
6. H. Kano and E. Yamazaki, Tetrahedron, 1964, 20, 159.
7. M. Ruccia, N. Vivona, G. Cusmano, and G. Macaluso, J. Chem. Soc., Perkin Trans. 1, 1977, 589.
8. M. Ruccia, N. Vivona, and D. Spinelli, Adv. Heterocycl. Chem., 1981, 29, 141.
9. See, e.g.: H. H. Credner, H. J. Friedrich, and G. Scheibe, Chem. Ber., 1962, 95, 1881.
10. G. M. Badger, J. H. Seidler, and B. Thomson, J. Chem. Soc., 1951, 3207.
11. S. Cusmano and T. Tiberio, Gazz. Chim. Ital., 1951, 81, 106.
12. F. Angelico and S. Cusmano, Gazz. Chim. Ital., 1936, 66, 3.
13. Comparison between rearrangement and oximation products was performed by tlc and hplc (normal and reverse-phase) analyses. The low solubility of the oximes did not allow a complete spectroscopic investigation aimed to confirm the structures of E- and Z-isomers. However, for oximes of acetyl- or benzoyl-azoles see, e.g.: ref. 7 and also: N. Vivona, G. Cusmano, and G. Macaluso, J. Chem. Soc., Perkin Trans. 1, 1977, 1616; N. Vivona, G. Macaluso, and V. Frenna, J. Chem. Soc., Perkin Trans. 1, 1983, 483; N. Vivona, S. Buscemi, V. Frenna, M. Ruccia, and M. Condo', J. Chem. Research, 1985, (S), 190, (M), 2184.
14. For the procedure adopted in the synthesis of hydrazidoyl chlorides, see: W. Dieckmann and L. Platz, Ber., 1905, 38, 2986.

Received, 19th February, 1990