

**WITTIG-HORNER REACTION OF DIMETHYL
PHTHALIDE-3-PHOSPHONATES WITH ALDEHYDES:
SYNTHESIS OF 3-YLIDENEPHTHALIDES AND
CHARACTERIZATION OF THEIR *E*- AND *Z*-ISOMERS**

Mitsuaki Watanabe,^{*a} Saori Ijichi,^b Hitoshi Morimoto,^b Kayoko Nogami,^b
and Sunao Furukawa^b


Center for Instrumental Analysis, ^a School of Pharmaceutical Sciences, ^b
Nagasaki University, 1-14 Bunkyo-machi, Nagasaki 852, Japan

Abstract- Wittig-Horner reaction of dimethyl phthalide-3-phosphonates with aldehydes in the presence of lithium bis(trimethylsilyl)amide was investigated. Using their NOE experiments, both the *E*- and *Z*-isomers of 3-ylidenephthalides were clearly characterized.

Recently, we reported a facile regioselective synthetic method for dimethyl phthalide-3-phosphonates having various substituents on the aromatic ring, based on the reaction of *N,N*-diethyl-2-formylbenzamides with dimethyl *tert*-butyldimethylsilyl phosphite.¹ In order to develop the utility of phthalide-3-phosphonates in organic synthesis, Wittig-Horner reactions² of these compounds with aldehydes were investigated. The Wittig-Horner reaction of phthalide-3-phosphonates in the presence of NaH at room temperature has already been reported,³ but characterization of the *E*- and *Z*-isomers of the synthesized 3-ylidenephthalides had not been reported except that of several thermally stable (*Z*)-3-benzylidenephthalides.^{3b} Although the 3-alkylidenephthalides may occur as the *E*- or *Z*-isomers, their configuration in natural or synthesized products had been confused for a long time.⁴ In 1980, Gijbels and

coworkers reported that natural 3-butyldenephthalide was confirmed to be the *Z*-isomer by means of ^1H and ^{13}C nmr measurements.⁴ However no nuclear Overhauser effect (NOE), which is necessary for the determination of stereochemistry, was made in their experiments. In this paper, we describe a convenient synthesis of 3-alkylidene- and 3-benzylidenephthalides using the Wittig-Horner reaction of dimethyl phthalide-3-phosphonates with aldehydes in the presence of lithium bis(trimethylsilyl)amide and the characterization of their *E*- and *Z*-isomers based on the NOE experiments.

Reaction of dimethyl phthalide-3-phosphonate (**1a**)^{1,3b} with 1.1 eq. of lithium bis(trimethylsilyl)amide in THF at -78°C for 1 h resulted in the formation of the yellow anion, which, upon treatment with 1.1 eq. of isobutyraldehyde (**2a**) at -78°C for 1 h followed by quenching with saturated NH_4Cl solution at -78°C , gave 3-isobutyldenephthalide (**3aa**) as *E/Z* mixtures in 87% yield. Lithium diisopropylamide (LDA) was also used instead of lithium bis(trimethylsilyl)amide and **3aa** was obtained in almost the same yield (85%). When the reaction mixture using LDA as a base was allowed to warm to room temperature for 1 h and then quenched, however, the yield of **3aa** was decreased to 50% yield.⁵ For the efficient synthesis of 3-ylidenephthalides, quenching of the reaction mixture at -78°C is necessary. Although we have tested NaH as a base (THF, 0°C , 6 h), the yield of **3aa** was much lower (64%). As shown in Scheme 1, various 3-ylidenephthalides (**3**), which were mixtures of *E/Z*-isomers, were synthesized in high yields starting from








Scheme 1


1a-c and **2b-f** under the reaction conditions using lithium bis(trimethylsilyl)amide as a base.

The pure samples of *E*-isomers and *Z*-isomers were easily obtained by silica gel column chromatography using benzene as the eluent for 3-alkylidenephthalides (**3aa**,^{3b,6 ab,6 ac,4 bc, cc}⁷) or dichloromethane for 3-benzylidenephthalides (**3ad, ae, af**^{3b,8}). In the cases of *E*-**3aa** and *Z*-**3aa**, their ir and mass spectra were almost identical. However, characteristic differences between *E*-**3aa** and *Z*-**3aa** were observed in their ¹H and ¹³C nmr spectra; *i. e.* the proton signal of the vinyl protons for these compounds appeared at different chemical shifts [*E*-**3aa**: δ 5.72 (δ , $J=10.26$ Hz); *Z*-**3aa**: δ 5.48 (δ , $J=9.53$ Hz)]. In the NOE experiments, the irradiation of the methine proton of *E*-**3aa** produced NOE enhancement at the signal of aromatic C4-H, but no NOE enhancement at the same proton was observed by the irradiation of the vinyl proton (Figure 1). In the NOE experiments of *Z*-**3aa**, the irradiation of the vinyl proton produced NOE enhancement at the signal of aromatic C4-H. On the other hand, no enhancement at the signals of aromatic protons was observed by the irradiation of the methine proton of *Z*-**3aa** (Figure 2).^{7,9}

In the cases of 3-benzylidenephthalides, bathochromic shifts of the uv spectra of the *Z*-isomers compared with those of the *E*-isomers were observed. In their ¹H nmr spectra; *i. e.* *E*-**3ad** and *Z*-**3ad**, the vinyl protons of *E*-**3ad** and *Z*-**3ad** appeared at δ 6.86 (s) and δ 6.35 (s), respectively. The irradiation of the vinyl proton of *E*-**3ad** gave NOEs at aromatic C2'-H and C6'-H and no NOE enhancement at the signal of aromatic C4-H was observed (Figure 3). On the other hand, the irradiation of the vinyl proton of *Z*-**3ad** caused NOEs at aromatic C4-H along with aromatic C2'-H and C6'-H (Figure 4). From the above NOE observations, the relative configurations of the 3-ylidenephthalides were clearly determined. The NOE observations and chemical shifts of the vinyl proton of *E*- and *Z*-ylidenephthalides synthesized here are summarized in Figure 5 and Table I, respectively. ¹³C Nmr data of a series of *E*- and *Z*-3-alkylidenephthalides are summarized in Table II. Using differentiation of the chemical shift of the vinyl protons, the *E/Z* ratios of each of the reaction products were calculated and the results are listed in Scheme 1. By comparison with the Perkin synthesis,⁴ which mainly gave thermally stable *Z*-isomers, the Wittig-Horner reaction of dimethyl phthalide-3-phosphonates (**1**) under the conditions described here predominantly lead to *E*-isomers due to the formation of *erythro*-intermediates.^{2,6,8} From our spectral data, the following observations allow the determination of the stereochemistry of 3-ylidenephthalides: i) in

Figure 1. ^1H Nmr Spectrum and NOE Spectra of *E*-3aaFigure 2. ^1H Nmr Spectrum and NOE Spectra of *Z*-3aaFigure 3. ^1H Nmr Spectrum and NOE Spectrum of *E*-3adFigure 4. ^1H Nmr Spectrum and NOE Spectrum of *Z*-3ad**(*E*)- and (*Z*)-3-Alkylidene-phthalides**

- 3aa:** $\text{R}^1=\text{R}^2=\text{Me}$, $\text{R}=\text{H}$
3ab: $\text{R}^1=\text{H}$, $\text{R}^2=\text{CHMe}$, $\text{R}=\text{H}$
3ac: $\text{R}^1=\text{H}$, $\text{R}^2=\text{CH}_2\text{Me}$, $\text{R}=\text{H}$
3bc: $\text{R}^1=\text{H}$, $\text{R}^2=\text{CH}_2\text{Me}$, $\text{R}=\text{OMe}$

(*E*)- and (*Z*)-3-Benzylidene-phthalides


- 3ad:** $\text{R}^1=\text{R}^2=\text{R}^3=\text{OMe}$
3ae: $\text{R}^1=\text{R}^2=\text{OMe}$, $\text{R}^3=\text{H}$
3af: $\text{R}^1=\text{R}^2=\text{OCH}_2\text{O}$, $\text{R}^3=\text{H}$

Figure 5. NOE Correlations of (*E*)- and (*Z*)-3-Ylidene-phthalides


Table I. Chemical Shift of Vinyl Protons of (*E*)- and (*Z*)-3-Ylidenephthalides
(400 MHz, δ from TMS in CDCl_3)

	3aa	3ab	3ac	3bc	3cc	3ad	3ae	3af
<i>E</i> -	5.72 (d)	5.87 (t)	5.83 (t)	5.78 (t)	5.67 (t)	6.86 (s)	6.87 (s)	6.82 (s)
<i>Z</i> -	5.48 (d)	5.67 (t)	5.64 (t)	5.59 (t)	5.47 (t)	6.35 (s)	6.36 (s)	6.35 (s)
Δ	0.24	0.20	0.19	0.19	0.20	0.51	0.51	0.47

Table II. ^{13}C Nmr Data for (*E*)- and (*Z*)-3-Alkylidenephthalides



E-3



Z-3

9 10
3aa: R=CHMe₂
9 10 11
3ab: R=CH₂CHMe₂
9 10 11
3ac: R=CH₂CH₂Me

Carbons	<i>E</i> -3aa	<i>E</i> -3ab	<i>E</i> -3ac	<i>Z</i> -3aa	<i>Z</i> -3ab	<i>Z</i> -3ac
1	166.97	166.89	166.67	167.13	167.20	167.13
3a	138.13	138.32	138.03	139.80	139.61	139.57
3	144.47	146.07	145.58	144.05	146.19	145.80
4	123.11	123.18	122.89	119.64	119.68	119.68
5	134.39	134.28	134.10	134.16	134.24	134.24
6	129.57	129.57	129.95	129.33	129.37	129.33
7	125.52	125.44	125.15	125.25	125.25	125.13
7a	126.26	126.26	125.88	124.51	124.55	124.43
8	121.12	113.03	113.62	116.33	108.39	109.41
9	26.00	34.99	27.73	25.92	34.76	27.83
10	23.12	29.19	22.63	22.81	28.72	22.54
11		22.34	13.52		22.38	13.78

Assignments were made by heteronuclear multiple bond connectivity.

the ^1H nmr spectra, the vinyl protons of *E*-isomers of 3-ylidenephthalides appear further downfield than those of *Z*-isomers (Table I); ii) in the ^{13}C nmr spectra of 3-alkylidenephthalides, the carbon signals of the aromatic C4 are remarkably different for *E*- and *Z*-isomers; in *E*-isomers, signals appear further downfield compared with those in *Z*-isomers (Table II).⁴

Among the (*Z*)-3-alkylidenephthalides synthesized here, both 3-isobutylidenephthalide (3aa) and 3-isovalidenephthalide (3ab) were isolated as components of the flavor of celery plants,¹⁰ and 3-butylidenephthalide (3ac) was isolated from umbelliferous plants.^{4,11} Moreover, (*Z*)-3-butylidene-6,7-dihydroxyphthalide isolated from *Ligusticum wallichii* Franch¹² was recently synthesized by Ogawa and coworkers through its dimethyl derivative (3cc: *E/Z*=1/1.8).⁷

In conclusion, we have shown that the Wittig-Horner reaction of dimethyl phthalide-3-phosphonate with aldehydes in the presence of lithium bis(trimethylsilyl)amide provided an efficient synthetic method for the preparation of various (*E*)- and (*Z*)-3-ylidenephthalide derivatives. Each assignment of configuration of the *E*- and *Z*-isomers in 3-alkylidene- and 3-benzylidenephthalides was clarified by their NOE experiments.

EXPERIMENTAL

General. Melting points were determined with a Yanagimoto micromelting point apparatus and were uncorrected. The IR spectra were measured with a JASCO 810 spectrophotometer. The UV spectra were recorded in 95% ethanol on a Hitachi 323 spectrophotometer. The NMR spectra were obtained on a JEOL GX-400 spectrometer operating at 399.65 MHz for ^1H and 100.40 MHz for ^{13}C nuclei using CDCl_3 as a solvent and tetramethylsilane as an internal reference. The MS were determined on a JEOL DX-303 mass spectrometer. Elemental analyses were performed at the Microanalytical Laboratory of the Center for Instrumental Analysis in Nagasaki University. Conventional column chromatography was carried out on a column of Kieselgel 60 (230–400 mesh).

General Procedure for the Reaction of Dimethyl Phthalide-3-phosphonates (1) with Aldehydes (2). A solution of lithium bis(trimethylsilyl)amide (3.3 mmol, 1.0 M solution of hexane) was injected into a stirred solution of $1^{1,3b}$ (3 mmol) in THF (50 ml) at -78°C under an argon atmosphere. The mixture was stirred at -78°C for 1 h and then a solution of aldehydes (2; 3.3 mmol) in THF (20 ml) was injected into the yellow solution at -78°C . Stirring was continued for an additional 1 h at -78°C . The reaction was quenched with saturated NH_4Cl solution and 10% HCl solution at -78°C . THF was removed under reduced pressure. The residue was extracted with CH_2Cl_2 and the CH_2Cl_2 layer was dried over Na_2SO_4 , and then evaporated to give 3-ylidenephthalides (3). The analytically pure samples of *E*- and *Z*-isomers of 3 were obtained by silica gel column chromatography. For the purification of 3-alkylidenephthalides, benzene was used as eluent and the *Z*-isomers eluted faster than the *E*-isomers. For a purification of 3-benzylidenephthalides, CH_2Cl_2 was used as eluent and the *E*-isomers eluted faster than the *Z*-isomers except for **3af** which was

reversed.

(E)-3-Isobutylidenephthalide (E-3aa): mp 56-57°C (ether/hexane). Ms *m/z*: 188 (M^+ , 18), 173 (100), 155 (7), 104 (10). Ir (KBr) vco: 1780 cm^{-1} . Uv nm (log ϵ): 210 (4.24), 215 (4.22), 235 (4.21), 250 (s)(4.21), 260 (4.26), 270 (s)(4.15), 308 (3.82). ^1H Nmr (400 MHz) δ : 1.21 (6H, d, $J=7.0$ Hz), 3.13-3.22 (1H, m), 5.72 (1H, d, $J=10.3$ Hz), 7.54 (1H, dd, $J=7.7, 0.7$ Hz), 7.72 (1H, dd, $J=8.1, 1.1$ Hz), 7.83 (1H, d, $J=8.1$ Hz), 7.92 (1H, d, $J=7.7$ Hz). Anal. Calcd for $\text{C}_{12}\text{H}_{12}\text{O}_2$: C, 76.57; H, 6.43. Found: C, 76.31; H, 6.46.

(Z)-3-Isobutylidenephthalide (Z-3aa): an oil. Ms *m/z*: 188 (M^+ , 33), 173 (100), 155 (9), 103 (79). Ir (neat) vco: 1780 cm^{-1} . Uv nm (log ϵ): 205 (4.11), 215 (s)(4.07), 233 (s)(3.99), 258 (s)(3.81), 270 (s)(3.65), 308 (3.26). ^1H Nmr (400 MHz) δ : 1.15 (6H, d, $J=7.0$ Hz), 3.09-3.18 (1H, m), 5.38 (1H, d, $J=9.5$ Hz), 7.50 (1H, dd, $J=8.1, 1.5$ Hz), 7.61-7.69 (2H, m), 7.89 (1H, d, $J=7.7$ Hz). Anal. Calcd for $\text{C}_{12}\text{H}_{12}\text{O}_2$: C, 76.57; H, 6.43. Found: C, 76.23; H, 6.74.

(E)-3-Isovalidenephthalide (E-3ab): an oil. Ms *m/z*: 202 (M^+ , 24), 159 (100), 148 (26), 132 (32), 104 (68). Ir (neat) vco: 1770 cm^{-1} . Uv nm (log ϵ): 210 (4.45), 215 (s)(4.43), 235 (s)(4.42), 260 (4.49), 270 (s)(4.37), 308 (4.06). ^1H Nmr (400 MHz) δ : 1.03 (6H, d, $J=6.6$ Hz), 1.84-1.91 (1H, m), 2.45 (2H, dd, $J=8.4, 6.6$ Hz), 5.87 (1H, t, $J=8.4$ Hz), 7.54 (1H, t, $J=7.3$ Hz), 7.72 (1H, dt, $J=7.7, 1.1$ Hz), 7.85 (1H, d, $J=7.7$ Hz), 7.91 (1H, d, $J=7.7$ Hz). Anal. Calcd for $\text{C}_{13}\text{H}_{14}\text{O}_2$: C, 77.20; H, 6.98. Found: C, 77.49; H, 7.27.

(Z)-3-Isovalidenephthalide (Z-3ab): an oil (this compound is gradually solidified in a refrigerator; mp 108-110°C). Ms *m/z*: 202 (M^+ , 12), 159 (48), 148 (82), 132 (9), 104 (100). Ir (neat) vco: 1780 cm^{-1} . Uv nm (log ϵ): 215 (s)(4.07), 232.5 (4.11), 257.5 (4.02), 270 (s)(3.89), 310 (3.51). ^1H Nmr (400 MHz) δ : 0.984 (6H, d, $J=7.0$ Hz), 1.77-1.88 (1H, m), 2.37 (2H, dd, $J=8.1, 7.0$ Hz), 5.66 (1H, t, $J=8.1$ Hz), 7.49-7.52 (1H, m), 7.66-7.68 (2H, m), 7.88 (1H, d, $J=7.7$ Hz). Anal. Calcd for $\text{C}_{13}\text{H}_{14}\text{O}_2$: C, 77.20; H, 6.98. Found: C, 77.31; H, 7.16.

(E)-3-Butylidenephthalide (E-3ac): an oil (lit.,^{11b} bp 125-135°C/2 mmHg). Ms *m/z*: 188 (M^+ , 36), 159 (100), 146 (29), 132 (19), 104 (33). Ir (neat) vco: 1770 cm^{-1} . Uv nm (log ϵ): 208 (4.15), 214 (s)(4.14), 235 (4.13), 258 (4.18), 268 (s)(4.06), 308 (3.74). ^1H Nmr (400 MHz) δ : 1.03 (3H, t, $J=7.3$ Hz), 1.62 (2H, sext, $J=7.3$ Hz), 2.53 (1H, q, $J=7.3$ Hz), 5.83 (1H, t, $J=8.1$ Hz), 7.53 (1H, dt, $J=7.0, 0.7$ Hz), 7.71 (1H, dt,

$J=7.3, 1.1$ Hz), 7.82 (1H, d, $J=7.7$ Hz), 7.89 (1H, dd, $J=7.7, 1.1$ Hz). *Anal.* Calcd for $C_{12}H_{12}O_2$: C, 76.57; H, 6.43. Found: C, 76.42; H, 6.50.

(Z)-3-Butylidenephthalide (Z-3ac): an oil. *Ms m/z*: 188 (M^+ , 19), 159 (52), 146 (63), 132 (42), 104 (100). *Ir* (neat) ν_{co} : 1780 cm^{-1} . *Uv nm* (log ϵ): 210 (s)(3.93), 215 (3.95), 234 (4.05), 256 (4.05), 268 (s)(3.93), 315 (3.61). $^1\text{H Nmr}$ (400 MHz) δ : 0.98 (3H, t, $J=7.3$ Hz), 1.55 (2H, sext, $J=7.3$ Hz), 2.44 (2H, q, $J=7.7$ Hz), 5.64 (1H, t, $J=7.7$ Hz), 7.46-7.50 (1H, m), 7.62-7.68 (2H, m), 7.84-7.86 (1H, m). *Anal.* Calcd for $C_{12}H_{12}O_2$: C, 76.57; H, 6.43. Found: C, 76.70; H, 6.56.

(E)-5-Methoxy-3-butylidenephthalide (E-3bc): an oil (this compound is gradually solidified in a refrigerator; mp 42-43°C). *Ms m/z*: 218 (M^+ , 34), 189 (100), 176 (72), 161 (15), 134 (33), 106 (20). *Ir* (neat) ν_{co} : 1770 cm^{-1} . *Uv nm* (log ϵ): 226 (4.17), 257 (4.41), 305 (s)(3.56), 318 (3.34). $^1\text{H Nmr}$ (400 MHz) δ : 1.03 (3H, t, $J=7.3$ Hz), 1.62 (2H, sext, $J=7.3$ Hz), 2.48 (2H, q, $J=7.7$ Hz), 3.93 (3H, s), 5.78 (1H, t, $J=8.1$ Hz), 7.04 (1H, dd, $J=8.4, 2.2$ Hz), 7.18 (1H, d, $J=2.2$ Hz), 7.77 (1H, d, $J=8.4$ Hz). $^{13}\text{C nmr}$ δ : 13.66, 22.73, 27.83, 55.77, 107.30, 113.57, 116.49, 118.55, 126.69, 140.31, 145.45, 164.64, 166.46. *Anal.* Calcd for $C_{13}H_{14}O_3$: C, 71.54; H, 6.47. Found: C, 71.62; H, 6.55.

(Z)-5-Methoxy-3-butylidenephthalide (Z-3bc): an oil (this compound is gradually solidified in a refrigerator; mp 50-51°C). *Ms m/z*: 218 (M^+ , 35), 189 (100), 176 (78), 161 (16), 134 (10), 116 (4), 106 (7). *Ir* (neat) ν_{co} : 1780 cm^{-1} . *Uv nm* (log ϵ): 220 (s)(4.06), 225 (4.13), 255 (4.50), 275 (s)(4.20), 303 (s)(3.67), 315 (3.47). $^1\text{H Nmr}$ (400 MHz) δ : 0.98 (3H, t, $J=7.3$ Hz), 1.55 (2H, sext, $J=7.3$ Hz), 2.43 (2H, q, $J=7.7$ Hz), 3.91 (3H, s), 5.59 (1H, t, $J=7.7$ Hz), 7.01 (1H, dd, $J=9.2, 2.2$ Hz), 7.02 (1H, d, $J=2.2$ Hz), 7.74 (1H, d, $J=9.2$ Hz). $^{13}\text{C Nmr}$ δ : 13.82, 22.54, 27.79, 55.89, 102.63, 109.17, 117.11, 117.77, 126.65, 142.18, 145.68, 164.98, 166.89. *Anal.* Calcd for $C_{13}H_{14}O_3$: C, 71.54; H, 6.47. Found: C, 71.84; H, 6.68.

(E)-6,7-Dimethoxy-3-butylidenephthalide (E-3cc): an oil (lit.,⁷ an oil). *Ms m/z*: 248 (M^+ , 43), 219 (100), 206 (9). *Ir* (neat) ν_{co} : 1780 cm^{-1} . *Uv nm* (log ϵ): 224 (4.29), 238 (4.26), 266 (4.15), 278 (s)(4.05), 338 (3.90). $^1\text{H Nmr}$ (400 MHz) δ : 1.02 (3H, t, $J=7.3$ Hz), 1.60 (2H, sext, $J=7.3$ Hz), 2.46 (2H, q, $J=7.7$ Hz), 3.95 (3H, s), 4.10 (3H, s), 5.67 (1H, t, $J=8.1$ Hz), 7.28 (1H, d, $J=8.4$ Hz), 7.45 (1H, d, $J=8.4$ Hz). $^{13}\text{C Nmr}$ δ : 13.78, 22.89, 27.95, 56.86, 62.27, 111.16, 118.28, 119.53, 131.94, 145.02, 147.98, 153.08, 164.56. The above *ms*, *ir*, *uv*, ^1H , and ^{13}C *nmr* data agreed well with those of an authentic sample of *E*-

3cc.⁷

(Z)-6,7-Dimethoxy-3-butylidenephthalide (Z-3cc): an oil (lit.,⁷ an amorphous powder). Ms *m/z*: 248 (M^+ , 44), 219 (100), 206 (10). Ir (neat) ν : 1770 cm^{-1} . Uv nm (log ϵ): 225 (4.06), 240 (4.07), 268 (3.95), 283 (s)(3.75), 340 (3.72). ^1H Nmr (400 MHz) δ : 0.97 (3H, t, $J=7.3$ Hz), 1.53 (2H, sext, $J=7.3$ Hz), 2.39 (2H, q, $J=7.7$ Hz), 3.93 (3H, s), 4.11 (3H, s), 5.47 (1H, t, $J=7.7$ Hz), 7.25 (1H, d, $J=8.4$ Hz), 7.29 (1H, d, $J=8.4$ Hz). ^{13}C Nmr δ : 13.82, 22.65, 27.75, 56.94, 62.20, 106.92, 114.58, 116.65, 120.03, 133.23, 145.10, 147.74, 152.80, 164.75. The above ms, ir, uv, ^1H , and ^{13}C nmr data agreed well with those of an authentic sample of Z-3cc.⁷

(E)-3-(3',4',5'-Trimethoxy)benzylidenephthalide (E-3ad): mp 148°C (CH_2Cl_2 /ether). Ms *m/z*: 312 (M^+ , 100), 297 (65), 269 (13). Ir (KBr) ν : 1780 cm^{-1} . Uv nm (log ϵ): 209 (4.50), 233 (s)(4.20), 253 (s)(4.13), 333 (4.08). ^1H Nmr (400 MHz) δ : 3.86 (6H, s), 3.93 (3H, s), 6.70 (1H, s), 6.71 (1H, s), 6.86 (1H, s), 7.72-7.57 (2H, m), 7.61-7.64 (1H, m), 7.91-7.96 (1H, m). Anal. Calcd for $\text{C}_{18}\text{H}_{16}\text{O}_5$: C, 69.22; H, 5.16. Found: C, 69.15; H, 5.14.

(Z)-3-(3',4',5'-Trimethoxy)benzylidenephthalide (Z-3ad): mp 162°C (ether/pentane). Ms *m/z*: 312 (M^+ , 100), 297 (59), 269 (10). Ir (KBr) ν : 1780 cm^{-1} . Uv nm (log ϵ): 210 (4.38), 242 (4.17), 305 (s)(4.17), 354 (4.35). ^1H Nmr (400 MHz) δ : 3.90 (3H, s), 3.95 (6H, s), 6.35 (1H, s), 7.11 (2H, s), 7.52-7.57 (1H, m), 7.70-7.79 (2H, m), 7.95 (1H, d, $J=7.7$ Hz). Anal. Calcd for $\text{C}_{18}\text{H}_{16}\text{O}_5$: C, 69.22; H, 5.16. Found: C, 68.90; H, 5.18.

(E)-3-(3',4'-Dimethoxy)benzylidenephthalide (E-3ae): mp 154-155°C (CH_2Cl_2 /hexane). Ms *m/z*: 282 (M^+ , 100), 267 (39), 239 (12), 221 (17). Ir (KBr) ν : 1775 cm^{-1} . Uv nm (log ϵ): 203 (4.53), 230 (s)(4.23), 255 (4.14), 345 (4.12). ^1H Nmr (400 MHz) δ : 3.89 (3H, s), 3.96 (3H, s), 6.87 (1H, s), 6.95 (1H, d, $J=8.1$ Hz), 6.97 (1H, d, $J=1.8$ Hz), 7.05-7.08 (1H, m), 7.50-7.54 (2H, m), 7.59-7.63 (1H, m), 7.90-7.94 (1H, m). Anal. Calcd for $\text{C}_{17}\text{H}_{14}\text{O}_4$: C, 72.33; H, 5.00. Found: C, 71.97; H, 5.00.

(Z)-3-(3',4'-Dimethoxy)benzylidenephthalide (Z-3ae): mp 128-129°C (ethyl acetate/ether). Ms *m/z*: 282 (M^+ , 100), 267 (22), 239 (9), 221 (11). Ir (KBr) ν : 1780 cm^{-1} . Uv nm (log ϵ): 201 (4.33), 235 (s)(4.32), 232 (4.18), 260 (s)(4.10), 303 (4.15), 362 (4.38). ^1H Nmr (400 MHz) δ : 3.92 (3H, s), 3.97 (3H, s), 6.36 (1H, s), 6.89 (1H, d, $J=8.4$ Hz), 7.35 (1H, dd, $J=8.4, 1.8$ Hz), 7.49-7.53 (2H, m), 7.67-7.74 (2H, m), 7.92 (1H, d,

$J=8.1$ Hz). *Anal.* Calcd for $C_{17}H_{14}O_4$: C, 72.33; H, 5.00. Found: C, 72.25; H, 5.10.

(*E*)-3-(3',4'-Methylenedioxy)benzylidenephthalide (*E*-3af): mp 125-126°C (CH_2Cl_2 /hexane) (lit.,⁸ mp 139-140°C). *Ms m/z*: 266 (M^+ , 100), 160 (10), 152 (23). Ir (KBr) ν_{CO} : 1770 cm^{-1} . Uv nm (log ϵ): 203 (4.56), 232 (s)(4.23), 255 (4.25), 348 (4.07). 1H Nmr (400 MHz) δ : 6.05 (2H, s), 6.82 (1H, s), 6.89 (1H, d, $J=8.1$ Hz), 6.93-6.98 (2H, m), 7.31-7.57 (2H, m), 7.61-7.64 (1H, m), 7.91-7.95 (1H, m). *Anal.* Calcd for $C_{16}H_{10}O_4$: C, 72.18; H, 3.79. Found: C, 71.77; H, 3.99.

(*Z*)-3-(3',4'-Methylenedioxy)benzylidenephthalide (*Z*-3af): mp 202-203°C (CH_2Cl_2 /hexane) (lit.,^{3b} mp 201-203°C; lit.,⁸ mp 202-204°C). *Ms m/z*: 266 (M^+ , 100), 160 (7), 152 (17). Ir (KBr) ν_{CO} : 1760 cm^{-1} . Uv nm (log ϵ): 201 (4.43), 220 (s)(4.36), 240 (s)(4.19), 255 (s)(4.08), 300 (4.08), 363 (4.36). 1H Nmr (400 MHz) δ : 6.01 (2H, s), 6.35 (1H, s), 6.84 (1H, d, $J=8.1$ Hz), 7.22 (1H, dd, $J=8.1, 1.8$ Hz), 7.51-7.55 (2H, m), 7.68-7.75 (2H, m), 7.93 (1H, d, $J=7.7$ Hz). *Anal.* Calcd for $C_{16}H_{10}O_4$: C, 72.18; H, 3.79. Found: C, 72.22; H, 3.95.

ACKNOWLEDGEMENTS

We are grateful to Drs T. Wakamatsu and Y. Ogawa in Tsumura & Co., for providing the spectra of *E*- and *Z*-3cc, and for valuable suggestions. We also thank Dr. I. Kouno in Nagasaki University for helpful discussions. We are indebted to the staff of the Center for Instrumental Analysis in Nagasaki University for ir, uv, ms, and nmr measurements and elemental analysis.

REFERENCES AND NOTES

1. M. Watanabe and S. Furukawa, *SYNLETT*, 1991, 481; M. Watanabe, S. Ijichi, and S. Furukawa, *Synthesis*, 1992, in press.
2. J. Boutagy and R. Thomas, *Chem. Rev.*, 1974, 74, 87 and references cited therein.
3. a) A. Yamaguchi and A. Okazaki, *Nippon Kagaku Kaishi*, 1973, 110; b) E. Napolitano, G. Spinelli, R. Fiaschi, and A. Marsili, *Synthesis*, 1985, 38.
4. M. J. M. Gijbels, J. J. C. Scheffer, and A. Baerheim Svendsen, *Planta Medica*, 1980, suppl., 41 and references cited therein.
5. It has been reported that 3-ylidenephthalides are converted to 2-substituted 1,3-indanediones under basic

- conditions, see: H. van der Goot, J. Ch. Eriks, M. Paula, J. van Rhijn-van der Schaar, O. P. Zuiderveld, and W. Th. Naura, *Eur. J. Med. Chem., Chimica Therapeutica*, **1978**, *13*, 425; Generally, 3-ylidenephthalides are unstable under basic conditions or uv irradiation.⁴
6. D. W. Knight and G. Pattenden, *J. Chem. Soc., Perkin Trans. I*, **1975**, 635.
 7. Y. Ogawa, K. Hosaka, M. Chin, and H. Mitsuhashi, *Heterocycles*, **1991**, *32*, 1737.
 8. R. K. Howe, *J. Org. Chem.*, **1973**, *38*, 4164.
 9. The NOE experiments of some 3-butyldenephthalides have been carried out by Ogawa *et al.*; private communication.
 10. H. J. Gold and C. W. Wilson, III, *J. Org. Chem.*, **1963**, *28*, 985.
 11. a) H. Mitsuhashi, T. Muramatsu, U. Nagai, T. Nakano, and K. Ueno, *Chem. Pharm. Bull.*, **1963**, *11*, 1317; b) T. Yamagishi and H. Kaneshima, *Yakugaku Zasshi*, **1977**, *97*, 237.
 12. M. P. Baronnet, M. Kaouadji, and A. M. Mariotte, *Planta Medica*, **1984**, *50*, 105; M. Kaouadji and C. Pouget, *J. Nat. Prod.*, **1986**, *49*, 184.

Received, 3rd September, 1992