

**SYNTHESIS OF THE FIRST EXAMPLES OF 1-BENZOSILEPINE,
1-BENZOGERMEPINE, AND 1-BENZARSEPINE RING SYSTEMS**

Jyoji Kurita, Shinichi Shiratori, Shuji Yasuike, and Takashi Tsuchiya*

Faculty of Pharmaceutical Sciences, Hokuriku University,
Kanagawa-machi, Kanazawa 920-11, Japan


Abstract - The first synthesis of 1,1-dimethyl-1-benzosilepine, 1,1-dimethyl-1-benzogermepine, and 1-phenyl-1-benzarsepines was achieved by flash vacuum pyrolysis of the dihydrocyclobut[*b*]benzoheteroles (**4,7**), prepared from the corresponding 1-benzoheteroles via three steps.

There has been considerable current interest in the synthesis of fully unsaturated seven-membered heterocyclic rings (heteroepines) containing an element other than nitrogen, oxygen, or sulfur and a variety of monocyclic and fused heteroepines have been prepared. With regard to benzoheteroepines, 3-benzoheteroepines containing B,¹ Si,² Sn,³ P,⁴ Sb,⁵ or Te⁶ are known, however, such 1-benzoheteroepines were not reported prior to several years ago. Therefore, we were interested in the synthesis of 1-benzoheteroepines containing Group 14, 15, and 16 heavier elements and have already reported the syntheses of 1-benzophosphepines,⁷ 1-benzoselenepines⁸ and 1-benzotellurepines.⁸ We report here on the synthesis of the title 1-benzoheteroepines.

The 1-benzosilole (**1a**)⁹ and 1-benzogermole (**1b**)¹⁰ were irradiated with methyl acrylate to give the adducts (**2**), which were obtained as mixtures of two stereoisomers, but used in the following reactions without separation. The adducts (**2**) were hydrolysed and then oxidatively decarboxylated by treatment with lead tetraacetate in the presence of cupric acetate and pyridine giving rise to the starting tricyclic compounds (**4**)¹¹

in *ca.* 20% yields from **1**, via the acids (**3**). Although when the tricyclic compounds (**4**) were heated in diphenyl ether at 180 °C for 20 h, no reaction occurred. However, flash vacuum pyrolysis (f.v.p.) of **4** resulted in valence isomerization with ring opening to give the desired 1-benzosilepine (**5a**) and 1-benzogermepine (**5b**),¹² along with naphthalene, in the yields shown in Table 1. The heteroepines (**5**) easily reverted back to the starting tricyclic compounds (**4**) in high yields on irradiation in benzene.

Next, f.v.p. of the dihydrocyclobut[*b*]arsindole oxide (**7**),¹¹ prepared from 1-phenylarsindole¹⁰ by similar procedures, also afforded the 1-benzarsepine 1-oxide (**8**)¹² and naphthalene in the yields shown in Table 1. Treatment of **8** with trichlorosilane gave the deoxygenated arsepine (**9**; 75% yield), which reverted to the oxide (**8**) on treatment with *m*-chloroperbenzoic acid (*m*-CPBA) in 80% yield.



Scheme 1

Table 1 F. V. P. of **4** and **7**

Compd.	Temp. (°C)	Press. (Torr)	Yield (%)		
			5 or 8	6	Recovery
4a	450	1.9×10^{-5}	59	tarce	40
	500	1.9×10^{-5}	84	12	2
	550	3.0×10^{-5}	77	21	...
4b	450	5.5×10^{-5}	33	15	50
	500	4.1×10^{-5}	47	42	10
	550	5.8×10^{-5}	15	83	...
7	500	1.7×10^{-5}	40	31	22
	530	2.2×10^{-5}	39	42	8
	550	1.1×10^{-5}	27	49	...

Both 1- and 3-benzoheteroepines containing heavier elements (S,¹³ Se,⁸ Te,⁸ P^{4,7}) are known to be thermolabile and gradually decomposed to naphthalene by extrusion of the hetero elements. However, the silepine (**5a**), germepine (**5b**) and arsepine oxide (**8**) are thermally stable and remained largely unchanged even when heated in solvents at 200 °C for 24 h, and undergo thermal decomposition to naphthalene only by further f.v.p. at 500-550 °C. The results shown in Table 1 indicate that the silepine (**5a**) is somewhat less thermolabile than the other heteroepines (**5b**) and (**8**). In contrast to the oxide (**8**), the deoxygenated arsepine (**9**) is extremely thermolabile and gradually decomposed even at room temperature. The half-life of **9** estimated by the ¹H-nmr spectral analysis in toluene at 80 °C is about 10 min. This thermal behaviour is similar to those of 1-phenyl-1-benzophosphepine ($t_{1/2} = 90$ min at 80 °C),⁷ 3-phenyl-3-benzophosphepine ($t_{1/2} = 120$ min at 80 °C),⁴ and 1-benzothiepine ($t_{1/2} = 58$ min at 47 °C),¹³ whose oxides are also thermally stable. Therefore, the oxide group in **7** is essential for the present thermal isomerization to benzarsepines, in fact, f.v.p. of the dihydrocyclobut[*b*]arsindole having no oxide group gave only naphthalene, derived from the initially formed benzarespine (**9**).

Further studies on the reactions of these new heterocyclic ring systems are in progress.

REFERENCES AND NOTES

1. A. J. Leusink, W. Drenth, J. G. Noltes, and G. J. M. van der Kerk, *Tetrahedron Lett.*, 1967, 1263; G. Axelrad and D. Halpern, *J. Chem. Soc., Chem. Commun.*, 1971, 291.
2. T. J. Barton, W. E. Volz, and J. L. Johnson, *J. Org. Chem.*, 1971, **36**, 3365; L. Birkofer and H. Haddad, *Chem. Ber.*, 1977, **110**, 3314.
3. A. J. Leusink, H. A. Budding, and J. G. Noltes, *J. Organomet. Chem.*, 1970, **24**, 375.
4. G. Markl and W. Burgen, *Tetrahedron Lett.*, 1983, **24**, 2545.
5. A. J. Ashe III, L. Goossen, J. W. Kampf, and H. Konishi, *Angew. Chem., Int. Ed. Engl.*, 1992, **31**, 1642.
6. H. Sashida, H. Kurahashi, and T. Tsuchiya, *J. Chem. Soc., Chem. Commun.*, 1991, 802.
7. J. Kurita, S. Shiratori, S. Yasuike, and T. Tsuchiya, *J. Chem. Soc., Chem. Commun.*, 1991, 1227.
8. H. Sashida, K. Ito, and T. Tsuchiya, *J. Chem. Soc., Chem. Commun.*, 1993, submitted.
9. T. J. Barton and B. L. Groh, *Organometallics*, 1985, **4**, 575; T. J. Barton and M. Juvet, *Tetrahedron Lett.*, 1975, **45**, 3893.
10. J. Kurita, M. Ishii, S. Yasuike, and T. Tsuchiya, *J. Chem. Soc., Chem. Commun.*, 1993, submitted.
11. Satisfactory elemental analyses and spectral data were obtained for all new compounds reported. **4a**: Oil, bp₂₅ 112-113 °C; ¹H-nmr (CDCl₃) δ 0.33 and 0.40 (each 3H, s), 2.84 (1H, d, J=3.7 Hz), 4.52 (1H, d, J=3.7 Hz), 6.29 (2H, s); **4b**: oil; **7**: mp 196-198 °C.
12. **5a**: Oil; ¹H-nmr (CDCl₃) δ 0.29 (6H, s, SiMe₂), 5.88 (1H, d, 2-H), 6.30 (1H, dd, 4-H), 6.79 (1H, dd, 3-H), 6.91 (1H, d, 5-H), J_{2,3}= 14.3, J_{3,4}= 5.9, J_{4,5}= 13.2 Hz; **5b**: oil; ¹H-nmr δ 0.38 (6H, s, GeMe₂), 5.97 (1H, d, 2-H), 6.26 (1H, dd, 4-H), 6.71 (1H, dd, 3-H), 6.81 (1H, d, 5-H), J_{2,3}= 13.2, J_{3,4}= 5.8, J_{4,5}= 13.2 Hz; **8**: mp 158-159 °C; **9**: oil, ¹H-nmr δ 6.17 (1H, d, 2-H), 6.46 (1H, dd, 4-H), 6.66 (1H, dd, 3-H), 7.02 (1H, d, 5-H), J_{2,3}= 11.3, J_{3,4}= 5.3, J_{4,5}= 12.5 Hz. Very recently, we have found a different synthetic route to the 1-benzoarsepine (**9**) from (Z)-o, β-dibromostyrene *via* five steps.
13. I. Murata and T. Tatsuoka, *Tetrahedron Lett.*, 1975, 2697.

Received, 29th July, 1993