


NOVEL CYCLIZATION OF UNSATURATED ALCOHOLS BY
PHENYL SELENOCYANATE IN THE PRESENCE OF
COPPER BIS(TRIFLUOROMETHANESULFONATE)

Hirofumi Inoue¹ and Shizuaki Murata*

Graduate School of Human Informatics, Nagoya University
Chikusa, Nagoya, 464-01 Japan













Abstract—Intramolecular oxyselenenylation of unsaturated alcohols by phenyl selenocyanate with copper bis(trifluoromethanesulfonate) proceeds stereoselectively to give cyclic ethers.

Organic selenocyanates, which have been known for more than 100 years, are very stable compounds and can be widely prepared from the corresponding organohalogen compounds or aromatic amines by simple substitution reactions using potassium selenocyanate and so on.² In addition, it is very easy to convert the $-\text{SeCN}$ group into other selenium functional groups.^{2,3} Therefore, aryl selenocyanates have been generally employed as fundamental precursors for various organoselenium nucleophiles such as ArSe^- .²⁻⁴ In the contrast, there were only a few examples of aryl selenocyanates which were activated by Lewis acids or metal salts (Cu^{2+} or Ni^{2+}) as an electrophilic organoselenium reagent in organic synthesis.⁵⁻⁹ In this communication, we describe about preparation of a new electrophilic reagent from phenyl selenocyanate (1) and copper bis(trifluoromethanesulfonate) ($\text{Cu}(\text{OTf})_2$)¹⁰ and its application to regio- and stereoselective cyclization of unsaturated alcohols.



3a, 4a: $\text{R} = \text{CH}_3$; **3b, 4b:** $\text{R} = \text{OCH}_3$; **3c, 4c:** $\text{R} = \text{OCOC}_6\text{H}_4\text{CF}_3\text{-}p$

Table 1. Cyclization of Unsaturated Alcohols (3) by PhSeCN–Cu(OTf)₂ (2).

unsaturated alcohol (3)	cyclic ether (4) (yield/%)
	 (95) ^b
	 (78) ^c
	 (78) ^d
	 (73)
	 (88)
	 (85) ^e

^aIsolated yield after silica gel column chromatography. ^b*trans/cis* = 96:4. ^c*trans/cis* = 13:87. ^d*trans/cis* = 23:77. ^e*trans/cis* = 89:11.

The reagent (PhSeCN–Cu(OTf)₂, **2**) was obtained from a 1:1 mixture of **1** and Cu(OTf)₂ in toluene (at 50 °C) or dichloromethane (under reflux) as a yellowish brown suspension.

Reactions of 4-methyl-5-hexen-1-ol (**3a**) and the 4-methoxy and 4-*p*-trifluoromethylbenzoyloxy analogs (**3b** and **3c**, respectively) with **2** proceeded at 25 °C to give *trans*-2-phenylselenomethyl-3-methyltetrahydropyran (*trans*-**4a**, 95% yield; 96:4 selectivity), *cis*-**4b** (78%; *trans/cis* = 13:87), and *cis*-**4c** (78%; *trans/cis* = 23:77) as major products, respectively. The results of cyclization of unsaturated alcohols (**3a** – **3f**) by **2** are summarized in Table 1. There, *exo*-cyclized 5- and 6-membered ring ethers (**4a** – **4f**) were regioselectively obtained in high yields. Existence of either *endo*-cyclized products or nitriles resulting from the cyanoselenenylation⁶ was not detected in the crude reaction mixtures. The *exo* selectivity and stereoselectivities depending on substituents at the allylic position were almost same as reactions which were carried out by benzeneselenenyl trifluoromethanesulfonate (PhSeOTf),^{11–15} and yields of **4** were better than those in reactions using PhSeOTf.

It was known that organic selenocyanate (RSeCN) was converted into selenenyl halide (RSeX) by a metal salt (CuX₂ or AgX),¹⁶ and formation of PhSeOTf by the same mechanism was likely to occur. However, the reactivity of **2** toward **3** was lower than that of typical PhSeOTf prepared from PhSeCl and AgOTf. For example, the *trans* isomer of **3d** (*trans*-2-allylcyclohexanol) which was smoothly cyclized by PhSeOTf even at -78 °C¹³ did not react with **2** in the same conditions. And even in the presence of trifluoromethanesulfonic acid, oxyselenenylation of **3a** to **4a** by **1** has not proceeded. Thus, we suppose that the reagent (**2**) contains a highly electrophilic organoselenium complex of copper. Mechanisms of stereoselectivities of **4a**, **4b**, **4c**, and **4f** were explained by the steric repulsion (**4a** and **4f**: *trans* predominating) and the electronic attraction (**4b** and **4c**: *cis* predominating) between the allylic substituent (R) and electropositive Se atom in 3-membered ring cationic seleniranium intermediates.^{15,17–19}

The following is a typical example of the reaction of **3** with **2**: A mixture of Cu(OTf)₂ (0.36 g, 1.0 mmol) and **1** (0.18 g, 1.0 mmol) in toluene (3.5 mL) was stirred at 50 °C for 1 h. To the resulting suspension was added **3a** (0.10 g, 0.90 mmol) at 25 °C, and the mixture was stirred for 1 h. To this was added saturated NaHCO₃ (3 mL) and dichloromethane (15 mL), and precipitate was removed by a centrifuge. The organic solution was concentrated, and the residue was subjected to silica gel column chromatography eluting with 5% (v/v) ethyl acetate in petroleum ether. A 96:4 mixture of *trans*-**4a** and *cis*-**4a** was obtained as pale yellow oil (0.23 g, 95% yield). Spectroscopic and analytical data of the products were listed in the reference.¹⁵

ACKNOWLEDGMENTS

We acknowledge to Central Glass Co. Ltd., for the supply of trifluoromethanesulfonic acid. The author, S. M., thanks Mr. Y. Hayashi in School of Informatics and Sciences, Nagoya University for technical help.

REFERENCES

1. Recent address: Kawasaki Plastics Laboratory, Showa Denko Co. Ltd., 3 – 2 Chidori-cho, Kawasaki-ku, Kawasaki, 210 Japan
2. Reviews for selenocyanates: (a) H. Rheinboldt, "Methoden der Organischen Chemie (Houben-Weyl)", Vol. 9 ed. by E. Mueller, Georg Thieme Verlag, Stuttgart, 1955, pp. 939–951; (b) E. Bulka, "The Chemistry of Cyanates and Their Thio Derivatives, Part 2", ed. by S. Patai, John Wiley and Sons, Chichester, 1977, Chapter 19.
3. ArSeCN is one of the most suitable precursors for ArSe⁻, see: K. B. Sharpless and M. W. Young, *J. Org. Chem.*, 1975, **40**, 947.
4. 2-Nitrophenyl selenocyanate in the presence of (C₄H₉)₃P has been employed to the preparation of alkyl aryl selenides from alcohols, see: P. A. Grieco, S. Gilman, and M. Nishizawa, *J. Org. Chem.*, 1976, **41**, 1485.
5. S. Tomoda, Y. Takeuchi, and Y. Nomura, *Tetrahedron Lett.*, 1982, **23**, 1361.
6. S. Tomoda, Y. Takeuchi, and Y. Nomura, *J. Chem. Soc., Chem. Comm.*, 1982, 871.
7. A. Toshimitsu, T. Aoi, S. Uemura, and M. Okano, *J. Org. Chem.*, 1980, **45**, 1953.
8. S. Uemura, A. Toshimitsu, T. Aoi, and M. Okano, *Chem. Lett.*, 1979, 1359.
9. A. Toshimitsu, T. Aoi, S. Uemura, and M. Okano, *J. Org. Chem.*, 1981, **46**, 3021.
10. Trifluoromethanesulfonate was abbreviated as OTf.
11. S. Murata and T. Suzuki, *Chem. Lett.*, 1987, 849.
12. S. Murata and T. Suzuki, *Tetrahedron Lett.*, 1987, **28**, 4297.
13. S. Murata and T. Suzuki, *Tetrahedron Lett.*, 1987, **28**, 4415.
14. S. Murata and T. Suzuki, *Tetrahedron Lett.*, 1990, **31**, 6535.
15. H. Inoue, S. Murata, and T. Suzuki, *Liebigs Ann. Chem.*, 1994, 901.
16. M. Hoelzle and W. Jenny, *Helv. Chim. Acta*, 1958, **41**, 331.
17. A. B. Reitz, S. O. Nortey, B. E. Maryanoff, D. Liotta, and R. Monahan III, *J. Org. Chem.*, 1987, **52**, 4191.
18. Since reactions of 3-methyl-4-penten-1-ol and 3-*p*-trifluoromethylbenzoyloxy-4-penten-1-ol, which are tetrahydrofuran forming analogs of **3a** and **3c** respectively, with PhSeOTf predominantly give trans isomers, we would like to propose the thermodynamic mechanism rather than the kinetic (stereoelectronic) mechanism which is employed for explanation of stereoselectivities in haloetherification, oxymercuration, and so on. In addition, a reaction of PhSeOTf with *trans*-2-allylcyclohexanol in CH₂Cl₂ at 0 °C affords a thermodynamically favored *endo*-cyclized product. See references 13, 15, 17, and S. Current and K. B. Sharpless, *Tetrahedron Lett.*, 1978, 5075.
19. Although the molecular orbital study on 3-buten-2-ol supported the kinetically controlled *cis*-cyclization (see: S. D. Kahn, C. F. Pau, W. J. Hehre, *J. Am. Chem. Soc.*, 1986, **108**, 7396), results of our preliminary calculations on 1-Se-methyl-2-(1-methoxyethyl)-seleniranium and 1-Se-methyl-2-(1-methylpropyl)seleniranium (PM3 and *ab-initio*) exhibit that conformations in which Se⁺ and OCH₃ groups are nearby are stabilized.