

## REACTION OF INDOLE-2,3-DICARBOXYLIC ANHYDRIDE WITH GRIGNARD REAGENTS: SYNTHESIS OF 2-ACYLINDOLES

Yasuyoshi Miki,\* Hiroko Hachiken, and Ichigo Yoshikawa


Faculty of Pharmaceutical Sciences, Kinki University, 3-4-1, Kowakae, Higashi-Osaka 577, Japan

**Abstract** - Reaction of indole-2,3-dicarboxylic anhydride with methylmagnesium bromide and phenylmagnesium bromide gave 2-acetyl- and 2-benzoyl-indole-3-carboxylic acids, but with *tert*-butylmagnesium chloride, 3-pivaloylindole-2-carboxylic acids were obtained as the main products. Treatment of 2-acylindole-3-carboxylic acids with copper chromite in quinoline or potassium hydroxide gave the corresponding 2-acylindoles.

We have reported that the reduction of 1-benzylindole-2,3-dicarboxylic anhydride (**3b**) with sodium borohydride gave a 2-hydroxymethylindole-3-carboxylic acid, which was converted to murrayaquinone-A via a Diels-Alder reaction of 4*H*-furo[3,4-*b*]indole with methyl acrylate.<sup>1</sup> Reduction of the 2-carbonyl group of **3b** is much easier than that of the 3-carbonyl group, because a nitrogen in the indole is more enolizable with the 3-carbonyl group than the 2-carbonyl group.<sup>2</sup> Reaction of substituted phthalic anhydrides with Grignard reagents was controlled by electronic and steric factors.<sup>3</sup> However, in reaction of pyridinedicarboxylic anhydride with Grignard reagents<sup>4</sup> and 1-benzenesulfonyl-2-lithioindole,<sup>5</sup> a nitrogen in the pyridine decides the reactivity of the anhydride toward nucleophiles. In this paper we report the reactivity of 1-benzyl- (**3b**) and 1-benzenesulfonylindole-2,3-dicarboxylic anhydrides (**3c**) with Grignard reagents and the synthesis of 2-acylindoles.

The anhydrides (**3b,c**) were prepared from dimethyl indole-2,3-dicarboxylate (**1**)<sup>6</sup> as follows. Hydrolysis

Scheme 1



of **1** with potassium hydroxide gave an indole-2,3-dicarboxylic acid (**2a**)(R=H)(60%). The acid (**2a**) was reacted with benzyl bromide in the presence of sodium hydride in *N,N*-dimethylformamide to afford the corresponding 1-benzylindole-2,3-dicarboxylic acid (**2b**)(R=CH<sub>2</sub>Ph)(87%), which was treated with trifluoroacetic anhydride in refluxing dichloromethane to yield anhydride (**3b**)(78%). Synthesis of 1-benzenesulfonylindole-2,3-dicarboxylic anhydride (**3c**) was performed by conversion of **2a** to the corresponding 2-methoxyethoxymethyl(MEM) ester, followed by benzenesulfonylation<sup>7</sup> and treatment of **4** with 10% hydrochloric acid, then trifluoroacetic anhydride.

Reaction of 1-benzylindole-2,3-dicarboxylic anhydride (**3b**) with phenylmagnesium bromide in tetrahydrofuran at -78°C gave 2-benzoyl-1-benzylindole-3-carboxylic acid (**5a**) in 84% yield. In a similar manner, treatment of **3b** with methylmagnesium bromide afforded a mixture of 2-acetyl-1-benzylindole-3-carboxylic acid (**5c**) and 1-benzyl-2-(1-hydroxy-1-methylethyl)indole-3-carboxylic acid (**7**) in 45% and 30% yields, respectively. When 1-benzenesulfonylindole-2,3-dicarboxylic anhydride (**3c**) reacted with phenylmagnesium bromide and methylmagnesium bromide, 1-benzenesulfonyl-2-benzoyl- (**5b**) and 2-acetylindole-3-carboxylic acid (**5d**) were isolated in 52% and 69% yield, respectively.

However, treatment of **3b** with *tert*-butylmagnesium chloride yielded a mixture of 1-benzyl-2-pivaloylindole-3-carboxylic acid (**5e**) and 3-pivaloylindole-2-carboxylic acid (**6e**) in 43% and 34% yields, respectively. In a similar manner, reaction of **3c** with *tert*-butylmagnesium chloride yielded a mixture of 1-benzenesulfonyl-2-pivaloylindole-3-carboxylic acid (**5f**) and 3-pivaloylindole-2-carboxylic acid (**6f**) in 2% and 24% yields, respectively.

Scheme 2




Table 1

R <sup>1</sup>	R <sup>2</sup> MgX	Time	Yield(%)		
			<b>5</b>	<b>6</b>	<b>7</b>
CH <sub>2</sub> Ph	PhMgBr	1 h	84	-	-
SO <sub>2</sub> Ph	PhMgBr	1 h	52	-	-
CH <sub>2</sub> Ph	MeMgBr <sup>1)</sup>	1 h	45	-	30
SO <sub>2</sub> Ph	MeMgBr	3 h	69	-	-
CH <sub>2</sub> Ph	<i>tert</i> -BuMgCl	1 h	43	34	-
SO <sub>2</sub> Ph	<i>tert</i> -BuMgCl <sup>1)</sup>	24 h	2	24	-

1) 5 eq

Treatment of **5a** with copper chromite in refluxing quinoline gave 2-benzoyl-1-benzylindole (**8a**) in 80% yield and an alkaline hydrolysis of **5b** with potassium hydroxide in hot methanol afforded 2-benzoylindole

(8)(R<sup>1</sup> = H, R<sup>2</sup> = Ph) in 70% yield. In a similar manner, the conversion of **5a** to **8a**, **5c-e** were changed to 2-acetyl-1-benzylindole (**8c**), 2-acetyl-1-benzenesulfonylindole (**8d**), and 1-benzyl-2-pivaloylindole (**8e**) in 80%, 69%, and 88% yields, respectively.

Scheme 3





Table 2

5	R <sup>1</sup>	R <sup>2</sup>	Method	Yield(%)
a	CH <sub>2</sub> Ph	Ph	A	80
b	SO <sub>2</sub> Ph	Ph	B	70
c	CH <sub>2</sub> Ph	Me	A	80
d	SO <sub>2</sub> Ph	Me	A	69
e	CH <sub>2</sub> Ph	<i>tert</i> -Bu	A	88

method A: copper chromite / quinoline

method B: KOH / MeOH

The structure of two isomeric products (**5e,f**) and (**6e,f**) was readily assigned by comparison of the C-4 protons of 2-pivaloylindole-3-carboxylic acids (**5e**;  $\delta$  8.24, **5f**;  $\delta$  8.06-8.13) with those of 3-pivaloylindole<sup>8</sup> ( $\delta$  8.45) and 3-pivaloylindole-2-carboxylic acids (**6e**;  $\delta$  7.01-7.57, **6f**;  $\delta$  7.20-7.50) with 3-pivaloylindole-2-carboxylic acid<sup>8</sup> ( $\delta$  6.90-7.62). A carbonyl group at the 2-position in indole-2,3-dicarboxylic anhydride (**3**) is more reactive toward a nucleophile than a carbonyl group at the 3-position because a benzenoid structure (**9**) is more stable than a quinoid structure (**10**). Therefore, reaction of **3b** with phenylmagnesium bromide and methylmagnesium bromide gave 2-acyl-1-benzylindole-3-carboxylic acids (**5**) but with the bulkier *tert*-butylmagnesium chloride the reaction afforded a mixture of 1-benzyl-2-pivaloylindole-3-carboxylic acid (**5e**) and 1-benzyl-3-pivaloylindole-2-carboxylic acid (**6e**) because of the steric hindrance of the benzyl group. We expected that the 1-benzenesulfonyl group, an electron-withdrawing group, in indole-2,3-dicarboxylic anhydride (**3c**) activates the carbonyl group at the 3-position rather than at the 2-position, but the results of the reactivity of **3c** toward Grignard reagents were similar to the reactivity of **3b**.



**b**; R<sup>1</sup> = CH<sub>2</sub>Ph, **c**; R<sup>1</sup> = SO<sub>2</sub>Ph

## EXPERIMENTAL

Melting points were determined on a Yanagimoto micromelting point apparatus and are uncorrected. The

$^1\text{H-NMR}$  spectra were determined on a JEOL JNM-GSX 270 spectrometer using tetramethylsilane as an internal standard. The IR spectra were recorded with a JASCO FT/IR-7000 spectrophotometer. The high MS were recorded on a JOEL JMS-HX100 spectrometer. Column chromatography was performed on E. Merck silica gel 60 (70-230 mesh or 230-400 mesh). Tetrahydrofuran (THF) was distilled from sodium and benzophenone prior to use. Phenylmagnesium bromide (1.0 M solution in THF) and methylmagnesium bromide (1.4 M solution in toluene and THF) were purchased from Aldrich. *tert*-Butylmagnesium chloride (1 M solution in THF) was purchased from Kanto Chemical.

#### Indole-2,3-dicarboxylic Acid (2a)

A solution of dimethyl indole-2,3-dicarboxylate (**1**)<sup>6</sup> (9.32 g, 40 mmol) in methanol (40 mL) and 4N potassium hydroxide solution (40 mL, 160 mmol) was heated at 70 °C for 2 h under stirring. The reaction mixture was diluted with water, and washed with Et<sub>2</sub>O. The aqueous solution was acidified with 10% hydrochloric acid to give the indole-2,3-dicarboxylic acid (**2a**) (4.88 g, 60%), mp 258-260°C (MeOH-H<sub>2</sub>O) as a pale yellow solid. IR (nujol) 3220, 1700 cm<sup>-1</sup>;  $^1\text{H-NMR}$  (DMSO-d<sub>6</sub>)  $\delta$ : 7.28 (1H, dt,  $J = 8, 1$  Hz, H-5 or H-6), 7.37 (1H, dt,  $J = 8, 1$  Hz, H-6 or H-5), 7.59 (1H, dd,  $J = 8, 1$  Hz, H-7), 8.16 (1H, br d,  $J = 8$  Hz, H-4), 12.66 (1H, br s, NH). *Anal.* Calcd For C<sub>10</sub>H<sub>7</sub>NO<sub>4</sub>: C, 58.39; H, 3.43; N, 6.81. Found: C, 58.40; H, 3.64; N, 6.76.

#### 1-Benzylindole-2,3-dicarboxylic Anhydride (3b)

Benzyl bromide (18 mL, 0.15 mol) was added dropwise to a suspension of sodium hydride (10.00 g, 0.25 mole, 60% dispersion in mineral oil) and indole-2,3-dicarboxylic acid (**2a**) (10.25 g, 0.05 mol) in *N,N*-dimethylformamide (100 mL) at 0 °C. After the mixture stirred for 1 h at rt, the mixture was poured into water and washed with Et<sub>2</sub>O. The aqueous layer was acidified (pH=1) with concentrated hydrochloric acid to give a precipitate, which was collected by filtration to afford 1-benzylindole-2,3-dicarboxylic acid (**2b**) (12.82 g, 87%), mp 203-204°C (lit.,<sup>9</sup> mp 198 °C) (MeOH-H<sub>2</sub>O) as a pale yellow solid. IR (nujol) 1693 cm<sup>-1</sup>;  $^1\text{H-NMR}$  (DMSO-d<sub>6</sub>)  $\delta$ : 6.13 (2H, s, CH<sub>2</sub>Ph), 7.04-7.44 (8H, m, aromatic protons), 8.45-8.50 (1H, m, H-4). *Anal.* Calcd for C<sub>17</sub>H<sub>13</sub>NO<sub>4</sub>: C, 69.15; H, 4.44; N, 4.74. Found: C, 69.19; H, 4.58; N, 4.72.

A suspension of 1-benzylindole-2,3-dicarboxylic acid (**2b**) (8.85 g, 30 mmol) and trifluoroacetic anhydride (8.5 mL, 60 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (60 mL) was refluxed for 2 h. The reaction mixture was evaporated off to afford a solid, which was washed with *n*-hexane-CH<sub>2</sub>Cl<sub>2</sub> (3 : 1) to give 1-benzylindole-2,3-dicarboxylic anhydride (**3b**) (6.52g, 78%), mp 150-152°C (AcOEt) as a pale yellow solid. IR (nujol) 1836, 1772 cm<sup>-1</sup>;  $^1\text{H-NMR}$  (DMSO-d<sub>6</sub>)  $\delta$ : 5.85 (2H, s, CH<sub>2</sub>Ph), 7.08-7.51 (8H, m, aromatic protons), 8.23-8.29 (1H, m, H-4). *Anal.* Calcd for C<sub>17</sub>H<sub>11</sub>NO<sub>3</sub>: C, 73.65; H, 4.00; N, 5.05. Found: C, 73.65; H, 4.19; N, 5.08.

#### Bis(2-methoxyethoxymethyl) 1-Benzenesulfonylindole-2,3-dicarboxylate (4)

To a mixture of indole-2,3-dicarboxylic acid (**2a**) (5.33 g, 26 mmol) and *N,N*-diisopropylethylamine (9.2 mL, 53 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL) was added 2-methoxyethoxymethyl chloride (6.0 mL, 52 mmol) at 0 °C under argon and the mixture was stirred for 1 h. Water was added to the reaction mixture, which was

extracted with  $\text{CH}_2\text{Cl}_2$ . The organic extracts were washed with water, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. The residue was purified by column chromatography (*n*-hexane : AcOEt = 10 : 1) to give bis(2-methoxyethoxymethyl) indole-2,3-dicarboxylate (8.16g, 82%). IR (neat) 3298, 1712  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 3.38 (3H, s,  $\text{OCH}_3$ ), 3.39 (3H, s,  $\text{OCH}_3$ ), 3.50-3.95 (4H, m,  $\text{OCH}_2\text{CH}_2\text{O}$ ), 5.58 (2H, s,  $\text{CO}_2\text{CH}_2\text{O}$ ), 5.62 (2H, s,  $\text{CO}_2\text{CH}_2\text{O}$ ), 7.20-7.45 (3H, m, aromatic protons), 8.09 (1H, dd,  $J = 8$ , 1 Hz, H-4), 9.78 (1H, br s, NH); HRMS  $m/z$  ( $\text{M}^+$ ) calcd for  $\text{C}_{18}\text{H}_{23}\text{NO}_8$ : 381.1424. Found: 381.1432.

Benzenesulfonyl chloride (2.8 mL, 22 mmol) was added to a suspension of bis(2-methoxyethoxymethyl) indole-2,3-dicarboxylate (8.16 g, 21 mmol) and tetra-*n*-butylammonium bisulfate (728 mg, 2.1 mmol) in 50% potassium hydroxide solution (21 mL) and methylene chloride (63 mL) at  $-20^\circ\text{C}$  and the reaction mixture was stirred for 2 h. The reaction mixture was poured into water, extracted with  $\text{CH}_2\text{Cl}_2$ , and washed with water. The extracts were dried over  $\text{Na}_2\text{SO}_4$ , and evaporated off to give a residue, which was chromatographed (*n*-hexane : AcOEt = 10 : 1) to afford bis(2-methoxyethoxymethyl) 1-benzenesulfonylindole-2,3-dicarboxylate (**4**) (8.98 g, 80%) as a brown oil. IR (neat) 1724  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 3.38 (3H, s,  $\text{OCH}_3$ ), 3.42 (3H, s,  $\text{OCH}_3$ ), 3.54-4.06 (8H, m,  $\text{OCH}_2\text{CH}_2\text{O}$ ), 5.58 (2H, s,  $\text{CO}_2\text{CH}_2\text{O}$ ), 5.74 (2H, s,  $\text{CO}_2\text{CH}_2\text{O}$ ), 7.33-7.65 (5H, m, aromatic protons), 7.99-8.20 (4H, m, aromatic protons); HRMS  $m/z$  ( $\text{M}^+$ ) calcd for  $\text{C}_{24}\text{H}_{27}\text{NO}_{10}\text{S}$ : 521.1356. Found: 521.1387.

### 1-Benzenesulfonylindole-2,3-dicarboxylic Anhydride (**3c**)

A mixture of bis(2-methoxyethoxymethyl) 1-benzenesulfonylindole-2,3-dicarboxylate (**4**) (8.98 g, 17 mmol) and 10% hydrochloric acid (9 mL) in THF (86 mL) was refluxed for 2 h. The mixture was poured into water and extracted with  $\text{CH}_2\text{Cl}_2$ . The organic extracts were washed with water, dried over with  $\text{Na}_2\text{SO}_4$ , and concentrated to give 1-benzenesulfonylindole-2,3-dicarboxylic acid (5.95 g, 100%), mp  $222-223^\circ\text{C}$  (decomp)(from  $\text{MeOH-H}_2\text{O}$ ). IR (nujol) 1729, 1682  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ )  $\delta$ : 7.33-7.77 (5H, m, aromatic protons), 7.96-8.14 (4H, m, aromatic protons). Anal. Calcd for  $\text{C}_{16}\text{H}_{11}\text{NO}_6\text{S}$ : C, 55.65; H, 3.21; N, 4.06. Found: C, 55.50; H, 3.35; N, 3.94.

A suspension of 1-benzenesulfonylindole-2,3-dicarboxylic acid (5.95g, 17 mmol) and trifluoroacetic anhydride (4.9 mL, 35 mmol) in  $\text{CH}_2\text{Cl}_2$  (34 mL) was refluxed for 2 h. After the mixture was cooled, a separated precipitate was collected by filtration to yield 1-benzenesulfonylindole-2,3-dicarboxylic anhydride (**3c**) (4.36 g, 77%), mp  $228-229^\circ\text{C}$  (AcOEt). IR (nujol) 1840, 1778  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ )  $\delta$ : 7.34-7.78 (5H, m, aromatic protons), 7.96-8.14 (4H, m, aromatic protons). Anal. Calcd for  $\text{C}_{16}\text{H}_9\text{NO}_5\text{S}$ : C, 58.71; H, 2.77; N, 4.28. Found: C, 58.52; H, 3.06; N, 4.52.

### Reaction of Indole-2,3-dicarboxylic Anhydride (**3b,c**) with Grignard Reagents (General Procedure)

To a solution of indole-2,3-dicarboxylic anhydride (**3**) (0.2 mmol) in THF (2-4 mL) at  $-78^\circ\text{C}$  under argon was added Grignard reagents (0.4 mmol). The reaction mixture was acidified with 10% hydrochloric acid and the mixture was extracted with  $\text{CH}_2\text{Cl}_2$ . The combined extracts were washed with water and dried

over  $\text{Na}_2\text{SO}_4$ . The solvent was evaporated off to afford a residue, which was purified by column chromatography ( $\text{CH}_2\text{Cl}_2$  :  $\text{MeOH}$  = 10 : 1).

**2-Benzoyl-1-benzylindole-3-carboxylic Acid (5a)**; mp 258-260°C (from EtOH). IR (nujol) 1669  $\text{cm}^{-1}$ ;  $^1\text{H}$ -nmr ( $\text{DMSO-d}_6$ )  $\delta$ : 5.35 (2H, s,  $\text{CH}_2\text{Ph}$ ), 7.00-7.74 (9H, m, aromatic protons), 8.09-8.15 (1H, m, H-4). Anal. Calcd for  $\text{C}_{23}\text{H}_{17}\text{NO}_3$ : C, 77.73; H, 4.82; N, 3.94. Found: C, 77.68; H, 5.00; N, 3.89.

**1-Benzenesulfonyl-2-benzoylindole-3-carboxylic Acid (5b)**; mp 238-240°C (from EtOH). IR (nujol) 1679  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR ( $\text{DMSO-d}_6$ )  $\delta$ : 7.38-8.18 (14H, m, aromatic protons). HRMS  $m/z$  ( $\text{M}^+$ ) calcd for  $\text{C}_{22}\text{H}_{15}\text{NO}_5\text{S}$ : 405.0671. Found: 405.0695. **5b** was identified as methyl 1-benzenesulfonyl-2-benzoylindole-3-carboxylate by treatment of diazomethane; mp 174-175.5°C (from AcOEt-*n*-hexane). IR (nujol) 1716, 1669  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR ( $\text{CDCl}_3$ )  $\delta$ : 3.67 (3H, s,  $\text{CH}_3$ ), 7.36-7.66 (8H, m, aromatic protons), 7.90-8.20 (6H, m, aromatic protons). Anal. Calcd for  $\text{C}_{23}\text{H}_{17}\text{NO}_5\text{S}$ : C, 65.86; H, 4.08; N, 3.34. Found: C, 65.66; H, 4.27; N, 3.37.

**2-Acetyl-1-benzylindole-3-carboxylic Acid (5c)**; mp 198.5-200°C (from EtOH). IR (nujol) 1707, 1658  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR ( $\text{DMSO-d}_6$ )  $\delta$ : 2.44 (3H, s,  $\text{CH}_3$ ), 5.42 (2H, s,  $\text{CH}_2\text{Ph}$ ), 7.02-7.58 (8H, m, aromatic protons), 8.03-8.11 (1H, m, H-4). Anal. Calcd for  $\text{C}_{18}\text{H}_{15}\text{NO}_3$ : C, 73.71; H, 5.15; N, 4.78. Found: C, 73.51; H, 5.30; N, 4.76.

**1-Benzyl-2-(1-hydroxy-1-methylethyl)indole-3-carboxylic Acid (7)**; mp 166-167°C (from EtOH). IR (nujol) 1633  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR ( $\text{DMSO-d}_6$ )  $\delta$ : 1.68 (6H, s, 2 x  $\text{CH}_3$ ), 5.93 (2H, s,  $\text{CH}_2\text{Ph}$ ), 6.92-7.30 (8H, m, aromatic protons), 7.88-7.93 (1H, m, H-4). Anal. Calcd for  $\text{C}_{19}\text{H}_{19}\text{NO}_3$ : C, 73.77; H, 6.19; N, 4.53. Found: C, 73.44; H, 6.21; N, 4.49.

**2-Acetyl-1-benzenesulfonylindole-3-carboxylic Acid (5d)**; mp 249-251°C (from EtOH). IR (nujol) 1714, 1679  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR ( $\text{DMSO-d}_6$ )  $\delta$ : 2.73 (3H, s,  $\text{CH}_3$ ), 7.36-7.80 (5H, m, aromatic protons), 7.93-8.09 (4H, m, aromatic protons). Anal. Calcd for  $\text{C}_{17}\text{H}_{13}\text{NO}_5\text{S}$ : C, 59.47; H, 3.82; N, 4.08. Found: C, 58.97; H, 4.03; N, 3.76.

**1-Benzyl-2-pivaloylindole-3-carboxylic Acid (5e)**; mp 228-230°C (from EtOH). IR (nujol) 1696, 1661  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR ( $\text{DMSO-d}_6$ )  $\delta$ : 1.28 (9H, s,  $\text{C}(\text{CH}_3)_3$ ), 5.27 (2H, s,  $\text{CH}_2\text{Ph}$ ), 7.07-7.35 (8H, m, aromatic protons), 8.24 (1H, br d,  $J$  = 8 Hz, H-4). Anal. Calcd for  $\text{C}_{21}\text{H}_{21}\text{NO}_3$ : C, 75.20; H, 6.31; N, 4.18. Found: C, 75.05; H, 6.38; N, 4.27.

**1-Benzyl-3-pivaloylindole-2-carboxylic Acid (6e)**; mp 219-220°C (from EtOH). IR (nujol) 1683  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR ( $\text{DMSO-d}_6$ )  $\delta$ : 1.21 (9H, s,  $\text{C}(\text{CH}_3)_3$ ), 5.86 (2H, s,  $\text{CH}_2\text{Ph}$ ), 7.01-7.55 (9H, m, aromatic protons). Anal. Calcd for  $\text{C}_{21}\text{H}_{21}\text{NO}_3$ : C, 75.20; H, 6.31; N, 4.18. Found: C, 74.82; H, 6.42; N, 4.09.

**1-Benzenesulfonyl-2-pivaloylindole-3-carboxylic Acid (5f)**; mp 92-97°C (from *n*-hexane). IR ( $\text{CHCl}_3$ ) 1707  $\text{cm}^{-1}$ ;  $^1\text{H}$ -NMR ( $\text{CDCl}_3$ )  $\delta$ : 1.39 (9H, s,  $\text{C}(\text{CH}_3)_3$ ), 7.30-7.57 (5H, m, aromatic

protons), 7.91-7.98 (2H, m, H-2'), 8.06 (1H, dd,  $J = 8, 1$  Hz, H-4 or H-7), 8.13 (1H, br d,  $J = 7$  Hz, H-7 or H-4). HRMS  $m/z$  ( $M^+$ ) calcd for  $C_{20}H_{19}NO_5S$ : 385.0984. Found: 385.0984.

**1-Benzenesulfonyl-3-pivaloylindole-2-carboxylic Acid (6f)**: mp 142-145°C (from AcOEt-*n*-hexane). IR ( $CHCl_3$ ) 1690  $cm^{-1}$ ;  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 1.16 (9H, s,  $C(CH_3)_3$ ), 7.20-7.50 (6H, m, arom), 7.92 (2H, br d,  $J = 8$  Hz, H-2'), 8.06 (1H, br d,  $J = 8$  Hz, H-7). HRMS  $m/z$  ( $M^+$ ) calcd for  $C_{20}H_{19}NO_5S$ : 385.0984. Found: 385.0974. **6f** was identified as methyl 1-benzenesulfonyl-3-pivaloylindole-2-carboxylate by treatment of diazomethane; mp 75-76°C (AcOEt-*n*-hexane). IR (nujol) 1717, 1670  $cm^{-1}$ ;  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 1.25 (9H, s,  $C(CH_3)_3$ ), 3.90 (3H, s,  $CH_3$ ), 7.26-7.63 (6H, m, aromatic protons), 7.99-8.04 (2H, m, H-2'), 8.11 (1H, dt,  $J = 8, 1$  Hz, H-7). Anal. Calcd for  $C_{21}H_{21}NO_5S$ : C, 63.14; H, 5.30; N, 3.51. Found: C, 63.05; H, 5.24; N, 3.86.

### Conversion of 2-Acylindole-3-carboxylic Acid (5) to 2-Acylindole (8)

#### 1) Copper Chromite in Quinoline

##### 2-Benzoyl-1-benzylindole (8a)

A mixture of 2-benzoyl-1-benzylindole-3-carboxylic acid (**5a**) (36 mg, 0.1 mmol) and copper chromite (4 mg) in quinoline (1 mL) was heated to reflux for 20 min. Water was added to the mixture and the aqueous mixture was extracted with  $CH_2Cl_2$ . The extracts were washed with water, then with 5% hydrochloric acid and water. The solution was dried over  $Na_2SO_4$  and evaporated off to give a residue, which was purified by column chromatography (*n*-hexane : AcOEt = 10 : 1) to yield 2-benzoyl-1-benzylindole (**8a**) (25 mg, 80%), mp 108-109°C (from AcOEt-*n*-hexane). IR (nujol) 1644  $cm^{-1}$ ;  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 5.87 (2H, s,  $CH_2Ph$ ), 7.06-7.92 (14H, m, aromatic protons). Anal. Calcd for  $C_{22}H_{17}NO$ : C, 84.86; H, 5.50; N, 4.50. Found: C, 84.97; H, 5.68; N, 4.55.

##### 2-Acetyl-1-benzylindole (8c)

Using a procedure similar to that described for the preparation of **8a**, **8c** (80%) was obtained from **5a**, mp 129.5-131°C (lit.,<sup>10</sup> mp 130-131.5°C) (MeOH). IR (nujol) 1665  $cm^{-1}$ ;  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 2.61 (3H, s,  $CH_3$ ), 5.86 (2H, s,  $CH_2Ph$ ), 7.20-7.40 (9H, m, aromatic protons), 7.73 (1H, dt,  $J = 8, 1$  Hz, H-4 or H-7).

##### 2-Acetyl-1-benzenesulfonylindole (8d)

Using a procedure similar to that described for the preparation of **8a**, **8d** (69%) was obtained from **5d**, mp 89-90°C (lit.,<sup>11</sup> mp 89-90°C) (MeOH). IR (nujol) 1690  $cm^{-1}$ ;  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 2.65 (3H, s,  $CH_3$ ), 7.14 (1H, d,  $J = 1$  Hz, H-3), 7.27-7.60 (6H, m, aromatic protons), 7.92-7.99 (2H, m, H-2'), 8.17 (1H, br d,  $J = 8$  Hz, H-4 or H-7).

##### 1-Benzyl-2-pivaloylindole (8e)

Using a procedure similar to that described for the preparation of **8a**, **8e** (88%) was obtained from **5e**, mp 77-78°C (from *n*-hexane). IR (nujol) 1657  $cm^{-1}$ ;  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 1.32 (9H, s,  $C(CH_3)_3$ ), 5.69 (2H, s,  $CH_2Ph$ ), 6.93-7.00 (2H, m, aromatic protons), 7.12-7.39 (7H, m, aromatic protons), 7.74 (1H, dt,

$J = 8, 1$  Hz, H-4). Anal. Calcd for  $C_{20}H_{21}NO$ : C, 82.43; H, 7.26; N, 4.81. Found: C, 82.27; H, 7.27; N, 4.63.

## 2) Potassium Hydroxide in Methanol

### 2-Benzoylindole (8b)

A solution of 1-benzenesulfonyl-2-benzoylindole-3-carboxylic acid (**5b**)(61 mg) in 4N potassium hydroxide solution (0.15 mL) and methanol (1.5 mL) was heated to reflux for 5 h and water was added to the mixture. The aqueous solution was extracted with  $CH_2Cl_2$  and the extracts were washed with water, dried over  $Na_2SO_4$ , and evaporated off to give a residue. The residue was purified by column chromatography (*n*-hexane : AcOEt = 20 : 1) to a solid, which was recrystallized from *n*-hexane- $CHCl_3$  to afford 2-benzoylindole (**10b**)(23 mg, 70%), mp 149-150°C (lit.,<sup>12</sup> mp 151-152°C)(from  $CHCl_3$ -*n*-hexane). IR ( $CHCl_3$ ) 3425, 1625  $cm^{-1}$ ;  $^1H$ -nmr ( $CDCl_3$ )  $\delta$ : 7.12-7.75 (8H, m, aromatic protons), 7.96-8.02 (2H, m, aromatic protons), 9.28 (1H, br s, NH).

## REFERENCES

1. Y. Miki and H. Hachiken, *Synlett*, 1993, 333.
2. R. J. Sundberg, 'Comprehensive Heterocyclic Chemistry,' Vol. 4, ed. by C. W. Bird and G. W. H. Cheeseman, Pergamon Press, New York, 1984, p. 288; R. J. Sundberg, 'The Chemistry of Indole', Academic Press, New York, 1970, p. 401.
3. M. S. Newman and C. D. McCleary, *J. Am. Chem. Soc.*, 1941, **63**, 1542; M. S. Newman and C. W. Muth, *J. Am. Chem. Soc.*, 1950, **72**, 5191; R. Goncalves, M. R. Kegelman, and E. V. Brown, *J. Org. Chem.*, 1952, **17**, 705; R. Melby, R. Crawford, D. McGreer, and R. B. Sandin, *J. Am. Chem. Soc.*, 1956, **78**, 3816; M. S. Newman and P. G. Scheurer, *J. Am. Chem. Soc.*, 1956, **78**, 5004; Z. Horii, H. Hakusui, T. Momose, and E. Yoshino, *Chem. Pharm. Bull.*, 1968, **16**, 1251; D. G. Miller, S. Trenbeath, and C. J. Sih, *Tetrahedron Lett.*, 1976, 1637; J. S. Swenton, D. K. Jackson, M. J. Manning, and P. W. Reynolds, *J. Am. Chem. Soc.*, 1978, **100**, 6182.
4. P. Canonne and M. Belley, *Can. J. Chem.*, 1987, **65**, 1885.
5. G. W. Grille, M. G. Saulnier, J. A. O.-Nutaitis, and D. M. Ketcha, *J. Org. Chem.*, 1992, **57**, 5891.
6. O. Diels and J. Reese, *Ann.*, 1934, **511**, 168.
7. E. Wenkert, P. D. R. Moeller, and S. R. Piettre, *J. Am. Chem. Soc.*, 1988, **110**, 7188.
8. Y. Murakami, M. Tani, M. Suzuki, K. Sudoh, M. Uesato, K. Tanaka, and Y. Yokoyama, *Chem. Pharm. Bull.*, 1985, **33**, 4707.
9. O. Diels and J. Rees, *Ann.*, 1935, **519**, 147.
10. Y. Murakami, T. Watanabe, A. Kobayashi, and Y. Yokoyama, *Synthesis*, 1984, 738.
11. D. M. Ketcha, B. A. Lieurance, D. F. J. Homan, and G. W. Gribble, *J. Org. Chem.*, 1989, **54**, 4350.
12. R. J. Sundberg, *J. Org. Chem.*, 1965, **30**, 3604.