

**A LUPIN ALKALOID, (–)-TENUAMINE (NORLUSITANINE),  
FROM *MAACKIA TENUIFOLIA***

Yong-Hong Wang<sup>a</sup>, Kimio Higashiyama<sup>a</sup>, Hajime Kubo<sup>a</sup>, Jia-Shi Li<sup>b</sup>,  
and Shigeru Ohmiya<sup>\*a</sup>

<sup>a</sup>Institute of Medicinal Chemistry, Hoshi University, Ebara 2-4-41, Shinagawa-ku,  
Tokyo 142-8501, Japan

<sup>b</sup>Beijing University of Traditional Chinese Medicine, 11, Beisan Huan Dong Ave,  
Beijing 100029, China

E-mail: ohmiya@hoshi.ac.jp

**Abstract**– A lupin alkaloid, (–)-tenuamine, was isolated from the stem parts of *Maackia tenuifolia* together with eight known alkaloids. The structure of (–)-tenuamine was determined to be (–)-acetylaminoethylindolizidine by comparison of its chemical and spectroscopic data with those of (–)-lusitanine, (+)-tashiromine and indolizidine.

In the course of our phytochemical study on *Maackia* (Leguminosae) plants, we have reported maackiamine,<sup>1</sup> tashiromine,<sup>2</sup> and camoensidine<sup>3</sup> which all contain a pyrrolizidine or indolizidine ring. It is interesting from the perspectives of both chemotaxonomy and biosynthesis that *Maackia* species accumulate unusual lupin alkaloids containing a pyrrolizidine or indolizidine ring together with common lupin alkaloids with a piperidine or quinolizidine ring (Figure 1). In the present paper, we report the isolation and structure determination of (–)-tenuamine (**1**), a new lupin alkaloid with an indolizidine ring, from the stems of *Maackia tenuifolia*, together with (–)-lusitanine<sup>4</sup> which contains a quinolizidine ring, and seven other known alkaloids.

From the air-dried stems of *Maackia tenuifolia*, a new alkaloid (**1**) was isolated as colorless needles, mp 136–138 °C,  $[\alpha]_D^{23}$  –4.5° (*c* 0.12, EtOH) in a yield of 1.2% of the total base by repeated column chromatography. We also isolated eight known lupin alkaloids, (–)-anagyrine, (–)-*N*-methylcytisine, (–)-*N*-formylcytisine, (–)-cytisine (main base), (–)-epibatifoline, (–)-lusitanine, (+)-epilupinine and (–)-12,12'-methylenedicytisine. The known alkaloids were identified by direct comparison with authentic samples (mp,  $[\alpha]_D$ , TLC, IR, NMR and MS).<sup>5</sup>

The chromatographic behavior of alkaloid (**1**) on a silica gel column was very similar to that of lusitanine (**2**). The IR spectrum of **1** also showed a similar pattern to that of **2**, with an amide carbonyl band at 1660 cm<sup>-1</sup> and a N-H band at 3330 cm<sup>-1</sup>. The molecular formula of **1** was determined to be C<sub>11</sub>H<sub>18</sub>N<sub>2</sub>O by the <sup>13</sup>C NMR spectrum (Table 1) and HREIMS spectrometry at *m/z* 194.1409 (*M*<sup>+</sup>, calcd for 194.1419). In the

EIMS spectrum of **1**.  $M^+$  at  $m/z$  (rel. int.) 194 (55) and fragment ions at  $m/z$  151 ( $M^+ - \text{COCH}_3$ , 40), 136 ( $M^+ - \text{NHCOCH}_3$ , 55), 122 (100) and 96 (38) were one methylene less than the corresponding ions of **2** at  $m/z$  208 ( $M^+$ ,  $\text{C}_{12}\text{H}_{20}\text{N}_2\text{O}$ , 70), 166 (90), 165 (35), 150 (30), 136 (100) and 110 (82), respectively. These suggested that **1** was a congener of lusitanine. The assignments in the  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were based on an analysis of  $^{13}\text{C} - ^1\text{H}$  and  $^1\text{H} - ^1\text{H}$  COSY. The  $^1\text{H}$  NMR spectrum of **1** also suggested the presence of an acetylaminomethylidene group based on signals of an olefinic methine proton at  $\delta$  6.62 (1H, d,  $J=10.4$  Hz), which was coupled with 11-NH at  $\delta$  7.65 (1H, d,  $J=10.4$  Hz), and also on the signal of the methyl of an acetylamine group at  $\delta$  2.04 (3H, s). In the  $^{13}\text{C}$  NMR spectrum, the signals of C-2 ( $\delta$  52.6, t), C-9 ( $\delta$  54.4, t) and C-6 ( $\delta$  65.8, d) adjacent to the nitrogen (N-1) coincided with those of indolizidine (**5**) and tashiromine (**3**), but were different from those of quinolizidine (**6**) and epilupinine (**4**). The substituent effects of the acetylaminomethylidene group in the  $^{13}\text{C}$  NMR signals at C-4 (+1.0), C-6 (+1.7) and C-7 (-4.0) of **1**, compared to those of an indolizidine ring, were also similar to those of **2** (Table 1).

Table 1.  $^{13}\text{C}$  NMR data of **1**, **2**, **3**, **4**, indolizidine (**5**) and quinolizidine (**6**) in  $\text{CDCl}_3$

C	<b>1</b>	<b>3</b>	<b>5</b>	<b>1-5</b>	C	<b>2</b>	<b>4</b>	<b>6</b>	<b>2-6</b>
2	52.6	52.7	52.7	-0.1	2	57.0	57.0	56.4	+0.6
3	25.5	25.2	25.1	+0.4	3	24.2	24.9	25.6	-1.4
4	25.2	29.2	24.2	+1.0	4	27.8	29.5	24.4	+3.4
5	120.2	44.7	30.7	+89.5	5	121.6	43.8	33.2	+88.4
6	65.8	66.4	64.1	+1.7	6	64.2	64.4	62.9	+1.3
7	26.1	27.6	30.1	-4.0	7	26.4	28.3	33.2	-6.8
8	20.3	20.3	20.3	0.0	8	25.1	24.6	24.4	+0.7
					9	25.4	25.5	25.6	-0.2
9	54.4	54.2	53.9	+0.5	10	56.4	56.6	56.4	0.0
10	114.7	65.9			11	116.2	64.1		
12	167.7				13	167.9			
13	23.2				14	23.3			

The above data suggested the presence of an indolizidine ring in the structure of **1**, instead of a quinolizidine ring in **2**, and an acetylaminomethylidene group at the 5-position. Accordingly, the structure of **1** was presumed to be 5-acetylaminomethylideneindolizidine. As far as we know, this is the first isolation of the compound which has an indolizidine ring in its structure, corresponding to lusitanine (**2**) with a quinolizidine ring. Thus, we propose **1** to be named tenuamine.

It is interesting that *M. tenuifolia* accumulates the two alkaloids, tenuamine and lusitanine, which contain an indolizidine ring and a quinolizidine ring, respectively. The unusual lupin alkaloids that contain a pyrrolizidine or indolizidine ring have so far been isolated from *Maackia* plants are shown in Figure 1, together with common lupin alkaloids with a piperidine or quinolizidine ring. It can be speculated that

*Maackia* species can utilize ornithine instead of lysine as a precursor amino acid for alkaloids or can transform the piperidine moiety to the corresponding pyrrolidine group.




Figure 1. Two typical lupin alkaloids that coexist in the *Maackia* plants

## EXPERIMENTAL

**General Experimental Procedures.** Melting points were determined on Yanagimoto micro melting point apparatus and are uncorrected. Optical rotations were measured on a JASCO DIP-1000 polarimeter. IR spectra were measured with a JASCO FT/IR-200 Fourier Transform Infrared Spectrophotometer. The high and low resolution MS were measured at 70 eV using direct inlet system with a JEOL JMS-600W instruments. NMR spectra were recorded with a 500 MHz JEOL NMR instruments using TMS as an internal standard and CDCl<sub>3</sub> as solvent. TLC was conducted on precoated silica gel plates (Merck 60 F<sub>254</sub>).

**Extraction and Isolation.** The stems of *M. tenuifolia* were collected in Zhejiang Province, China, September, 1998, and identified by Director of Jiangxi Jioujiang Forest and Plant Research, Ce-ming Tan. A voucher specimen (No. 981013) is deposited in the Herbarium of the same Forest and Plant Research. The air-dried stem parts (1.6 kg) were extracted with 75% MeOH (x 3) at room temperature for 24 h. The combined extracts were concentrated and acidified with 10% HCl to pH 2. The acid phase was washed with

Et<sub>2</sub>O ( x 3), basified with 25% NH<sub>4</sub>OH to pH 11 and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The aqueous phase was saturated with K<sub>2</sub>CO<sub>3</sub> and extracted with CH<sub>2</sub>Cl<sub>2</sub> repeatedly until it became negative to Dragendorff's reagent. The all CH<sub>2</sub>Cl<sub>2</sub> extracts were combined, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to dryness *in vacuo*. The crude alkaloid was obtained as a pale brown oil in a 0.28% yield (4.5 g) of the dry stem parts and subjected to silica gel column chromatography (230-400 mesh, 400 g) with Et<sub>2</sub>O-CH<sub>2</sub>Cl<sub>2</sub>-MeOH-25%NH<sub>4</sub>OH (2.5:2.5:1:0.1), monitoring with TLC to give 11 fractions. Separation of frs 2, 3 (0.5 g) by silica gel column with Et<sub>2</sub>O-CH<sub>2</sub>Cl<sub>2</sub>-MeOH-25%NH<sub>4</sub>OH (10:10:1:0.1) yield (-)-anagyrine {[α]<sub>D</sub><sup>23</sup> -163° (c 0.28, EtOH), 0.4 g} and (-)-*N*-methylcytisine (80 mg). Fr 4 (0.4 g) yielded (-)-*N*-methylcytisine {mp 135~136°C, [α]<sub>D</sub><sup>23</sup> -216° (c 0.3, EtOH), 0.15 g} and (-)-*N*-formylcytisine {mp 169~170°C, [α]<sub>D</sub><sup>23</sup> -226° (c 0.16, EtOH), 0.2 g}. Frs 5~7 (1.5 g) were subjected to silica gel column chromatography with CH<sub>2</sub>Cl<sub>2</sub>-MeOH (4:1) to give (-)-epibatifoline {mp 207~208 °C, [α]<sub>D</sub><sup>23</sup> -132° (c 0.35, EtOH), 0.4 g}, (-)-cytisine {mp 153~155 °C, [α]<sub>D</sub><sup>23</sup> -110° (c 0.52, EtOH), about 1 g} and **2** (30 mg). In those frs. the dimer of cytisine, 12,12'-methylenedicytisine, was confirmed by <sup>1</sup>H NMR spectrum, but decomposed into cytisine on the isolation process. Fr 8 (0.6 g) contains (-)-cytisine and **2**. From frs 9,10 (0.5 g), **2** {mp 186~187 °C, [α]<sub>D</sub><sup>23</sup> -6.0° (c 0.3, EtOH), 0.35 g} and (+)-epilupinine {mp 75~76 °C, [α]<sub>D</sub><sup>23</sup> +16.8° (c 0.13, EtOH), 55 mg} were separated. Fr 11 (0.2 g) was purified by silica gel column chromatography (20 g) using Et<sub>2</sub>O-CH<sub>2</sub>Cl<sub>2</sub>-MeOH-25%NH<sub>4</sub>OH (3:1:1:0.1) to yield **1** (57 mg) and **2** (40 mg).

**Tenuamine (1)**. Colorless needles, mp 136~138°C (CHCl<sub>3</sub>), [α]<sub>D</sub><sup>23</sup> -4.5° (c 0.12, EtOH), IR (KBr): 3330, 1660 cm<sup>-1</sup> (-CONH-). HRMS *m/z* 194.1409 [M]<sup>+</sup> (calcd for C<sub>11</sub>H<sub>18</sub>N<sub>2</sub>O, 194.1419). MS *m/z* (% rel. int.): 194 [M<sup>+</sup>] (55), 179 (10), 159 (60), 151 (40), 136 (55), 122 (100), 96 (38). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 500 MHz): δ 7.65 (1H, d, *J*=10.4 Hz, 11-NH), 6.62 (1H, d, *J*=10.4 Hz, 10-H), 3.14 (2H, m, 2, 9-Heq), 2.56 (1H, dd, *J*=14.9, 4.2 Hz, 4-H), 2.43 (1H, m, 6-H), 2.22 (1H, dd, *J*=11.6, 8.5 Hz, 9-Hax), 2.16 (1H, dt, *J*=11.7, 2.8 Hz, 2-Hax), 2.04 (3H, s, 13-CH<sub>3</sub>). <sup>13</sup>C NMR data: see Table 1.

## REFERENCES

1. K. Saito, T. Yoshino, T. Sekine, S. Ohmiya, H. Otomasu, and I. Murakoshi, *Phytochemistry*, 1989, **28**, 2533.
2. S. Ohmiya, H. Kubo, H. Otomasu, K. Saito, and I. Murakoshi, *Heterocycles*, 1990, **30**, 537.
3. S. Ohmiya, H. Kubo, Y. Nakaaze, K. Saito, H. Otomasu, and I. Murakoshi, *Chem. Pharm. Bull.*, 1991, **39**, 1123.
4. K. Saito, T. Yoshino, S. Tsai, S. Ohmiya, H. Kubo, H. Otomasu, and I. Murakoshi, *Chem. Pharm. Bull.*, 1987, **35**, 1308.
5. S. Ohmiya, K. Saito, and I. Murakoshi, 'The Alkaloids' Vol. 47, ed. by Academic Press, inc., New York, 1995, pp. 4-43.