

ELECTRON TRANSFER-INITIATED PHOTOCYCLIZATION OF SUBSTITUTED α -DEHYDRO(1-NAPHTHYL)ALANINES

Tadamitsu Sakurai,* Yoshiko Morioka, Kei Maekawa, and Kanji Kubo[†]


Department of Applied Chemistry, Faculty of Technology, Kanagawa University, Kanagawa-ku, Yokohama 221-8686, Japan

[†]Institute of Advanced Material Study, 86, Kyushu University, Kasuga-koen, Kasuga, Fukuoka 816-8580, Japan

Abstract—On irradiation in the presence of triethylamine, the title compounds (**1**) having *n*-butyl and benzyl groups gave preferentially dihydrobenzoquinolinones (**2**) *via* electron transfer, whereas the introduction of the bulky *tert*-butyl and phenyl substituents into **1** resulted in the formation of benzoisoquinoline (**3**) and 1-azetine (**4**) derivatives without affording **2**.

In recent years much attention has been devoted to the synthetic application of excited-state processes initiated by electron transfer (ET), owing to the fact that many photoinduced ET reactions proceed in high chemical and quantum yields enabling the construction of heteroatom-containing polycyclic compounds.^{1,2} In the course of our systematic study towards the characterization of the excited-state reactions of substituted α -dehydrophenylalanines, we have discovered interesting photocyclization reactions giving isoquinoline and 1-azetine derivatives.³ If we introduce a naphthyl group (instead of a phenyl) into α -dehydroamino acids, it is expected that we might be able to explore ET-initiated photoreactions of naphthyl-substituted α -dehydroamino acid derivatives in the presence of an aliphatic amine.^{1,4} In this communication we demonstrate that the photoreaction of *N*-acyl α -dehydro(1-naphthyl)alanines (**1**) bearing *n*-butyl and benzyl groups on its carboxamide side chain in MeOH containing triethylamine (TEA) proceeds by an ET mechanism to give 1,2-dihydro[*f*]benzoquinolinone derivative (**2**) as the major product, whereas the introduction of bulky *tert*-butyl and phenyl substituents into **1** completely suppresses the appearance of **2** resulting in the formation of benzo[*f*]isoquinoline (**3**) and 1-azetine (**4**) derivatives (Scheme 1). The starting (*Z*)-isomers (**1a–e**) were prepared in nearly quantitative yields by the ring-opening reactions of 1-naphthyl-substituted oxazolones with primary amines.⁵ After a nitrogen-purged MeOH solution of **1a** (5.0×10^{-3} mol dm⁻³) containing TEA (0.10 mol dm⁻³) was irradiated with Pyrex-filtered light (>280 nm) from a 450 W high-pressure Hg lamp for 1.5 h at room temperature, the product mixture obtained was subjected to preparative thin-layer chromatography over silica gel, which allowed us to isolate (*Z*)-**1a** (22%: isolated yield), (*E*)-**1a** (20%), **2a** (12%), and **3a** (5%).⁶ A careful ¹H NMR analysis of the product mixture suggested the detectable formation of the *cis*-azetine isomer (**4a**) whose ring-proton signals with the $J_{3,4}$ value of 10.7 Hz were detected at 5.04 and 6.50 ppm,³ though no attempt was made to isolate **4a** owing

to its poor yield (Scheme 1).



Scheme 1 Product distribution derived from the photoreaction of (Z)-1a–e in MeOH containing TEA.

Table 1. Relation between irradiation time and composition (%) of each compound in methanol^a

Compound	Irradiation time /h						
	0	0.5	1.0	1.5	2.0	3.0	5.0
(Z)-1a	100	66.0	45.9	37.4	27.7	16.3	3.2
(E)-1a	0	28.8	38.2	36.7	32.8	20.3	4.5
2a	0	3.9	10.3	18.3	28.6	47.9	71.7
3a	0	1.0	5.1	6.9	9.9	13.9	18.4
4a	0	0.2	0.4	0.7	1.0	1.6	2.1

^a At regular time intervals, an appropriate amount of the solution being irradiated was pipetted off and concentrated to dryness in vacuo, giving the residue which was subjected to ¹H NMR analysis in DMSO-*d*₆. ¹H NMR compositions were estimated from the area ratio of a given signal for each compound.

The finding that the photoproducts (**2–4**) are stable enough such that they undergo only negligible decomposition under the irradiation conditions made it possible to monitor the reactions by means of ¹H NMR spectroscopy, as typically shown in Table 1. The result obtained for (Z)-1a demonstrates the rapid production of (E)-1a and the subsequent increase in compositions for 2a–4a with the decrease of (E)- and (Z)-isomer compositions, being consistent with the mechanism in which these excited-state isomers serve as precursors of these products. On the other hand, there was no formation of the benzoquinolinone derivative (2a) when an oxygen-free MeOH solution of (Z)-1a (5.0 × 10⁻³ mol dm⁻³) containing no TEA was irradiated with Pyrex-filtered light (UV and ¹H NMR analyses), suggesting that ET from TEA to the excited-state naphthylmethylene moiety in 1a participates in the appearance of 2a as the primary process.

In Table 2 are shown substituent and solvent effects on the product distribution and composition. Interestingly, the introduction of bulky *tert*-butyl (**1c**) and phenyl (**1d**) groups into **1** completely inhibits the appearance of the corresponding benzoquinolinone derivative (**2**), while the phenyl substituent in **1d** substantially lowers the excited-state reactivity of this naphthylalanine derivative. Because the fluorescence of **1c** was also quenched by TEA ($I_0/I = 1 + 3.0 \pm 0.4[\text{TEA}]$), ET from TEA to the excited-state (*E*)-**1c** must take place giving the **1c** anion and TEA cation radical pair (**I**) (Scheme 2). Taking into account that the enol-type biradical (**II**) [generated by ET and the subsequent proton transfer within its precursor (**I**)] may adopt a nearly planar structure, it is reasonable to conclude that the bulky *tert*-butyl and phenyl substituents exert their great steric effects on the coupling process of **II** so as to completely inhibit this process, providing also a piece of evidence for the mechanism shown in Scheme 2. There were negligible products other than (*E*)-**1**, **3**, and **4** in the reaction mixtures obtained by the irradiation of **1c** and **1d** in the presence (and even in the absence) of TEA (¹H NMR analysis). This observation suggests that on prevention of the cyclization of **II** affording **2** *via* **III**, the intermediate (**II**) eventually reverts to the (*E*)-isomer.

Table 2. Substituent and solvent effects on the composition of each compound obtained by 1.0 and 5.0 h irradiations of the starting (*Z*)-**1a–e** (5.0×10^{-3} mol dm⁻³) in a given solvent

Compound	Irradiation time /h	Composition (%)				
		(<i>Z</i>)- 1	(<i>E</i>)- 1	2	3	4
1a ^a	1.0 (5.0)	45.9 (3.2)	38.2 (4.5)	10.3(71.7)	5.1(18.4)	0.4 (2.1)
1b ^a	1.0 (5.0)	49.8 (4.7)	37.9 (6.6)	8.0(68.6)	3.8(17.5)	0.4 (2.7)
1c ^a	1.0 (5.0)	52.1(27.6)	40.0(37.6)	0 (0)	6.4(25.8)	1.4 (9.0)
1d ^a	1.0 (5.0)	82.2(51.9)	17.1(38.6)	0 (0)	0.7 (4.6)	0 (4.9)
1e ^a	1.0 (5.0)	65.4(11.8)	27.3(11.7)	6.0(64.1)	0 (1.7)	1.3(10.8)
1a ^b	1.0 (5.0)	70.0(38.0)	24.3(26.3)	3.6(25.3)	2.0(10.5)	0 (0)

^a In MeOH. ^b In MeCN.

The previous finding^{3b} that intramolecular cyclization in the excited-state (*Z*)-*N*-benzoyl-*α*-dehydrophenylalanines leading to isoquinoline derivatives is almost completely suppressed (owing to stereoelectronic effects of the bulky benzoyl group) to selectively give 1-azetines *via* the (*E*)-isomer provides a good explanation for the increased composition of the azetine (**4e**) (10.8%) as compared to that of **4a** (2.1%, 5.0 h irradiation).

An analysis of solvent effects on the reactivity of **1a** shows that the excited-state reactivities of both isomers are much higher in the protic polar solvent, MeOH, than in the aprotic polar solvent, MeCN, which lowers not only the relative rate for the *Z*–*E* isomerization (1.0 h irradiation, Table 2) but also the product-composition ratio **2a**/(**2a**+**3a**+**4a**) (77.8–70.7% at 5.0 h irradiation). The fact that a tertiary amine undergoes hydrogen-bonding solvation to decrease its electron-donating ability,⁴ therefore, confirms that this solvation of the excited singlet-state (**1a**) by MeOH accelerates the *Z*–*E* isomerization and then greatly

enhances the electron-accepting ability of the naphthylmethylene moiety.

Although there are several synthetic routes to lactams fused to an aromatic ring,⁷ no convenient photochemical route to substituted dihydroquinolinones is known. The procedure for preparing the starting **1** is very simple and is easily applicable to its related compounds carrying less bulky alkyl substituents on the carboxamide side chain. Taking into consideration the fact that the benzoquinolinone derivatives (**2**) obtained are photochemically very stable, we are led to conclude that the photoreaction of substituted α -dehydro(1-naphthyl)alanines (**1a–e**) in the presence of TEA presents a new method for constructing the quinolinone skeleton and also offers significant information regarding the mechanism for the novel photocyclization reactions of **1** via ET.

REFERENCES AND NOTES

1. P. S. Mariano and J. L. Stavinoha, 'Synthetic Organic Photochemistry,' ed. by W. M. Horspool, Plenum Press, New York, 1984, pp. 145–257.
2. F. D. Lewis and G. D. Reddy, *J. Am. Chem. Soc.*, 1989, **111**, 6465; F. D. Lewis, G. D. Reddy, and D. M. Bassani, *J. Am. Chem. Soc.*, 1993, **115**, 6468; F. D. Lewis, D. M. Bassani, and G. D. Reddy, *J. Org. Chem.*, 1993, **58**, 6390; F. D. Lewis, G. D. Reddy, D. M. Bassani, S. Schneider, and M. Gahr, *J. Am. Chem. Soc.*, 1994, **116**, 597; F. D. Lewis, D. M. Bassani, E. L. Burch, B. E. Cohen, J. A. Engleman, G. D. Reddy, S. Schneider, W. Jaeger, P. Gedeck, and M. Gahr, *J. Am. Chem. Soc.*, 1995, **117**, 660.
3. a) K. Kubo, S. Yaegashi, K. Sasaki, T. Sakurai, and H. Inoue, *Tetrahedron Lett.*, 1996, **37**, 5917; b) K. Kubo, M. Koshiba, H. Hoshina, and T. Sakurai, *Heterocycles*, 1998, **48**, 25.
4. T. Sakurai, K. Miyoshi, M. Obitsu, and H. Inoue, *Ber. Bunsenges. Phys. Chem.*, 1996, **100**, 46; K. Miyoshi, K. Kubo, T. Sakurai, and H. Inoue, *Nippon Kagaku Kaishi*, 1999, 37 and references cited therein.
5. Y. S. Rao and R. Filler, *Synthesis*, 1975, 749; B. Rzeszotarska, J. Karolak-Wojciechowska, M. A. Broda, Z. Galdecki, B. Trzezwinska, and A. E. Koziol, *Int. J. Peptide Protein Res.*, 1994, **44**, 313.
6. Selected data for (*Z*)-**1a**: mp 187.0–188.5 °C (EtOH-hexane); IR (KBr) ν/cm^{-1} = 3306, 1660, 1625, 1550; ¹H NMR (500 MHz, DMSO-*d*₆) δ = 0.90 (3H, t, *J* = 7.3 Hz), 1.33 (2H, tq, *J* = 7.3, 7.6 Hz), 1.48 (2H, tt, *J* = 6.7, 7.6 Hz), 1.83 (3H, s), 3.17 (2H, dt, *J* = 6.4, 6.7 Hz), 7.46 (1H, s), 7.50–7.57 (4H, m), 7.89 (1H, d, *J* = 7.9 Hz), 7.93–7.96 (2H, m), 8.06 (1H, t, *J* = 6.4 Hz), 9.20 (1H, s); ¹³C NMR (DMSO-*d*₆) δ = 13.7, 19.5, 22.6, 31.2, 38.8, 123.7, 124.1, 125.4, 125.9, 126.2(2C), 128.2, 128.3, 131.0, 131.3, 132.7, 133.1, 164.6, 169.3. Anal. Calcd for C₁₉H₂₂N₂O₂: C, 73.52; H, 7.14; N, 9.03%. Found: C, 73.54; H, 7.02; N, 9.13%.
For (*E*)-**1a**: mp 148.0–149.0 °C (EtOAc-hexane); IR (KBr) ν/cm^{-1} = 3250, 1650, 1629, 1539; ¹H NMR (500 MHz, DMSO-*d*₆) δ = 0.66 (3H, t, *J* = 7.3 Hz), 0.88 (2H, tq, *J* = 7.3, 7.6 Hz), 1.06 (2H, tt, *J* = 7.0, 7.3 Hz), 2.00 (3H, s), 2.87 (2H, dt, *J* = 6.1, 7.0 Hz), 7.36–7.40 (2H, m), 7.43 (1H, s), 7.52 (2H, m), 7.75 (1H, t, *J* = 6.1 Hz), 7.77–7.79 (1H, m), 7.89 (1H, d, *J* = 7.3 Hz), 7.98 (1H, d, *J* = 7.6 Hz), 9.69 (1H, s); ¹³C NMR (DMSO-*d*₆) δ = 13.6, 19.4, 23.5, 30.3, 38.3, 112.6, 124.6, 125.3, 125.6, 125.8, 126.0, 127.1, 128.2, 131.3, 132.6, 133.1, 135.1, 164.4, 168.6. Anal. Calcd for C₁₉H₂₂N₂O₂: C, 73.52; H, 7.14; N, 9.03%. Found: C, 73.29; H, 7.19; N, 9.14%.
For **2a**: mp 157.0–158.0 °C (EtOH); IR (KBr) ν/cm^{-1} = 3308, 1668, 1642; ¹H NMR (500 MHz, DMSO-*d*₆) δ = 0.89 (3H, t, *J* = 7.3 Hz), 1.53 (2H, m), 1.95 (3H, s), 2.96 (1H, dd, *J* = 14.3, 15.6 Hz), 3.64 (1H, dd, *J* = 6.1, 15.6 Hz), 3.95–4.14 (2H, m), 4.56 (1H, ddd, *J* = 6.1, 7.9, 14.3 Hz), 7.45 (1H, dd, *J* = 7.6, 7.9 Hz), 7.51 (1H, d, *J* = 8.2 Hz), 7.55 (1H, dd, *J* = 7.6, 8.2 Hz), 7.91 (1H, d, *J* = 8.5 Hz),

7.92 (1H, d, $J = 8.5$ Hz), 8.00 (1H, d, $J = 8.2$ Hz), 8.34 (1H, d, $J = 7.9$ Hz); ^{13}C NMR (DMSO- d_6) $\delta = 13.7, 19.4, 22.6, 27.0, 29.4, 41.7, 48.1, 116.2, 118.2, 123.1, 124.6, 127.1, 128.1, 128.3, 129.6, 130.8, 136.0, 168.2, 169.4$. Anal. Calcd for $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_2$: C, 73.52; H, 7.14; N, 9.03%. Found: C, 73.24; H, 7.13; N, 9.11%.

For **3a**: mp 61.5–62.5 °C (EtOAc-hexane); IR (KBr) $\nu/\text{cm}^{-1} = 3382, 1668, 1530$; ^1H NMR (500 MHz, DMSO- d_6) $\delta = 0.94$ (3H, t, $J = 7.3$ Hz), 1.37 (2H, tq, $J = 7.3, 7.6$ Hz), 1.60 (2H, tt, $J = 7.0, 7.6$ Hz), 3.04 (3H, s), 3.41 (2H, dt, $J = 6.1, 7.0$ Hz), 7.81–7.85 (2H, m), 8.12 (1H, d, $J = 9.2$ Hz), 8.13 (1H, m), 8.17 (1H, d, $J = 9.2$ Hz), 8.83 (1H, t, $J = 6.1$ Hz), 8.92 (1H, m), 9.12 (1H, s); ^{13}C NMR (DMSO- d_6) $\delta = 13.7, 19.6, 22.6, 31.5, 38.6, 113.0, 122.8, 123.8, 126.2, 127.9, 128.65, 128.71, 129.1, 129.5, 132.9, 134.6, 144.4, 157.0, 164.1$. Anal. Calcd for $\text{C}_{19}\text{H}_{20}\text{N}_2\text{O}$: C, 78.05; H, 6.89; N, 9.58%. Found: C, 77.85; H, 7.09; N, 9.74%.

7. I. Ninomiya, C. Hashimoto, T. Kiguchi, and T. Naito, *J. Chem. Soc., Perkin Trans. 1*, 1985, 941; K. Jones, M. Thompson, and C. Wright, *J. Chem. Soc., Chem. Commun.*, 1986, 115; G. P. Johnson and B. A. Marples, *J. Chem. Soc., Perkin Trans. 1*, 1988, 3399; B. E. Ali, K. Okuro, G. Vasapollo, and H. Alper, *J. Am. Chem. Soc.*, 1996, **118**, 4264.