

SYNTHESIS AND REACTIONS OF THIABENZENES

Mikio Hori, Tadashi Kataoka, Hiroshi Shimizu,
Sachio Ohno, Ken'ichi Narita, and Harumi Aoki
Gifu College of Pharmacy
Mitahora, Gifu 502

In order to elucidate the structure of thiabenzenes, synthesis of new cyclic sulfur 1,4-ylides and their reactions with electrophiles have been studied: from 9-cyano and 9-ethoxycarbonyl-10-methylthioxanthenium salts (I and II) and base stable 9-cyano and 9-ethoxycarbonyl-10-methylthiaanthracenes (III and IV) have been synthesized. The conformation of I and II was considered by NMR spectral data.

The treatment of III (example 1) with dimethyl acetylenedicarboxylate gave 9-cyano-9-methyl-(V), 9-cyano-9-(cis-1,2-dimethoxycarbonylvinyl)-(VI), 9-cyano-9-(1,2-dimethoxycarbonylcyclopropyl)-(VII), and 9-cyano-9-(1,2-dimethoxycarbonylallyl)thioxanthene (VIII). By the reaction of III and tetracyanoethylene, V, 9-cyano-9-(dicyanomethyl)-(IX), 9-dicyanomethylidenethioxanthene (X), malononitrile (XI), and 2,2,3,3-tetracyanobutane (XII) were isolated as main products. Mechanisms of these reactions were also discussed.

In the course of the studies, the synthesis, stereochemistry and reactivity of sulfoxides, sulfones and 10-alkyl or arylsulfonium salts derived from 9-pentafluorophenyl-(XIII), 9-mesityl-(XIV), and 9-phenyl(d_5)thioxanthene (XV) were also investigated.