

A NOVEL SYNTHESIS OF THE PHENANTHROINDOLIZIDINE
ALKALOID RING SYSTEM FROM PHENANTHRENE


Seichi Takano,* Kohtaro Yuta, and Kunio Ogasawara

Pharmaceutical Institute, Tohoku University, Aobayama, Sendai,
Japan

A novel synthesis of the phenanthroindolizidine alkaloid ring system(16) from phenanthrene(2) is described.

In connection with our recent finding on the efficient synthesis of phenanthrene derivatives¹, its utility for the construction of the phenanthroindolizidine alkaloids² was examined by synthesizing the ring system(16) from phenanthrene(2). Since there has been no example starting from 9,10-unsubstituted phenanthrene derivatives in the synthesis of the phenanthroindolizidine alkaloids, e.g., tylophorine(1)³, the attempt seemed to be worthwhile.

Chart 1




(1) R=OMe

(16) R=H

Following Billups' method⁴, 7,7-dichlorodibenzo[*a,c*]bicyclo-[4.1.0]heptane(3)⁵, derived from phenanthrene(2) and dichlorocarbene, was treated with potassium *ter*.butoxide(2eq.) in tetrahydrofuran at 0° to give an oily 9-(chloro-2-tetrahydrofurylmethyl)-phenanthrene(5) in good yield *via* the chlorocarbene insertion to the solvent(Chart 2). Dehydrochlorination of 5 was effected by boiling with pyridine in dimethylformamide(1:8) to give the enol ether(6) as a stereoisomeric mixture which,


Chart 2



without separation, on hydrolysis with ethanolic hydrochloric acid(2%) at room temperature afforded the ketol(7), mp 110°, ir(nujol) 3300, 1695 cm^{-1} , NMR(CDCl_3) (δ) 1.75(3H, m, disapp. 1H with D_2O), 3.47(2H, t, J=6Hz), 3.52(2H, t, J=7Hz), 4.09(2H, s), 7.45-7.95(7H, m), 8.50-8.75(2H, m), Mass(m/e) 278(M^+), in 30.1% yield from 5. The ketol(7) was converted to the benzylamino-alcohol(9), an oil, ir(neat) 3300 cm^{-1} , NMR(CDCl_3) (δ) 1.5-1.9 (4H, m), 3.15-4.25(8H, m, disapp. 1H with D_2O), 7.2-7.8(12H, m),

8.5-8.8(2H, m), 9.8-10.2(1H, br.s, disapp. with D₂O), in 55% yield, by treating with benzylamine, followed by reduction with sodium borohydride. After treatment of 9 with thionyl chloride in chloroform, the crude chloride(10) was stirred with ethanolic potassium carbonate at room temperature to give the N-benzylpyrrolidine derivative(11), an oil, NMR(CDCl₃) (δ) 1.5-1.8(4H, m), 2.8-4.2(7H, m), 7.2-8.02(12H, m), 8.45-8.75(2H, m), Mass(m/e) 351(M⁺), in 91% from 9. The carbamate(12), an oil, ir(neat) 1680cm⁻¹, NMR(CDCl₃) (δ) 1.6-2.1(4H, m), 2.7(1H, m), 3.5(2H, br.t), 4.0-4.5(2H, m), 5.2(2H, s), 7.15-7.9(12H, m), 8.5-8.75(2H, m), converted from 11 with carbobenzoxy chloride in the presence of potassium hydrogen carbonate in boiling chloroform⁶, was heated at 120° in ethanolic hydrochloric acid(5%) to produce the secondary amine(13), mp 205-207°, ir(nujol) 3300cm⁻¹, NMR(CDCl₃) (δ) 1.4-2.1(4H, m), 2.84(1H, br.s, disapp. with D₂O), 2.6-3.6(5H, m), 7.4-8.2(7H, m), 8.5-8.73(2H, m), in total yield 73.6% from 11. Standard formic acid treatment of 13 yielded the formamide(14), an oil, which on cyclization with phosphorus oxychloride, followed by reduction with sodium borohydride, gave the required ring system(2,3,6,7-tetrademethoxytylophorine)(16), mp 167°(lit⁷., 170°), NMR(CDCl₃) (δ) 1.5-2.2(4H, m), 2.3-2.7(3H, m), 2.9-3.4(2H, m), 3.65(1H, d, J=18Hz), 4.7(1H, d, J=18Hz), 7.4-8.1(6H, m), 8.5-8.8(2H, m), Mass(m/e) 273(M⁺), 204(base peak), in 72% yield from 13.

Chart 3



The synthesis just described would be promising to the conversion of 9,10-unsubstituted phenanthrene derivatives into the phenanthroindolizidine alkaloids. Application of this method to the synthesis of an antitumor phenanthroindolizidine alkaloid, tylophorine(1), is currently being carried out in this laboratory.

REFERENCES

- 1 To be published elsewhere.
- 2 T. Kametani, 'The Chemistry of the Isoquinoline Alkaloids' Vol. 2, The Sendai Institute of Heterocyclic Chemistry, Sendai, 1974, pp. 289-293.
- 3 T.R. Govindachari, M.V. Lakshmikantham, and S. Rajadurai, Tetrahedron, 1961, 14, 284; R.B. Herbert and C.J. Moody, Chem. Comm., 1970, 121; B. Chauncy and E. Gellert, Austral. J. Chem., 1970, 23, 2503.
- 4 W.E. Billups, L.P. Lin, and W.Y. Chow, J. Amer. Chem. Soc., 1974, 96, 4026.
- 5 G.C. Joshi, N. Singh, and L.M. Pande, Synthesis, 1972, 317.
- 6 K.C. Rice, J. Org. Chem., 1975, 40, 1850.
- 7 T.R. Govindachari, M.V. Lakshmikantham, K. Nagarajan, and B.R. Pai, Tetrahedron, 1958, 4, 311.

Received, 6th February, 1976