




## A SYNTHESIS OF APORPHINE DERIVATIVES

Shinzo Kano<sup>\*</sup>, Youko Takahagi, Eiji Komiyama,  
Tsutomu Yokomatsu, and Shiroshi Shibuya  
Tokyo College of Pharmacy, 3-20-1, Kitashinjuku,  
Shinjuku-ku, Tokyo, Japan

A variety of 1-hydroxyaporphines have been synthesized via 6a,7-dehydroaporphines, obtained by the reaction between a series of 1-halogenobenzyl-3,4-dihydro-2-methylisoquinolinium iodides and dimethylsodium.

We have studied the benzyne reaction of 1-(3-bromo-4-methoxybenzyl)-1,2,3,4-tetrahydro-7-hydroxy-6-methoxy-2-methylisoquinoline (1) using dimethylsodium as a base to give 12a-methyldibenz[b,g]indolizine (2). In this reaction, formation of neither the aporphine (3) nor the morphinandienone (4) was observed<sup>1</sup>. We widely investigated a synthesis of aporphine derivatives via 6a,7-dehydroaporphines, obtained by the reaction of 1-halogenobenzyl-3,4-dihydro-2-methylisoquinolinium salts with dimethylsodium, as an extension of the previous works<sup>1,2</sup>, although a synthesis of the 6a,7-dehydroaporphine (5) was studied<sup>3</sup>.


















1-(2-Bromo-4,5-methylenedioxybenzyl)-3,4-dihydro-7-hydroxy-6-methoxy-2-methylisoquinolinium iodide (6) was treated with dimethylsodium in DMSO at room temperature to give the 6a,7-dehydroaporphine (7), mp 178-179° (from methanol), in 25 % yield. Reduction of (7)



with zinc amalgam in a mixture of 50 % acetic acid and conc. hydrochloric acid (1:1) yielded ( $\pm$ )-domesticine (8), mp 185-186° (lit.<sup>4</sup> mp 186-187°), in quantitative yield, which was identical with the authentic specimen<sup>4</sup>. Similarly, the reaction of 1-(3-bromo-4-methoxybenzyl)-3,4-dihydro-7-hydroxy-6-methoxy-2-methylisoquinolinium iodide (9) with dimethylsodium, followed by reduction of (10) with zinc amalgam, afforded the aporphine (3), mp 199-200° (from methanol), which showed  $\delta$  (CDCl<sub>3</sub>), 2.50 (NCH<sub>3</sub>), 3.88, 3.92 (6H, 2xOCH<sub>3</sub>), 6.50 (3-H), 6.78 (d,d, J=2, 11 Hz, 10-H), 7.22 (d, J=11 Hz, 8-H), 8.08 (d, J=2 Hz, 11-H), m/e 311 (M<sup>+</sup>, C<sub>19</sub>H<sub>21</sub>NO<sub>3</sub>), 310, 294. Gibson<sup>5</sup> also examined the benzyne reaction of (1) to obtain (3) by the use of sodium amide, but unsuccessful.

Furthermore, the aporphine derivatives possessing a substituent at the 5-position were synthesized by the same methods. The reaction of the quaternary salts (11), (12), (13) with dimethylsodium, followed by reduction of the dehydroaporphines (14), (15), (16) yielded the corresponding 5-methylaporphines (17), mp 165-167° (from methanol-ether) [ $\delta$  (CDCl<sub>3</sub>) 0.93 (d, J=6 Hz, 5-CH<sub>3</sub>), 2.52 (NCH<sub>3</sub>), 3.80, 3.85, 3.88 (9H, 3xOCH<sub>3</sub>), 6.45 (3-H), 6.73 (8-H), 8.07 (11-H), m/e 355 (M<sup>+</sup>, C<sub>21</sub>H<sub>25</sub>NO<sub>4</sub>), 354, 338], (18), mp 196-198° (from methanol) [ $\delta$  (CDCl<sub>3</sub>) 0.98 (d, J=6 Hz, 5-CH<sub>3</sub>), 2.55 (NCH<sub>3</sub>), 3.90 (OCH<sub>3</sub>), 5.93 (OCH<sub>2</sub>O), 6.52 (3-H), 6.73 (8-H), 7.95 (11-H), m/e 339 (M<sup>+</sup>, C<sub>20</sub>H<sub>21</sub>NO<sub>4</sub>), 338, 322], and (19), mp 182-183° (from methanol) [ $\delta$  (CDCl<sub>3</sub>) 0.97 (d, J=6 Hz, 5-CH<sub>3</sub>), 2.57 (NCH<sub>3</sub>), 3.87, 3.90 (6H, 2xOCH<sub>3</sub>), 6.58 (3-H), 6.78 (d,d, J=2, 11 Hz, 9-H), 7.20 (d, J=11 Hz, 8-H), 8.87 (d, J=2 Hz, 11-H), m/e 325 (M<sup>+</sup>, C<sub>20</sub>H<sub>23</sub>NO<sub>3</sub>), 324, 308], respectively.

Since the aporphine (17) synthesized by this method was identical



with the product obtained by photolysis<sup>6</sup> of (21) which was prepared from the 3,4-dihydroisoquinoline (20) by Gal's method<sup>7</sup>, the relative configuration of 5-CH<sub>3</sub> and 6a-H would be trans.

In the carbon-13 nmr spectroscopies of these aporphines in deuteriochloroform, the assignments of the chemical shifts, based on splitting patterns recorded under off-resonance decoupled conditions and on the comparison of the spectra<sup>8</sup>, are shown in Table I. The C<sub>6a</sub> and NCH<sub>3</sub> signals shifted to the higher region by introducing a substituent at the 5-position.

Table 1 Carbon-13 Chemical Shifts of Aporphines<sup>9</sup>

(All shifts are in parts per million from TMS)

Carbon	Compound				
	(8)	(17)	(18)	(19)	(22) <sup>3</sup>
C-1	140.73	140.79	140.86	141.32	140.86
C-2	145.83	146.23	145.89	146.03	146.03
C-3	109.73	109.67	109.67	110.33	109.01
C-3a	123.64	122.38	122.38	122.38	123.64
C-4	28.81	35.30	34.83	35.23	29.01
C-5	53.31	55.63	55.43	55.63	53.64
N-CH <sub>3</sub>	43.91	40.86	40.53	40.79	44.04
C-6a	62.45	54.44	54.44	54.24	62.78
C-7	34.94	34.30	34.57	34.64	34.70
C-7a	125.76	125.17	126.03	127.35	125.10
C-8	108.74	112.65	109.27	128.34	112.58
C-9	145.83	147.81	146.09	112.52	147.81
C-10	145.83	147.35	146.09	158.34	147.35
C-11	108.15	111.26	108.34	114.11	111.26
C-11a	130.20	129.54	130.60	133.31	129.01
C-11b	127.22	126.49	126.49	128.94	127.35
C-6b	119.47	119.54	119.40	119.43	119.74
5-CH <sub>3</sub>		10.07	10.20	9.93	
OCH <sub>3</sub>	56.03	56.03(x3)	56.09	56.03(x2)	55.96 56.16(x2)
OCH <sub>2</sub> O	100.73		100.79		

## REFERENCES

1. S. Kano, T. Yokomatsu, N. Yamada, K. Matsumoto, S. Tokita, and S. Shibuya, Chem. and Pharm. Bull.(Japan), 1974, 22, 1607.
2. S. Kano, E. Komiyama, T. Ogawa, Y. Yakahagi, T. Yokomatsu, and S. Shibuya, Chem. and Pharm. Bull.(Japan), 1975, 23, 2058.
3. T. Kametani, S. Shibuya, and S. Kano, J. C. S. Perkin I, 1973, 1212.
4. T. Kametani, S. Shibuya, K. Kigasawa, M. Hிரagi, and O. Kusama, J. Chem. Soc. (C), 1971, 2712.
5. M. S. Gibson, G. W. Prenton, and J. M. Walthew, J. Chem. Soc. (C), 1970, 2234.
6. T. Kametani, K. Fukumoto, S. Shibuya, H. Nemoto, T. Nakano, T. Sugahara, T. Takahashi, Y. Aizawa, and M. Toriyama, J. C. S. Perkin I, 1972, 1435.
7. J. Gal, R. J. Wienkam, and N. Castagnoli, Jr., J. Org. Chem., 1974, 39, 418.
8. G. C. Levy and G. L. Nelson, "Carbon-13 Nuclear Magnetic Resonance for Organic Chemist", John Wiley & Sons., New York, 1972.
9. The spectra were taken in deuteriochloroform with Varian NV-14 spectrometer operating at 15.1 MHz.

Received, 28th January, 1976