

SYNTHESIS OF s-TRIAZOLO[1,5-b]ISOQUINOLINES.

NEW STABLE o-QUINONOID HETEROCYCLES




Yasumitsu Tamura,* Midori Iwaisaki, Yasuyoshi Miki,
and Masazumi Ikeda

Faculty of Pharmaceutical Sciences, Osaka University,
133-1, Yamada-kami, Suita, Osaka, Japan

The synthesis and some properties of new stable o-quinonoid heterocycles, s-triazolo[1,5-b]isoquinolines, are described.

During the last ten years, interest in the syntheses and chemistry of o-quinonoid heterocycles has been noticeable.¹ This class of compounds, particularly ones lacking ring substituents, are usually unstable and sometimes isolated only as their stable N-phenylmaleimide adducts. We now wish to report the synthesis of new stable o-quinonoid heterocycles, s-triazolo[1,5-b]isoquinolines (3).²

Treatment of 3-acetamidoisoquinoline (1a) with an equimolar quantity of O-mesitylenesulfonylhydroxylamine (MSH)³ in methylene chloride at room temperature for 10 min gave 74% yield of 2a, m.p. 134°. ⁴ Heating 2a in polyphosphoric acid (PPA) at 80° for



1 hr gave 3a, m.p. 182-183°, in 76% yield, which was stable even in refluxing xylene for 10 hr. The structure of 3a was determined from its spectral data [nmr δ (CDCl₃) 2.70 (s, 3H, CH₃), 7.3-8.0 (m, 4H, arom. H), 8.10 (bs, 1H, H-10), and 9.30 (bs, 1H, H-5); uv λ_{\max} . (EtOH) 238 (log ϵ 4.61), 250 sh (4.38), 346 sh (3.41), 361 (3.52), and 378 nm (3.38)] and chemical evidence. Refluxing 3a with N-phenylmaleimide or N-methylmaleimide in benzene gave the corresponding cycloadducts 4, m.p. 298° [ir ν_{\max} . (KCl) 1780, 1705 cm⁻¹; nmr δ (CDCl₃) 2.39 (s, 3H, CH₃), 3.58 (dd, 1H, J_{bc}=9 Hz, J_{cd}=3 Hz, H_c), 3.82 (dd, 1H, J_{bc}=9 Hz, J_{ab}=4 Hz, H_b), 5.15 (d, 1H, J_{cd}=3 Hz, H_d), 6.17 (d, 1H, J_{ab}=4 Hz, H_a), 6.4-6.6 (m, 2H, arom. H), 7.2-7.5 (m, 7H, arom. H)] and 5, m.p. 261.5-263° [ir ν_{\max} . (KCl) 1780, 1700 cm⁻¹; nmr δ (CDCl₃) 2.30 (s, 3H, CH₃), 2.57 (s, 3H, CH₃), 3.39 (dd, 1H, J_{bc}=9 Hz, J_{cd}=3 Hz, H_c), 3.62 (dd, 1H, J_{bc}=9 Hz, J_{ab}=4 Hz, H_b), 5.01 (d, 1H, J_{cd}=3 Hz, H_d), 6.03 (d, 1H, J_{ab}=4 Hz, H_a), 7.0-7.3 (m, 4H, arom. H)], in quantitative yields.

Similarly, reaction of 3-formamidoisoquinoline (1b), m.p. 138°, with MSH gave 2b, m.p. 130-131°, in 82% yield, which was cyclized with PPA to 3b, m.p. 160-161°, in 57% yield [nmr δ (CDCl₃) 7.3-8.0 (m, 4H, arom. H), 8.22 (bs, 1H, H-10), 8.58 (s, 1H, H-2), 9.38 (bs, 1H, H-5); uv λ_{\max} . (EtOH) 235 (log ϵ 4.67), 249 sh (4.34), 345 sh (3.45), 360 (3.59), 378 nm (3.47)]. This compound was rather unstable and gradually decomposed in refluxing benzene.

The thermal stability of compounds 3 may be attributed, at least in part, to the contribution of the mesoionic structures, e.g., 6.



REFERENCES

1. D.W.H. MacDowell, A.T. Jeffries, M.B. Meyers, J. Org. Chem., 1971, 36, 1416, and references cited therein; M.P. Cava, N.M. Pollack, O.A. Mamer, and M.J. Mitchell, ibid., 1971, 36, 3932; M.J. Haddadin and N.C. Chelhot, Tetrahedron Letters, 1973, 5185.
2. Although a U.S. patent has claimed the synthesis of a pyrrolo[1,2-b]isoquinoline, we were unable to repeat this observation. [R.H. Sprague, U.S.P., 2622082 (Dec. 16, 1952)].
3. Y. Tamura, J. Minamikawa, and M. Ikeda, Synthesis, 1977, 1.
4. All new compounds gave satisfactory elemental analyses.

Received, 20th April, 1977