


SYNTHESIS OF METHYL (R,S)-LICHENSTERINATE¹Seppo I. Pennanen

Department of Chemistry, Helsinki University of Technology,
02150, Otaniemi, Finland.

The methyl ester of (R,S)-lichensterinic acid was synthesized from methyl α -ketopalmitate and 1-diethylaminopropyne in three steps in 58% overall yield.


Recently a new synthesis of the 3-methyl-2(5H)-furanoid structural unit (1) from α -epoxyketones and 1-diethylaminopropyne was reported from this laboratory.² A modification of this method in the synthesis of a natural product is now reported.

(R,S)-Lichensterinic acid¹ (2, R = H) has been isolated from some lichen species such as Icelandic moss (*Cetraria islandica*)^{3,4,5} and its synthesis has been reported involving the isomerization^{3,5,6} of protolichensterinic acid (3), another lichen constituent. Both 2 and 3 show antibacterial activity toward Gram positive organisms.⁵



Insofar β -epoxy- α -ketoesters are not very readily available, it was reasoned that the furanoid ether oxygen of 1 could perhaps be introduced at a later stage of synthesis, allowing α -ketoester to be used as the starting material. Thus, using methyl α -ketopalmitate⁷ (4) only three synthetic steps were required to convert

it to 2 (R = Me), the two last steps being carried out in one pot (Scheme 1).



Scheme 1.

The ynamine 5 reacted smoothly with the keto group of 4 giving via the oxetene rearrangement the expected product 6 in 86% yield. The allylic Wohl-Ziegler bromination⁸ (methylene attack preferred to methyl) of 6 led to the intermediate 7 which was not isolated, but the reaction mixture was treated directly with aqueous sodium bicarbonate. Hydrolysis of the reactive allylic bromide⁹ occurred, followed by spontaneous lactonization and expulsion of diethylamine. After purification a pale yellow, viscous oil was obtained. According to the spectral data the oil was identified as the expected (R,S)-mixture of 2 (R = Me). Asano and Kanematsu⁴ give the melting point of pure R-form of 2 (R = Me) as 53-4°C. The yield was 67% and thus the overall yield was 58%.

ExperimentalCompound 6:²

To a mixture of 200 mg (0.7 mmol) of methyl α -ketopalmitate⁷ and 0.10 ml (0.7 mmol) of 1-diethylaminopropyne¹⁰ in 10 ml of dry ether 200 mg (1.1 mmol) of dry magnesium bromide was added and the reactants were stirred under argon for 30 mins. During this time a very viscous syrup developed on the walls of the reaction flask. Water was added and the ether solution was dried with Na_2SO_4 followed by the evaporation of ether. The product 6 was isolated from the residue by preparative TLC (silica gel; elution with EtOAc / CHCl_3 , 1:9). The yield was 238 mg (86%) of a pale yellow, viscous oil.

δ : 3.63 (3H, s), 3.24 (4H, q, 7 Hz), 2.30 (2H, t, 7 Hz),
1.90 (3H, s), 1.32 (22H, broad s), 1.10 (6H, t, 7 Hz),
1.00 (3H, t, 7 Hz).

m/e: 395 (M^+).

Compound 2 (R = Me):

100 mg (0.25 mmol) of the adduct 6 and 50 mg (0.28 mmol) of NBS in 5 ml of CCl_4 were refluxed for 3 hrs.⁸ The mixture was cooled and CCl_4 was evaporated under reduced pressure. The residue was dissolved into 10 ml of H_2O / THF (1:2) and 200 mg of NaHCO_3 was added and the mixture was refluxed for 5 hrs.⁹ After evaporation of the solvents the product 2 was isolated by preparative TLC (silica gel; elution with EtOAc / CHCl_3 , 1:9). The yield was 57 mg (67%) of a pale yellow, viscous oil.

δ : 3.70 (3H, s), 3.68 (1H, t, 7 Hz), 1.90 (3H, broad s),
1.30 (24 H, broad s), 1.00 (3H, t 7 Hz).

m/e: 338, 323 ($\text{M}^+ - 15$; base peak).

The author wishes to thank Professor T. Hase for his kind discussion.

References and footnotes

1. Systematic name: Methyl 2,5-dihydro-4-methyl-5-oxo-2-tridecyl-3-furoate.
2. Pennanen S.I., Tetrahedron Lett., 1977, 2631.
3. Asano M. and Kanematsu T., J. Pharm. Soc. Japan, 1931, 51, 390.
4. Asano M. and Kanematsu T., Chem. Ber., 1932, 65B, 1175.
5. Cavallito C.J., Fruehauf D.M. and Bailey J.H., J. Am. Chem. Soc., 1948, 70, 3724.
6. Bloomer J.L., Eder W.R. and Hoffman W.F., J. Chem. Soc. (C), 1970, 1848.
7. Kuwata T., J. Am. Chem. Soc., 1938, 60, 559.
8. Greenwood F.L., Kellert M.D. and Sedlak J., Org. Synth., Coll. Vol. IV, 1963, 108 and references therein.
9. Burgstahler A.W. and Nordin I.C., J. Am. Chem. Soc., 1961, 83, 198.
10. Commercially available.

Received, 17th April, 1978