


REACTIONS OF BENZO-1,2-DITHIOLE-3-THIONE WITH N,N'-
DIALKYL-1,2-DIAMINOETHANES. SYNTHESIS OF o-THIO-
QUINONEMETHIDES WITH A KETENE AMINAL GROUP

Renji Okazaki, Kyung-Tae Kang, and Naoki Inamoto*

Department of Chemistry, Faculty of Science, The University
of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan


Benzo-1,2-dithiole-3-thione (2) reacted with N,N'-dialkyl-
1,2-diaminoethanes (3) to give o-thioquinonemethides with
a ketene aminal group (4-6) in high yields, which were
found to be of considerable betain nature significantly
different from previously known o-thioquinonemethides
with a ketene thioacetal group (1).

Synthesis of o-thioquinonemethides (1) from the photo-
reaction of 1,2-dithiole-3-thiones with olefins has recently been
reported by de Mayo¹ and by us.²





We now describe another type of o-thioquinonemethides (4)
with a ketene aminal group which show significantly different
properties from 1.

Reaction of benzo-1,2-dithiole-3-thione (2) with N,N'-dialkyl-1,2-diaminoethane (3) in ethanol afforded 4 in high yields. In the reaction of 2 with 3a, the formation of hydrogen sulfide (91% as PbS) and sulfur (94%) was also confirmed. Thioquinonemethides (5) (78%) and (6) (85%) were also prepared in a similar way from the corresponding 1,2-dithiole-3-thiones. Compound 7 (65%) was synthesized by the reaction of more reactive dithiolium salt (8)³ with N,N'-dimethyl-o-phenylene-diamine using ion exchange resin as base.



a: R=Me reflux, 4h 99% mp 153-5°C
 b: R=Et reflux, 6h 64% mp 148-9°C
 c: R=i-Pr 160 °C, 30h 78%
 (sealed tube)



The electronic spectra of 4b⁴ showed no concentration dependence, indicating it exists as monomer in solution. This is in marked contrast with 1 which is in equilibrium with a dimer.^{1,2} The NMR spectra of 4-7 are very similar to each other, implying other thioquinonemethides synthesized here are also monomeric in solution. Of particular interest is the isolation and stability of 5 since compounds with 2,3-naphthoquinone structure are usually very unstable⁵ and isolable only in special cases.⁶ The NMR chemical shifts (CD₃OD) of N-methyl protons of 4a, 5, and 6 (δ 2.90-2.92) are almost the same as that of 9 (δ 2.92; prepared from 4a and methyl iodide) and those of ethylene protons (δ 4.00-4.06) are fairly close to that of 9 (δ 4.14), indicating the large contribution of ionic canonical structure (e.g. 10 for 4) in these thioquinonemethides. This is reflected in the solubility of these compounds; for instance, 4a is very soluble in water and alcohols but only slightly soluble in chloroform and benzene.

The importance of the ionic contribution 10 to the ground state of 4 was also shown by solvent effect of the UV spectra of 4b.³ The more polar is the solvent, the shorter becomes the absorption maximum, implying the stabilization of the ionic ground state by polar solvents. Here again, the thioquinonemethides with a ketene aminal group are quite different from those with a ketene thioacetal group (1) whose electronic spectra show no essential solvent dependence.

Compound 4a reacted with dimethyl acetylenedicarboxylate and dibenzoylacetylene in acetonitrile at room temperature to give

[4+2] adduct (11), suggesting a possible route to six-membered sulfur heterocycles.



REFERENCES and NOTES

- 1 P. de Mayo and H.Y. Ng, J.C.S. Chem. Commun., 1974, 877; Can. J. Chem., 1977, 55, 3763.
- 2 R. Okazaki, and N. Inamoto, Chem. Lett., 1974, 1439; R. Okazaki, K. Sunagawa, M. Kotera, and N. Inamoto, Tetrahedron Lett., 1976, 3815.
- 3 J. Faust and R. Mayer, Justus Liebigs Ann. Chem., 1965, 688, 150.
- 4 Solvent, λ_{\max} (nm), log ϵ : H₂O, 252, 4.00; EtOH, 278, 4.18; MeCN, 299, 4.26; CH₂Cl₂, 299, 4.06; PhH, 307, 4.27.
- 5 J.E. Shields and J. Bornstein, Chem. Ind. (London), 1967, 1404; M.P. Cava, N.M. Pollack, O.A. Mamer, and M.J. Mitchell, J. Org. Chem., 1971, 36, 3932; Cf., G.J. Gleicher, D.D. Newkirk, and J.C. Arnold, J. Am. Chem. Soc., 1973, 95, 2526.
- 6 M.P. Cava and J.P. Van Meter, J. Org. Chem., 1969, 34, 538.

Received, 27th September, 1978