

OXOANLOBINE, A NEW OXOAPORPHINE ALKALOID FROM GUATTERIA MELOSMA


Charles H. Phoebe, Jr., Paul L. Schiff, Jr., Joseph E. Knapp, and David J. Slatkin*

Department of Pharmacognosy, School of Pharmacy, University of Pittsburgh, Pittsburgh, PA 15261 U.S.A.

Abstract - Oxoanolobine (1), a new alkaloid from an alcoholic extract of Guatteria melosma Diels (Anonaceae), was characterized as 1,2-methylenedioxy-9-hydroxyoxoaporphine (10-hydroxy-8H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-8-one) by physicochemical data and conversion to lanuginosine (2).

An alcoholic extract (654 g) of Guatteria melosma¹ (27.2 kg) was submitted to an acid-base partition procedure. Subsequent silicic acid chromatography of the basic fraction afforded oxoanolobine (1) (50 mg) as an orange amorphous solid from methanol; mp 270-275° (dec); $[\alpha]_D^{28}$ 0° (c 1.0, MeOH);
 uv λ_{max}^{MeOH} nm 217 (log ϵ 4.24) 249 (4.43), 274 (4.35), 324(sh) (3.84), 370 (3.65) and 442 (3.76),
 $\lambda_{max}^{MeOH + 0.1N HCl}$ nm 220 (log ϵ 4.29) 260 (4.41), 287 (4.31), 345 (3.72), 395 (3.73) and 510 (3.51);
 $\lambda_{max}^{MeOH + 0.1N NaOH}$ nm 222 (log ϵ 4.29), 253 (4.38), 291 (4.42), 334(sh) (4.01), 372 (3.48) and 506 (3.42); ir ν_{max}^{KBr} cm⁻¹ 3420 (br), 1660(C=O). The nmr spectrum (60 MHz, TFA, TMS, δ in ppm) indicated the presence of one methylenedioxy group at 6.60 (2H,s), a C-3 aromatic proton at 7.47 (1H,s) an aromatic AB system for C-4 at 8.37 (1H,d,J=6Hz) and C-5 at 8.65 (1H,d,J=6Hz) and an aromatic AMX system for C-8, C-10 and C-11 at 7.98 (1H,d,J=2.5Hz), 7.59 (1H,dd,J=8,2.5Hz) and 8.70 (1H,d,J=8Hz), respectively. The ms showed a M⁺ at m/e 291 (100%) for C₁₇H₉NO₄, 263(8), 233(15) and 178(10) with metastable ions at m/e 237.3 for the transition 291→263 (m*_{calc} 237.7) and 206.4 for the transition 263→233 (m*_{calc} 206.4). These spectral data indicated that 1 was 1,2-methylenedioxy-9-hydroxyoxoaporphine.

Treatment of oxoanolobine (1) with ethereal diazomethane gave an O-methyl derivative (lanuginosine) (2) as yellow needles from methanol; mp 314°; $[\alpha]_D^{28}$ 0° (c 1.0, MeOH); ms M⁺ m/e 305(100%), 304(56), 290(5), 276(20), 275(40) and 234(16). O-Methylxoanolobine was identical (ir, uv, ms, mp) with authentic lanuginosine², thus confirming that 1 was 1,2-methylenedioxy-9-hydroxyoxoaporphine (10-hydroxy-8H-benzo[g]-1,3-benzodioxolo[6,5,4-de]quinolin-8-one)³.



- 1 R = OH
 2 R = OCH₃
 3 R = H

Oxoaporphine alkaloids have been found in several plant families in addition to the Anonaceae. These include the Araceae, Hernandiaceae, Lauraceae, Magnoliaceae, Menispermaceae, Monimiaceae, Papaveraceae, Ranunculaceae and Eupomatiaceae.^{4,5} Although liriodenine (3) has reported broad spectrum antimicrobial activity^{6,7} and activity against 9-KB tissue culture cells⁸, little is known about the pharmacological activity of other members of this group of alkaloids.

Acknowledgements and References

1. The authors express their appreciation to Drs. Norman R. Farnsworth and Geoffrey A. Cordell, School of Pharmacy, University of Illinois at the Medical Center, Chicago, Illinois for providing the plant material used in this study. The stem bark was collected in Peru in 1974 and a herbarium specimen is deposited at the National Arboretum USDA Acc. No. PR-80719; ID No. CA-2202.
2. The authors are grateful to Drs. Andre Cave and Michel Leboeuf, Laboratoires de Matière Médicale, Center 'Etudes Pharmaceutiques, Besancon, France for spectra and an authentic sample of lanuginosine.
3. The authors are grateful to Kurt L. Loening, Nomenclature Director, Chemical Abstracts Service, Columbus, Ohio for providing the correct IUPAC name for oxoanolobine.
4. M. Shamma, "The Isoquinoline Alkaloids", Academic Press, New York, 1972, p. 245.
5. M. Shamma and J.L. Moniot, "Isoquinoline Alkaloids Research", Plenum Press, New York, 1978, p. 173.
6. C.R. Chen, J.L. Beal, R.W. Doskotch, L.A. Mitscher and G.H. Svoboda, *Lloydia*, 1974, 37, 493.
7. C.D. Hufford, M.J. Funderburk, J.M. Morgan and L.W. Robertson, *J. Pharm. Sci.*, 1975, 64, 789.
8. P.E. Sonnet and M. Jacobson, *J. Pharm. Sci.*, 1971, 60, 1254.

Received, 30th June, 1980