

STRUCTURE OF MULBERROFURAN C, A NATURAL HYPOTENSIVE DIELS-ALDER
ADDUCT FROM ROOT BARKS OF THE CULTIVATED MULBERRY TREE
(MORUS BOMBYCIS KOIDZUMI)

Taro Nomura^{*,a)} Toshio Fukai^{a)} Junko Matsumoto^{a)} Kazutaka Fukushima^{b)}
and Yasunori Momose^{c)}

- a) Faculty of Pharmaceutical Sciences, Toho University,
2-2-1, Miyama, Funabashi-Shi, Chiba 274, Japan
- b) Research Institute for Chemobiodynamics, Chiba University,
1-8-1, Inohana-Cho, Chiba-Shi, Chiba 280, Japan
- c) Faculty of Medicine, Toyama Medical and Pharmaceutical University,
2630, Sugitani, Toyama-Shi, Toyama 930-01, Japan

A new 2-arylbenzofuran derivative with a fused dihydro-chalcone partial moiety was isolated from root barks of the cultivated mulberry tree (Morus bombycis Koidzumi) and named mulberrofuran C. The structure was shown to be I on the basis of spectral data. Mulberrofuran C (I) is regarded biogenetically as a Diels-Alder adduct of a chalcone derivative and dehydromoracin C (II) or its equivalent. Intravenous injection of I (1 mg/Kg) produced a significant hypotension in rabbit.

In previous communications,^{1,2} we reported that two new flavone derivatives having a hypotensive effect, kuwanon G (=moracenin B³=albanin F,⁴ III) and H (moracenin A⁵=albanin G,⁴ IV), were isolated from root barks of the cultivated mulberry tree (a variety of Morus alba L.), and the structures were shown to be III and IV, respectively. Further survey for hypotensive constituents has led to isolation of a novel hypotensive derivative of 2-arylbenzofuran, mulberrofuran C, from mulberry root barks (a variety of Morus bombycis Koidzumi). In this paper, we report the isolation and structure determination of mulberrofuran C.

The dried root bark of the cultivated mulberry tree (a variety of *Morus bombycis* Koidzumi) was extracted successively with hexane, benzene, and ethyl acetate. The ethyl acetate extract was fractionated sequentially by the polyamide column chromatography and by the preparative thin layer chromatography over silica gel to give chalcomoracin (V)^{6,7} and a new 2-arylbenzofuran derivative, mulberrofuran C (I)⁸ in $2.7 \times 10^{-3}\%$ and $2.5 \times 10^{-3}\%$ yield, respectively. The compound (I) showed a marked hypotensive effect (1 mg/Kg, i.v.) to rabbit.

Mulberrofuran C (I), amorphous powder,⁹ $[\alpha]_D^{18} +153^\circ$ (c=0.33 in methanol), gave the FD-MS spectrum which showed the molecular ion peak at m/e 580, and the ¹³C nmr spectrum which indicated the presence of thirty-four carbons [nine aliphatic carbons (1xCH₃-, 1x-CH₂-, 3x>CH-, 1x>C=CH, 1x-CH=C-O-), twenty-four aromatic carbons (11xCH, 5xC, 8xC-O) and one carbonyl carbon]. These spectra suggested composition of mulberrofuran C to be C₃₄H₂₈O₉. Treatment of I with dimethyl sulfate and potassium carbonate in acetone (reflux, 9 h) effected exhaustive methylation to give its heptamethyl ether (Ia), amorphous powder,^{9a} $[\alpha]_D^{14} +175^\circ$ (c=0.12 in methanol), FeCl₃ test (negative). The molecular formula of Ia was determined to be C₄₁H₄₂O₉ by the high-resolution mass spectrum (m/e 678.2826, M⁺), and hence mulberrofuran C (I) could be formulated as C₃₄H₂₈O₉. The compound (I) showed the following color reactions: Mg-HCl test (negative), FeCl₃ test (brown), and showed the following spectra: ir $\nu_{\max}^{\text{Nujol}} \text{ cm}^{-1}$: 3350, 1625, 1620; uv $\lambda_{\max}^{\text{MeOH}} \text{ nm}(\log \epsilon)$: 215(4.75), 281(4.43), 320(4.62), 333(4.56); $\lambda_{\max}^{\text{MeOH+AlCl}_3}$: 220(4.80), 316(4.73), 360(sh 4.03). While the uv spectrum was similar to that of chalcomoracin (V),⁶ comparison of the spectrum of I with that of moracin M (VI)¹⁰




Fig. 1 The uv spectra of mulberrofuran C




Fig. 2 The uv spectra















Chart 1


revealed the presence of an extra absorption at ~ 280 nm in the former (I) which would be ascribed to a conjugated carbonyl group³ (Figs. 1,2). Moreover, the absorption at 280 nm induced a bathochromic shift in the presence of AlCl_3 . Sherif *et al.* reported that when a prenyl group was located *ortho* to a chelated hydroxyl group no AlCl_3 induced shift was observed in the uv spectra.¹¹ These data, together with the fact that chalcomoracin (V) coexists with mulberrofuran C (I), led us to presume that I would probably be a deprenylchalcomoracin.

The mass spectra of I and Ia showed the following fragments: I, m/e 580(M^+),¹² 562,¹² 470,¹² 452,¹² 320, 308(VII), 242(VIII), 110(IX); Ia,¹³ 678(M^+), 513 ($\text{C}_{31}\text{H}_{29}\text{O}_7$), 350(VIIa), 297($\text{C}_{18}\text{H}_{17}\text{O}_4$), 284(VIIIa), 165(X), 151($\text{C}_9\text{H}_{11}\text{O}_2$). On the other hand, the mass spectrum of chalcomoracin (V) showed the fragments at m/e 648(M^+),¹² 630,¹² 470,¹² 388, 308(VII), 242(VIII), 178(XI), 110(IX). These mass spectra also suggested that mulberrofuran C (I) would be a Diels-Alder adduct such as chalcomoracin (V),⁶ regarded as a cycloaddition product with the chalcone and the 2-arylbenzofuran derivative. This was substantiated by detailed analysis of the ^1H nmr spectrum (200 MHz, CD_3COCD_3) using sequential decoupling and by comparison of the ^1H nmr spectrum of chalcomoracin (V).⁶ The chemical shifts (δ) and coupling constants (Hz) of protons of the relevant cyclohexene ring are shown in Fig. 3, while the remaining protons are summarized as follows: protons in a 2-arylbenzofuran moiety, 6.96(2H,s, C_2 - and C_6 -H), 6.78-6.86(3H,m, C_3 -, C_5 -, and C_7 -H), 7.41(1H,d,J=8.5, C_4 -H); aromatic protons in a 2,4-dihydroxybenzoyl moiety, 6.27(1H,d,J=2.5, C_{11} "-H), 6.40(1H,dd,J=2.5 and 9, C_{13} "-H), 8.56(1H,d,J=9, C_{14} "-H), 12.61(1H,s, C_{10} "-OH); aromatic protons in a 2,4-dihydroxyphenyl moiety, 6.34(1H, dd,J=2.5 and 8.5, C_{19} "-H), 6.55(1H,d,J=2.5, C_{17} "-H), 7.04(1H,d,J=8.5, C_{20} "-H). All these results indicate that the structure of mulberrofuran C is possibly represented by I or I' (except the stereochemistry).

Detailed examination of the ^1H nmr spectrum of I with that of V,⁶ revealed that the chemical shifts and coupling constants of protons of the relevant cyclohexene ring of I resembled closely those of V as shown in Figs. 3 and 4. This strongly suggested that mulberrofuran C (I) and chalcomoracin (V) have the same disposition concerning location of the dihydroxyphenyl and dihydroxybenzoyl moieties on the cyclohexene ring and also the same relative configuration. The ^{13}C nmr spectra of I and V were also measured, in which all the carbon atoms were assigned (Table 1) by comparison with the ^{13}C nmr spectra of model compounds, benzofuran derivatives,^{8a,14,15} and kuwanon G¹ and H.² In the ^{13}C nmr spectrum



Flg. 3 (I)


 Fig. 4 (V)⁶

of I, the signal of the C-11" appeared at a higher applied magnetic field than that of the C-11" of V, whereas the chemical shift values of all the carbon atoms other than except the C-11" were essentially the same as to those of the relevant carbon atoms of V (Table 1).

 Table 1 ¹³C nmr chemical shifts in DMSO-d₆

	I	V		I	V		I	V
C-2	155.33* s	155.40*	C-1"	133.37 s	133.21	C-15"	122.27 s	122.04
C-3	102.70 d	102.89	C-2"	128.09 d	127.90	C-16"	157.14* s	155.21*
C-3a	121.27 s	121.19	C-3"	33.63 d	33.05	C-17"	101.47 d	102.62
C-4	121.54 d	121.56	C-4"	47.07 d	46.88	C-18"	155.90* s	155.21*
C-5	112.60 d	112.53	C-5"	33.28 d	33.86	C-19"	106.40 d	107.17
C-6	153.94* s	153.83*	C-6"	34.51 t	33.86	C-20"	133.37 d	131.17
C-7	97.69 d	97.65	C-7"	23.38 q	23.24	C-21"		21.03
C-7a	157.14* s	155.83*	C-8"	207.45 s	207.68	C-22"		122.04
C-1'	129.24 s	129.17	C-9"	113.41 s	114.99	C-23"		130.36
C-2'	103.01 d	103.16	C-10"	164.61 s	162.42	C-24"		25.38
C-3'	155.33* s	156.95*	C-11"	102.70 d	114.37	C-25"		17.56
C-4'	112.83 s	112.83	C-12"	164.61 s	162.03			
C-5'	155.33* s	156.95*	C-13"	108.13 d	106.21			
C-6'	103.01 d	103.16	C-14"	128.47 d	131.17			

* Assignments may be reversed.

This result establishes that mulberrofuran C (I) is deprenylchalomoracin.

Mulberrofuran C (I) is optically active and the second example of a natural

product which is considered to be formed by a Diels-Alder type of enzymatic reaction process of a chalcone derivative and dehydromoracin C (II) or its equivalent.⁶ The 2-arylbenzofuran derivatives were isolated as phytoalexins from diseased mulberry tree.^{6,16} It is interesting that mulberrofuran C is a hypotensive constituent of *Morus* root bark.

ACKNOWLEDGEMENT We are grateful to Dr. M. Takasugi, Faculty of Science, Hokkaido University, for a gift of spectral data and authentic sample of chalcomoracin, and grateful to Prof. Dr. S. Sakai, Faculty of Pharmaceutical Sciences, Chiba University, for mass spectrum measurement, and also grateful to Dr. S. Urano, Tokyo Metropolitan Institute of Gerontology, for ¹H nmr spectrum (200 MHz) measurement. We also thank to Mr. T. Ohmori for his technical assistance.

REFERENCES AND FOOTNOTES

- 1 T. Nomura and T. Fukai, Chem. Pharm. Bull., 1980, 28, 2548.
- 2 T. Nomura, T. Fukai, and T. Narita, Heterocycles, 1980, 14, 1943.
- 3 Y. Oshima, C. Konno, H. Hikino, and K. Matsushita, Tetrahedron Letters, 1980, 21, 3381.
- 4 M. Takasugi, S. Ishikawa, S. Nagao, T. Masamune, A. Shirata, and K. Takahashi, Chem. Letters, 1980, 1577.
- 5 Y. Oshima, C. Konno, H. Hikino, and K. Matsushita, Heterocycles, 1980, 14, 1287.
- 6 M. Takasugi, S. Nagao, T. Masamune, A. Shirata, and K. Takahashi, Chem. Letters, 1980, 1573.
- 7 Although this compound could not be isolated as crystalline form, it was indistinguishable (only tlc and ¹H nmr) from chalcomoracin which was sent from Dr. M. Takasugi, Hokkaido University.
- 8 Mulberrofuran A and B described in the references, 8a and 8b, respectively.
 - a) T. Nomura, T. Fukai, J. Uno, and T. Arai, Heterocycles, 1978, 9, 1593.
 - b) T. Nomura and T. Fukai, Planta medica, in press.
- 9a) Although only one spot was detected on tlc, this compound could not be isolated as crystalline form.
- b) Elemental analysis gave a following result: Anal. Calcd. for C₃₄H₂₈O₉·H₂O: C, 68.22; H, 5.06. Found: C, 67.88; H, 5.02.

- 10 M. Takasugi, S. Nagao, L. Muñoz, S. Ishikawa, T. Masamune, A. Shirata, and K. Takahashi, Abstracts of papers, "22nd Symposium the Chemistry of Natural Products", Fukuoka, October 24, 1979, p 275.
- 11 C.A. Sherif, R.K. Gupta, and M. Krishnamurti, Tetrahedron Letters, 1980, 21, 641.
- 12 This fragment could not be detected in EI-MS and only detected in FD-MS.
- 13 The formulae of the fragment ions of this compound were supported by the high-resolution mass spectrometry.
- 14 T. Okuyama and T. Fueno, Bull. Chem. Soc. Jpn., 1974, 47, 1263.
- 15 M. Komatsu, I. Yokoe, and Y. Shirataki, Chem. Pharm. Bull., 1978, 26, 1274.
- 16 M. Takasugi, S. Nagao, T. Masamune, A. Shirata, and K. Takahashi, Tetrahedron Letters, 1979, 48, 4675 and refs cited therein.

Received, 9th February, 1981