

STUDIES ON FURAN DERIVATIVES. XIII.<sup>1</sup>

## THERMOLYSIS OF METHYL 5-(2-AZIDOPHENOXY)-2-FUROATE

Akira Tanaka\* and Toshinao Usui

Faculty of Pharmaceutical Sciences, Josai University, Keyakidai  
1-1, Saitama 350-02, Japan

**Abstract** — In thermolysis of methyl 5-(2-azidophenoxy)-2-furoate, the first conversion of this into 2-(2-methoxallylvinyl)benzoxazole has been found.

Meth-Cohn and co-workers<sup>2</sup> have proposed a new mechanism on the pyrrolo[2,1-*b*]-benzothiazole formations employing photolysis and thermolysis of 2-azidophenyl thienyl sulfides, which involves the cleavage of thiophene ring to  $\alpha,\beta$ -unsaturated thioketones via a five membered ring intermediate by intramolecular insertion of nitrene. They also have emphasized the propriety of this ring opening mechanism rather than the 1,4-cycloaddition pathway of nitrene already suggested in other related reactions.<sup>3</sup> Jones and Mckinley<sup>4</sup> have also assumed similar  $\alpha,\beta$ -unsaturated indolenyl ketone intermediates in the deoxygenation of ortho-nitrophenyl-di-(2-furyl)methanes to give diethyl 2-alkylfuro[3,2-*c*]carbazol-5-ylphosphates.

We now confirmed that the thermolysis of methyl 5-(2-azidophenoxy)-2-furoate (I) leads to the formation of 2-(2-methoxallylvinyl)benzoxazole (II) which corresponds to  $\alpha,\beta$ -unsaturated ketones (or thioketones) as mentioned above.


Compound (I), prepared from methyl 5-(2-nitrophenoxy)-2-furoate<sup>1</sup> by the usual method,<sup>5</sup> was allowed to thermolyse in ortho-dichlorobenzene at 160-170° for 3 hr under nitrogen. After cooling, an excess of solvent was removed and the residue was chromatographed on silica gel, eluting with benzene, to give II; yield 22%; m.p. 134-136° (from petroleum benzin); <sup>1</sup>H-N.M.R.  $\delta$  (acetone-*d*<sub>6</sub>): 3.89 (3H, s, CH<sub>3</sub>), 7.10-7.85 (6H, m, benzene ring and vinylene H); I.R.  $\nu$  (nujol): 1725, 1695 (C=O), and 1625 cm<sup>-1</sup> (C=N); mass spectrum m/e: 231 (40.5%), 203 (33), 172 (100), 144 (89), 116 (21.7), and 89 (23.4).

Hydrolysis of II in 5% sodium hydroxide solution gave 3-(2-benzoxazolyl)acrylic acid which was identified with a sample.<sup>6</sup>

From these data, the structure of II was assigned as 2-(2-methoxallylvinyl)benz-

oxazole. The elemental analysis of II supported a molecular formula of  $C_{12}H_9NO_4$ . Thus, we could be obtained the stable  $\alpha,\beta$ -unsaturated ketone (II) which corresponds to the ring opening intermediate proposed by Meth-Cohn et al.<sup>2</sup> and a new synthetic rout for a benzoxazole ring system was also provided. More recently,  $\alpha,\beta$ -unsaturated ketone and thioketone as minor components ( $\sim 1.5\%$ ) were isolated by Jones et al.<sup>7</sup> in the reaction of 2-nitrenophenyl-di-(5-t-butyl-2-thienyl)methane.

Scheme



#### References and Notes

1. Part XII of this series: A. Tanaka, T. Usui, and M. Shimadzu, J. Heterocyclic Chem., submitted.
2. J.M. Lindley, O. Meth-Cohn, and H. Suschitzky, J. Chem. Soc. Perkin I, 1978, 1198.
3. K. Hafner and W. Kaiser, Tetrahedron Letters, 1964, 2185; G.R. Cliff, G. Jones, and J.M. Woollard, J. Chem. Soc. Perkin I, 1974, 2072.
4. G. Jones and W.H. Mckinley, J. Chem. Soc. Perkin I, 1979, 599.
5. Unpublished results; m.p. 63.5-65.5°.
6. W. Ried and H. Keller, Chem. Ber., 1956, 89, 2578.
7. G. Jones, C. Keates, I. Kladko, and P. Radley, Tetrahedron Letters, 1979, 1445; P.C. Hayes, G. Jones, C. Keates, I. Kladko, and P. Radley, J. Chem. Res. (S), 1980, 288; idem, J. Chem. Res. (M), 1980, 3523-3535.

Received, 9th March, 1981