

STUDIES OF POTENTIAL HETEROCYCLIC SULFONES AS ANTIMICROBIAL AGENTS

SYNTHESIS AND ANTIMICROBIAL ACTIVITY OF SOME NOVEL SUBSTITUTED ARYLSULFONYLBENZO[h]QUINOLINE DERIVATIVES

RAJENDRA PRASAD BAHUGUNA, YOGESH CHANDRA JOSHI, MAHAVEER PRASAD DOBHAL AND BHUWAN CHANDRA JOSHI*

Department of Chemistry, University of Rajasthan, Jaipur

AND

HARI NARAIN MANGAL

Department of Mycrobiology, S.M.S. Medical College, Jaipur

Abstract - A series of substituted arylthiobenzo[h]quinoline (sul fides, Ia~If and IIa~IIc) synthesised earlier were oxidised (with 1% KMnO₄) to get their corresponding sulfones (IA~IF and IIA~IIC). All these compounds have been characterised by their elemental analysis, molecular weight determination (Rast Method) and spectral studies (IR and ¹H NMR). These compounds have been screened for antimicrobial activity against Gram positive Cocci and Gram negative Bacilli.

INTRODUCTION

A perusal of literature revealed that no work has so far been reported on aryl-sulfonylbenzo[h]quinolines. As such the syntheses and structural studies of these sulfones, hitherto unknown, would be of considerable importance. In our earlier communication we have synthesised arylthiobenzo[h]quinoline¹. However, the present communication describes the synthesis and structural elucidation of arylsulfonylbenzo[h]quinolines (sulfones).

Heterocyclic sulfones have been found to exhibit antimicrobial activity^{2,3}. Besides this, sulfones have also been used as antimalarials^{4,5} and antileprotic agents⁶. Realising the medicinal importance of sulfones and benzo[h]quinoline compounds, various arylsulfonylbenzo[h]quinoline compounds were successfully synthesised in this laboratory. These were tested for antimicrobial activity.

In spite of the number of methods known for oxidation of sulfides to sulfones the one involving the oxidation (with 1% KMnO_4)⁷ was found to be most elegant and convenient. Various arylthiobenzo[h]quinolines (Ia~If) and (IIa~IIc) were prepared and subsequently oxidised with 1% KMnO_4 to the corresponding arylsulfonylbenzo[h]quinolines (IA~IF and IIA~IIC). The structures of TLC pure sulfones (IA~IF and IIA~IIC) have been established through elemental analysis, molecular weight determination (Rast method, cf. Table I) and spectral studies (IR and ^1H NMR).

SPECTRAL STUDIES

The compounds showed characteristic absorption bands in IR spectra in the vicinity of $1600\text{--}1450\text{ cm}^{-1}$ along with characteristic absorptions bands at 1350 cm^{-1} , 1145 cm^{-1} , $1180\text{--}1070\text{ cm}^{-1}$ (for sulfonyl groups) and at $655\text{--}610\text{ cm}^{-1}$ (due to S-aryl vibration) were observed. This clearly indicates these compounds as aromatic heterocyclic sulfones.

By and large the presence of 10 protons were established in the aromatic range at $\delta 7.5$ to $\delta 8.4$. In the NMR spectra at $\delta 9.40$ a broad signal for 1H was accounted for the presence of proton at C-10. The shifting of this signal to downfield with a broad appearance was due to its partial bonding characteristic with the angular nitrogen. In IA one more proton (i.e. 11 H) was observed in the range of $\delta 7.8$ to $\delta 8.2$ since there was no substituent in the aromatic ring of arylsulfonyl group. The presence of C- CH_3 , O- CH_3 and NH_2 groups are further established in these compounds on the basis of their NMR spectra (cf. Table II).

ANTIBACTERIAL ACTIVITY

The antibacterial activity of nine compounds was studied against Gram positive Cocci i.e. Staphylococcus aureus 'a', Streptococcus faecalis 'b' and Gram negative Bacilli i.e. Escherichia coli 'c' Klebsiella pneumoniae 'd' and Pseudomonas aeruginosa 'e' in 200 ug/disc (cf. Table III).

After comparing the antimicrobial activity of the compounds with various bacteria the following conclusions were drawn:

1. The substitution of arylsulfonyl group at p-position to nitrogen favours the activity as the compounds having arylsulfonyl group at o-position to nitrogen.
2. In arylsulfonyl moiety the substitution does not favour the activity. IA and IIA are found to be most active against almost all bacteria.

TABLE I

Com- pound	Name of the Compound	Yield in %	m.p. °C	Rf	Analysis		M.W. (Rast Method)	-SO ₂ absor- ptions
					Calcd.	Found		
IA	2-(Benzenesulfonyl)- 4-methylbenzo[h]quino- line.	77	140	0.69	S=9.60 N=4.20 For C ₂₀ H ₁₅ O ₂ NS	9.42 4.04	328.5	1350, 1145
IB	2-(<i>o</i> -Carboxyl-benzene- sulfonyl)-4-methyl- benzo[h]quinoline.	79	170	0.76	S=8.44 N=3.71 For C ₂₁ H ₁₅ O ₄ NS	8.00 3.40	371.5	1350, 1140
IC	2-(<i>o</i> -Amino-benzene- sulfonyl)-4-methyl- benzo[h]quinoline.	75	184	0.58	S=9.19 N=8.04 For C ₂₀ H ₁₆ O ₂ N ₂ S	9.10 8.00	342.2	1350 1145
ID	2-(<i>o</i> -Nitro-benzene- sulfonyl)-4-methyl- benzo[h]quinoline.	78	192	0.59	S=8.46 N=7.40 For C ₂₀ H ₁₄ O ₄ N ₂ S	8.20 7.10	372.7	1360, 1140
IE	2-(<i>p</i> -Methylbenzene- sulfonyl)-4-methyl- benzo[h]quinoline.	72	122	0.60	S=9.22 N=4.03 For C ₂₁ H ₁₇ O ₂ NS	9.00 4.00	342.5	1360, 1145
IF	2-(<i>p</i> -Methoxybenzene- sulfonyl)-4-methyl- benzo[h]quinoline.	76	146	0.62	S=8.81 N=3.85 For C ₂₁ H ₁₇ O ₃ NS	8.60 3.60	368.5	1360 1150
IIA	4-(Benzenesulfonyl)- 2-methylbenzo[h]quino- line.	78	144	0.68	S=9.60 N=4.20 For C ₂₀ H ₁₅ O ₂ NS	9.50 4.00	329.2	1350 1145
IIB	4-(<i>o</i> -Carboxylbenzene- sulfonyl)-2-methyl- benzo[h]quinoline.	76	168	0.73	S=8.44 N=3.71 For C ₂₁ H ₁₅ O ₄ NS	8.00 3.40	372.1	1350, 1140
IIC	4-(<i>o</i> -Aminobenzene- sulfonyl)-2-methyl- benzo[h]quinoline.	73	172	0.59	S=9.19 N=8.04 for C ₂₀ H ₁₆ O ₂ N ₂ S	9.10 8.80	343.2	1350, 1145

TABLE II
(NMR; CDCl_3 ; Chemical shifts in δ values)

No.	Aromatic Protons	-C-CH ₃	Ar-CH ₃	Ar-OCH ₃	Ar-NH ₂
IA	7.8-8.2(m)-11H 9.40(bs)-1H	2.85(s)-3H	-	-	-
IB	7.90-8.3(m)-10H 9.40(bs)-1H	2.85(s)-3H	-	-	-
IC	7.7-8.2(m)-10H 9.40(bs)-1H	2.80(s)-3H	-	-	4.5(s)-2H
1D	7.90-8.4(m)-10H 9.40(bs)-1H	2.90(s)-3H	-	-	-
IE	7.7-8.4(m)-10H 9.40(bs)-1H	2.80(s)-3H	2.50(s)-3H	-	-
IF	7.5-8.3(m)-10H 9.40(bs)-1H	2.80(s)-3H	-	3.85(s)-3H	-
IIA	7.8-8.2(m)-11H 9.40(bs)-1H	2.85(s)-3H	-	-	-
IIB	7.90-8.30(m)-10H 9.40(bs)-1H	2.85(s)-3H	-	-	-
IIC	7.7-8.2(m)-10H 9.40(bs)-1H	2.80(s)-3H	-	-	4.5(s)-2H

m = multiplet, bs = broad singlet; s = singlet.

3. The substitution of $-\text{NO}_2$ group decreases the activity to minimum.
4. The activity is in the following order:
 $\text{IIA, IA} > \text{IIB} > \text{IIC, IC} > \text{IB, IE, IF} > \text{ID}$
5. The substitution of COOH group and NH_2 group decreases the activity but still the activity is more than the compounds having $-\text{CH}_3$ and $-\text{OCH}_3$ groups.
- The summary of the results are given in Table IV.

TABLE III

Sensitivity of *Staphylococcus aureus* (a), *Streptococcus faecales* (b), *Escherichia coli* (c), *Klebsiella pneumoniae* (d), *Pseudomonas aeruginosa* (e) against different substituted arylsulfonylbenzo h quinoline.

Compounds	Concentration of different compound used was 200ug/disc				
	a	b	c	d	e
IA	++	+++	++	++	+++
IB	+	+	+	+++	++
IC	+++	+	++	+	++
ID	+	+	+	+	+
IE	+++	++	+	+	+
IF	+	+	+++	++	++
IIA	+++	++	++	++	+++
IIB	+	+++	++	++	++
IIC	+++	+++	+++	+	+

6-8 mm (+); 9-12 mm (++), 13-16 mm (+++)

TABLE IV

Name of Bacteria	Maximum activity	Average activity	Least activity
a	IC, IE, IIA, IIC	IA	IB, ID, IF, IIB
b	IA, IIB, IIC	IE, IIA	IB, IC, ID, IF
c	IF, IIC	IA, IC, IIA, IIB	IA, IB, IE
d	IB	IA, IF, IIA, IIB	IC, ID, IE, IIC
e	IIA	IA, IB, IC, IF, IIB	ID, IE, IIC

The plates of culture medium were uniformly seeded with 18 hr broth culture of the bacterium to be tested. Small (6 mm) sterile discs of standard

filter papers, impregnated with standard solutions of test compounds in one strength (vide supra), were placed on plates of culture medium at proper spacing. One disc impregnated with solvent (used for the preparation of solutions) was also placed on culture plates as control. These culture plates were kept for 18-20 hr in an incubator at 37°C. The zone of inhibition of growth of bacteria produced by the diffusion of compounds from the disc into the surrounding medium was measured. Each test compound in one concentration was tested with ten strains of each organism.

EXPERIMENTAL


The melting points are uncorrected. The purity of compounds was checked by TLC using silica gel 'G' as stationary phase and benzene-ethanol-ammonia (7:2:1; v/v, upper layer) as mobile phase. Compounds were characterised by elemental analysis (sulfur and nitrogen) and spectral (IR and NMR) studies. IR spectra were recorded on Perkin Elmer-Infracord spectrometer in KBr pellets. NMR spectra were recorded on Varian-A-60 MHz spectrometer using CDCl₃ as solvent and TMS as an internal reference. The chemical shifts were expressed in δ values.

Substituted arylthiobenzo[h]quinoline derivatives (Ia~If and IIA~IIC) were synthesised by the method described earlier¹.

2-(Benzenesulfonyl)-4-methylbenzo[h]quinoline (IA)

To 1.50 g (0.005M) of 2-thiophenyl-4-methylbenzo[h]quinoline (Ia) dissolved in glacial acetic acid (15-20 ml), 1% KMnO₄ solution was added with occasional shaking as rapidly as decolorisation occurred. Excess of KMnO₄ solution was added with shaking followed by the addition of ethanol (ca. 50 ml). The reaction mixture was allowed to stand till the colour of KMnO₄ disappeared. The mixture was poured into ice cold water and filtered. The crude residue was exhaustively extracted with hot acetone and the solvent was evaporated off. The residue was crystallized from petroleum ether (bp 40-60°), when colourless shining crystals of IA were obtained.

On similar lines IB~IF and IIA~IIC were also synthesised from their corresponding sulfides (Ib~If and IIA~IIC) (cf. Chart I).



R	SULFIDES		SULFONES	
	I-SERIES	II-SERIES	I-SERIES	II-SERIES
	Ia	IIa	IA	IIA
	Ib	IIb	IB	IIB
	Ic	IIc	IC	IIC
	Id	-	ID	-
	Ie	-	IE	-
	If	-	IF	-

CHART-1

REFERENCES

1. R.P. Bahuguna, R. Sharma, and B.C. Joshi, Acta Ciencia Indica,
in press.
2. (a) F. Gialdi and R. Ponci, Farmaco (Pavia) Ed. Sci., 1957, 12, 370-4.
(b) G. Gialdi and R. Ponci, Farmaco (Pavia) Ed. Sci., 1959, 14, 288.
3. (a) T. Makikado, Japan Pat., 70, 25,896. Chem. Abstr., 1970, 73, 109707u;
(b) H. Johnston and S.H. Ructman, Ger. Offen, 1,911,023 (1970);
Chem. Abstr., 1971, 74, 31692j.
4. P.E. Thompson, B.J. Olszewski, and J.A. Waitz, Am. J. Trop. Med. Hyg.,
1965, 14, 343.
5. P.E. Thompson, A. Bayles, B.J. Olszewski, and J.A. Waitz, Am. J. Trop.
Med. Hyg., 1965, 14, 198.
6. E. Titus and J. Bernstein, Ann. N.Y. Acad. Sci., 1949, 52, 719.
7. N.D. Sharma, S. Kumar, P.V. Bakore, and B.C. Joshi, Croatica Chemica
Acta, CCACCA, 1977, 49, 517.

Received, 8th July, 1981