

CONDENSATION OF 2-HYDROXYTRYPTAMINE WITH
DIMETHYL 4-ETHYL-4-FORMYLPIMELATE

Esahak Ali, Pulak K. Chakraborty and
Satyesh C. Pakrashi*

Indian Institute of Chemical Biology,
Calcutta-700 032, India

Abstract - Condensation of 2-hydroxytryptamine (1) with dimethyl 4-ethyl-4-formylpimelate (2) yielded the stereoisomeric oxindoles, 8A, 8B, 8C, and 8D which were separated and their stereochemistry assigned. The compounds 8A-D or their N-methyl derivatives, 5A-D could be selectively reduced to the oxindole-amines 9 or 3 by treatment with phosphorus oxychloride followed by sodium borohydride but the amines were isomerised under the reaction condition.








The condensation of 2-hydroxytryptamine (1) with dimethyl 4-ethyl-4-formylpimelate (2) is a potentially useful route for the total synthesis of several indole alkaloids. In an attempted synthesis of the alkaloid vincatine (3)¹, the reaction of 1-methyl-2-hydroxytryptamine (4) with 2 was reported² to yield two stereoisomers of the oxindole 5. Condensation of 1 with 2 was later used for the total synthesis of (+) - vincadifformine (6)³ and (+) - quebrachamine (7)⁴. This reaction was also reported to yield two isomers of 8, the more polar one being amenable to cyclisation to vincadifformine³ or aspidospermidine⁴ skeleton. A recent publication⁵ by the French group prompted us to report the work carried out in this laboratory on the condensation of 1 with 2, which showed that the reaction yields all the four possible isomers of 8.

Reaction of 2-hydroxytryptamine (1) with the ester 2 in refluxing methanol in presence of acetate buffer (pH 4.6) yielded, as reported^{2,3} earlier, two products (ratio 6:4, total yield 76%) separable by thin layer or column chromatography over silica gel. The nmr spectra, however, indicated that each of them is a mixture of two compounds which could be resolved by reversed phase HPLC (μ -Bondapak C-18, MeOH:H₂O/2:1). We have also been able to successfully separate the four isomers 8A-D by fractional crystallisation, their relative proportion (by HPLC) being A:B:C:D = 18:22.5:32:27.5. The corresponding N-methyl derivatives 5A-D were prepared by methylation with MeI/NaH in DMF.


As expected, the uv spectra of all the compounds 8A-D and 5A-D were almost superimposable⁶. The mass spectra of the isomers 8A-D were also almost identical [for 8A, m/z 370 (M^+ , 100%), 341 (M-Et, 6%), 339 (M-OMe, 11%), 283 (M-CH₂CH₂COOMe, 6%), 225 (a, 16%), 212 (b, 23%), 196 (c, 39%), 159 (d, 59%) and 138 (e, 67%)], with appropriate mass shifts in the spectra of the N-methyl derivatives. The ir spectra of the compounds were also very close. Since the chemical shifts of the C-21 methine proton in all the eight compounds were observed in the narrow range of δ 4.12-4.20 (Table) all of them must have the same stereochemistry at the C/D ring junction, i.e. trans. The C-7 configuration of the isomers could be unambiguously deduced by comparison of the ¹³C chemical shifts of C-7 and C-9 with those reported for model compounds⁷. A distinction between the isomers 8A and 8B and between 8C and 8D could be made from the chemical shifts of the C-17 and C-19 which were expected to resonate at higher field in the axial isomer than in the corresponding equatorial isomer⁸. The shielding of the CH₃-CH₂- groups in the pmr spectra of 8B and 8C can then be explained as due to anisotropy of the aromatic ring.

The isolation of all the four possible isomers of 8 thus, disproves the earlier contention² that two of the isomers are unlikely to be produced because of steric interactions. We have also observed that the lactams 8A-D are not isomerised by acetic acid or pyridine at 100°C, the usual conditions for equilibration of oxindole alkaloids⁹.

For the synthesis of indole alkaloids, it is necessary to selectively reduce the δ -lactam functions of 5 and 8. Diborane or lithium aluminium hydride reduction of 5 and 8 have been reported to cleave the C₇-C₂₁ bond^{1,4}. Potassium borohydride reduction of a derivative of 9 was also reported⁴ to cleave this bond. We could, however, achieve the desired reduction by treatment of the lactams 5 and 8 with POCl₃ followed by sodium borohydride reduction¹⁰ when the oxindole amines 3 and 9 were produced in reproducible yields of 60-65%. However, the stereochemical integrity was not preserved during the reduction. In fact, reduction of any of the isomers of 5 or 8 produced apparently the same mixture of the isomers of 3 or 9. It could be confirmed that isomerisation of 5 and 8 did not take place during the POCl₃ treatment. Evidently, the different stereoisomers of 3 and 9 are equilibrated under the conditions of borohydride reduction. Thus, though the δ -lactam group in 5 and 8 could be selectively reduced, the stereospecific synthesis of vincatine (3) could not be achieved. That goal could, however, be reached by an alternative route reported in the succeeding communication⁸.



$\underline{5C}$, $\underline{8C}$ $\text{R}' = \text{CH}_2\text{CH}_2\text{COOMe}$, $\text{R}'' = \text{Et}$
 $\underline{5D}$, $\underline{8D}$ $\text{R}' = \text{Et}$, $\text{R}'' = \text{CH}_2\text{CH}_2\text{COOMe}$



$\underline{5A}$, $\underline{8A}$ $\text{R}' = \text{Et}$, $\text{R}'' = \text{CH}_2\text{CH}_2\text{COOMe}$
 $\underline{5B}$, $\underline{8B}$ $\text{R}' = \text{CH}_2\text{CH}_2\text{COOMe}$, $\text{R}'' = \text{Et}$

Table. Physical data of the lactams 8A-D and 5A-D

Compound	mp ($^{\circ}$ C)	ir (nujol), cm^{-1}	pmr (90/100 MHz), δ CDCl_3			
			CH_3CH_2	OMe	21-H	NH/NMe
8A*	122-124	3413,1705,1618	0.86	3.60	4.20	9.46
8B*	238-240	3100,1718,1613	0.63	3.63	4.20	9.56
8C**	236-238	3236,1720,1646,1616	0.74	3.56	4.16	9.43
8D**	256-258	3170,1720,1611	0.56	3.56	4.20	9.35
5A	126-128	1730,1706,1643,1613	0.83	3.60	4.16	3.30
5B	152-154	1730,1706,1643,1613	0.58	3.70	4.20	3.30
5C	198-200	1738,1693,1640,1613	0.74	3.52	4.12	3.22
5D	212-214	1724,1693,1635,1613	0.57	3.63	4.13	3.24

*Isolated from the less polar product. **Isolated from the more polar product.

REFERENCES AND FOOTNOTES

1. W. Döpke, H. Meisel, and H. W. Fehlhaber, Tetrahedron Letters, 1969, 1701.
2. L. Castedo, J. Harley-Mason, and M. Kaplan, J. Chem. Soc. Chem. Comm., 1969, 1444.
3. J. Y. Laronze, J. Laronze-Fontaine, J. Lévy, and J. Le Men, Tetrahedron Letters, 1974, 491.
4. V. S. Giri, E. Ali, and S. C. Pakrashi, J. Heterocycl. Chem., 1980, 17, 1133.
5. G. Hugel, J. Y. Laronze, J. Laronze, and J. Lévy, Heterocycles, 1981, 16, 581.
6. For 8A, λ_{max} . (EtOH) (Log ϵ): 209 (4.50), 256 (3.88) and 286 (3.20) nm.
7. E. Wenkert, J. S. Bindra, C. -J. Chang, D. W. Cochran, and F. M. Schell, Accounts Chem. Res., 1974, 7, 46.
8. E. Ali, P. K. Chakraborty, A. K. Chakravarty, and S. C. Pakrashi, Heterocycles, succeeding communication.
9. M. Shamma, R. J. Shine, I. Kompis, T. Sticzáy, F. Morsing, J. Poisson, and J. L. Pousset, J. Amer. Chem. Soc., 1967, 89, 1739.
10. A. Rahman, A. Basha, N. Waheed, and S. Ahmed, Tetrahedron Letters, 1976, 219.

Received, 6th October, 1981