

ON THE CONFIGURATION OF THE C<sub>5</sub>-SUBSTITUENTS OF LYCOPODIUM  
ALKALOIDS, 8-DEOXSERRATINIDINE, EPIDIHYDROFAWCETTIDINE,  
SERRATINIDINE, AND DIHYDROFAWCETTIDINE

Takashi Harayama,\* Tooru Taga, and Kenji Osaki

Faculty of Pharmaceutical Sciences, Kyoto University, Sakyo-ku,  
Kyoto 606, Japan


Kaoru Kuriyama

Shionogi Research Laboratories, Shionogi & Co., Ltd., Sagisu,  
Fukushima-ku, Osaka 553, Japan


**Abstract** — The configuration of the C<sub>5</sub>-substituents of  
8-deoxyserratinidine, epidihydrofawcettidine, serratinidine,  
and dihydrofawcettidine was determined as formulas 2, 3, 4,  
and 5, respectively, by <sup>1</sup>H-NMR, Brewster's benzoate rule, and  
an X-ray analysis of 8-deoxyserratinidine.

In previous papers, we reported the configurational establishment of the C<sub>3</sub>-C<sub>4</sub>  
bond of fawcettidine (1)<sup>1</sup> and the structure elucidation of 8-deoxyserratinidine  
(2)<sup>2</sup>, epidihydrofawcettidine (3)<sup>2</sup>, and serratinidine (4)<sup>3</sup>. However, the  
configuration of the C<sub>5</sub>-substituents of the latter three alkaloids had been left  
to be determined. In this communication, we describe the configurational  
establishment of the C<sub>5</sub>-substituents of these alkaloids and dihydrofawcettidine  
(5)<sup>3</sup>, which is a reduction product of 1 with NaBH<sub>4</sub> and is an epimer of 3 with  
respect to the C<sub>5</sub>-hydroxy group.


In the <sup>1</sup>H-NMR (60 MHz) spectrum in CDCl<sub>3</sub>-D<sub>2</sub>O, the signals due to a proton geminal  
to a C<sub>5</sub>-substituent were observed at δ 4.07 (W<sub>1/2</sub>=22 Hz) in 2, at δ 3.98 (W<sub>1/2</sub>=  
10 Hz) in 3, at δ 4.33 (W<sub>1/2</sub>=24 Hz) in 4<sup>4</sup>, and at δ 4.17 (W<sub>1/2</sub>=20 Hz) in 5,  
respectively. The half-band width suggests that the C<sub>5</sub>-substituents of 2, 4,  
and 5 have the same relative configuration. Since 3 and 5 possess the same  
absolute configuration<sup>2,3</sup> and have an epimeric relationship with respect to the  
C<sub>5</sub>-hydroxy group as mentioned above, the Brewster's benzoate rule<sup>5</sup> was applied



- (1)  $R^1=R^2=O, R^3=H$
- (2)  $R^1=R^3=H, R^2=NHAc$
- (3)  $R^1=OH, R^2=R^3=H$
- (4)  $R^1=H, R^2=NHAc, R^3=OH$
- (5)  $R^1=R^3=H, R^2=OH$
- (6)  $R^1=OCOC_6H_5, R^2=R^3=H$
- (7)  $R^1=R^3=H, R^2=OCOC_6H_5$



(6)



(7)

Chart 1




Fig. 1. Molecular Structure of  
8-Deoxyserratinidine (2)

in order to assume the configuration of the C<sub>5</sub>-substituents of both compounds. Then, 3 and 5 were benzoylated to give benzoylepидihydrofawcettidine (6)<sup>6</sup> and benzoyldihydrofawcettidine (7)<sup>6</sup>, respectively. The Brewster's benzoate rule was applied to 3 and 6 [ $\Delta[M]_D^{20} - 32.3^\circ = [M]_D^{20} + 333.5^\circ (\text{benzoate}) - [M]_D^{20} + 365.8^\circ (\text{OH})$ ], and to 5 and 7 [ $\Delta[M]_D^{20} + 176.2^\circ = [M]_D^{20} + 496.3^\circ (\text{benzoate}) - [M]_D^{20} + 320.1^\circ (\text{OH})$ ]. As can be seen from the Newman's projections of the surroundings of the bond between O and C<sub>5</sub> of 6 and 7 in chart 1, benzoylepидihydrofawcettidine, which shows a negative  $\Delta[M]_D$ , can be represented by formula 6, and benzoyldihydrofawcettidine, which shows a positive  $\Delta[M]_D$ , can be represented by formula 7. These assumptions were also supported by the results derived from the benzoate sector rule<sup>7</sup> applied to 6 and 7.<sup>8</sup> Therefore, the stereostructures of 8-deoxyserratinidine, serratinidine, and dihydrofawcettidine can be represented by formulas 2, 4, and 5, respectively, since these three bases possess the same configuration with respect to the C<sub>5</sub>-substituent.

Next, in order to confirm these assumptions, 8-deoxyserratinidine (2) was subjected to an X-ray analysis. Compound (2) crystallized from ether in the monoclinic, space group P2<sub>1</sub>, with two molecules per unit cell of dimensions a=12.783(3), b=9.364(2), c=7.326(2) Å, β=104.96(2)°, Z=2, and Dx=1.131 g/cm<sup>3</sup>. A total of 1017 unique reflections having Fo>3σ(Fo) were measured on a Rigaku AFC-5 diffractometer using CuKα radiation. The structure was solved by the direct method using the program MULTAN-78<sup>9</sup> and refined by the full-matrix least-squares method to R=0.066. The molecular structure of 2 so derived is depicted in Fig. 1, proving the above-mentioned assumptions. Consequently, the stereostructures of 8-deoxyserratinidine, epидihydrofawcettidine, serratinidine, dihydrofawcettidine can be definitely represented by formulas 2, 3, 4, and 5, respectively.

**ACKNOWLEDGEMENTS** The authors wish to express their sincere thanks to Emeritus Professor Y. Inubushi of Kyoto University for his hearty encouragement and helpful discussions, and to Ministry of Education, Science and Culture (Japan) for Grant-in-Aid for Scientific Research.

#### REFERENCES

- 1) Y. Inubushi, T. Harayama, K. Yamaguchi, and H. Ishii, Chem. Pharm. Bull.,

1981, 29, 3418.

- 2) Y. Inubushi and T. Harayama, Yakugaku Zasshi, 1982, 102, 434.
- 3) H. Ishii, B. Yasui, R. Nishino, T. Harayama, and Y. Inubushi, Chem. Pharm. Bull., 1970, 18, 1880.
- 4) In our previous paper<sup>3</sup>, the relative configurational establishment of the C<sub>3</sub>-C<sub>4</sub> bond and the C<sub>5</sub>-NHAc group of serratinidine (4) had been left behind. However, it seems reasonable to assume that the C<sub>3</sub>-C<sub>4</sub> bond of 4 possesses the same configuration as that of 1, since fawcettidine (1), the stereostructure of which has been recently determined<sup>1</sup>, and 4 were derived from serratinine via the reductive product of the same type of enolate<sup>3</sup> (a reductive fission of the C<sub>4</sub>-N bond of serratinine).
- 5) a) J. H. Brewster, Tetrahedron, 1961, 13, 106. b) J. H. Brewster, J. Amer. Chem. Soc., 1959, 81, 5475, 5483, 5493.
- 6) Benzoylepidihydrofawcettidine (6) : m/e 351 (M<sup>+</sup>). IR (CHCl<sub>3</sub>),  $\nu$  1710, 1660, 1602, 1586. <sup>1</sup>H-NMR (60 MHz in CDCl<sub>3</sub>),  $\delta$  1.01 (3H, d, J=7 Hz, >CH-CH<sub>3</sub>), 5.08 (1H, m, W<sub>1/2</sub>=9 Hz, -CH-OCO), 5.53 (1H, d, J=3 Hz, olefinic proton), 7.33-8.20 (5H, m, aromatic protons).  $[\alpha]_D^{20}$ =+95.0° (c=1.0) in EtOH.  
Benzoyldihydrofawcettidine (7) : m/e 351 (M<sup>+</sup>). IR (CHCl<sub>3</sub>),  $\nu$  1705, 1658, 1600, 1581. <sup>1</sup>H-NMR (60 MHz in CDCl<sub>3</sub>),  $\delta$  1.00 (3H, d, J=7 Hz, >CH-CH<sub>3</sub>), 5.33 (1H, m, W<sub>1/2</sub>=20 Hz, -CH-OCO), 5.55 (1H, d, J=2 Hz, olefinic proton), 7.34-8.20 (5H, m, aromatic protons).  $[\alpha]_D^{20}$ =+141.4° (c=0.99) in EtOH.
- 7) N. Harada, H. Ohashi, and K. Nakanishi, J. Amer. Chem. Soc., 1968, 90, 7349.
- 8) The CD data  $[\theta](nm)$  in MeOH. 3 : +43800 (231) (c=1.74×10<sup>-3</sup>M), 5 : +56900 (233) (c=1.86×10<sup>-3</sup>M), 6 : +47900 (230) (c=0.99×10<sup>-3</sup>M), 7 : +56400 (235) (c=0.99×10<sup>-3</sup>M). Details will be discussed in a full paper.
- 9) P. Main, S. E. Hull, L. Lessinger, G. German, J-P. Declercq, and M. M. Woolson, MULTAN 78. A System of Computer Programmes for the Atomic Solution of Crystal Structures from X-ray Diffraction Data. University of York, England and Louvain-LaNeuve, Belgium, 1978.

Received, 1st March, 1984