




THE INFLUENCE OF ALKYL SUBSTITUENTS ON INTERNAL PHOTODIMERIZATION
OF PYRIMIDINE-6-AZAPYRIMIDINE DINUCLEOTIDE ANALOGUES

Jadwiga Jankowska, Henryk Koroniak, and Krzysztof Golankiewicz⁺
Faculty of Chemistry, Adam Mickiewicz University, 60-780 Poznań
Poland

Abstract - The new internal photodimers of dinucleotide analogues in which 5-alkyl derivatives of uracil and 6-azaauracil are connected with trimethylene chain have been obtained. The structure is discussed on the basis of spectroscopic data.

In the recent communication¹ we described the formation of cis-syn internal photodimers during UV irradiation of dinucleotide analogues in which uracil or thymine was connected with 6-azaauracil with a trimethylene bridge. The unusual properties of some protons at the azacyclobutane ring - according to ¹H-NMR spectra - inspired us to examine the influence of alkyl substituents in 5,5'-positions on the formation of such internal photodimers.



In order to reach some quantitative data on unbonding base-base interaction in compounds 1a, 1b and 1c the hypochromism values² have been calculated and compared with the proper pyrimidine-pyrimidine counterparts. The obtained results are presented in the Table 1.

Compound	% H
Ura(1(CH ₂) ₃ 1)Ura ²	1.2
Ura(1(CH ₂) ₃ 1)6-azaUra	-1.2
Thy(1(CH ₂) ₃ 1)Ura ³	2.6
Thy(1(CH ₂) ₃ 1)6-azaUra	0.3
5-EtUra(1(CH ₂) ₃ 1)Ura ³	4.5
5-EtUra(1(CH ₂) ₃ 1)6-azaUra	0.6
Thy(1(CH ₂) ₃ 1)Thy ²	7.0
Thy(1(CH ₂) ₃ 1)6-azaThy	5.1

Table 1. Values of hypochromism % H of some dinucleotide analogues.

It is evident from the data given in the Table 1 that the hypochromism of the analogues having 6-azauracil moiety is lower than their uracil counterparts. Nevertheless the differences are too small to be responsible of the different behaviour of these two series under UV irradiation in aqueous solution. The short time irradiation of compounds 1a, 1b and 1c in aqueous solution with $\lambda > 290$ nm led exclusively to photohydrates, whereas the same experiments carried out with pyrimidine-pyrimidine analogues gave as a result cyclobutane type photodimers^{2,3}. The prolongation of the irradiation resulted in complicated mixture of products as shown by TLC. Quite different results were obtained when acetone was used as solvent and sensitizer. The irradiation of compounds 1a, 1b and 1c in acetone with $\lambda > 290$ nm for 15 min gave exclusively the azacyclobutane photodimers 2a, 2b and 2c in high yields. The obtained photodimers showed only end UV absorption at $\lambda = 270$ nm as shown in the Figure 1. The photodimers 2a, 2b and 2c are unstable in aqueous solution and they decompose gradually to 1a, 1b and 1c, respectively /within 4 days 2a, 2b and 2c stored in water solution at room temperature completely reverses to 1a, 1b and 1c, respectively/. The short time irradiation of photodimers

2a, 2b and 2c with $\lambda = 254$ nm reverses the dimerization what is visualized by appearance of UV absorption at 270 nm /similarly to pyrimidine-pyrimidine analogues ^{2,3}/.






Figure 1. UV spectra of compounds 1c and 2c in water.

The ¹H-NMR and ¹³C-NMR spectra of known internal photodimers and newly obtained in 6-azauracils series showed their close similarity. Obtained results simply suggested that these compounds should have the cis-syn geometry. The NMR data are collected in Table 2 and Table 3.


The chemical shifts of H^{6'} and H⁵ in all reported compounds are almost identical. Also geometry of seven-member diazacycloheptane ring on the basis of NMR data seems to be identical. The very complex pattern of splittings of CH₂^C signals and also suprisingle of H_B and H_B, need some additional comment. This complicated picture of splittings may be probably attributed to second order splitting pattern or internal isomerization. In case of cis-syn geometry of obtained photodimers the expected internal isomerization is due to the possibility of equilibrium between two relatively stable rotamers.





	R^5	$R^{5'}$	$H^{6'}$	H_A, H_A'	H_B, H_B'	H^C
Thy[1(CH ₂) ₃ 1]6-azaUra 4	4.97, s	1.92, s	5.13, s	4.66, dt, $^2J_{H_A H_B} = 14$ Hz $^3J_{H_A H^C} = 3.5$ Hz	3.0, /dt?/ complex mult.	1.6-1.9 complex mult.
<u>2a</u>	4.95, s	CH ₃ , 1.05, t, $^3J = 7.5$ Hz	5.22, s	4.64, dt, $^2J_{H_A H_B} = 14$ Hz $^3J_{H_A H^C} = 3.5$ Hz	3.0, /dt?/ complex mult.	1.9-2.8 complex mult.
<u>2b</u>	4.95, s	CH ₃ , 0.86, t, $^3J = 7.5$ Hz	5.23, s	4.68, dt, $^2J_{H_A H_B} = 14$ Hz $^3J_{H_A H^C} = 3.5$ Hz	3.0, /dt?/ complex mult.	1.6-2.7 complex mult.
<u>2c</u>	1.94, s	4.10, d, $^3J = 7.5$ Hz	5.25, d $^2J = 7.5$ Hz	4.56, dt, $^2J_{H_A H_B} = 14$ Hz $^3J_{H_A H^C} = 4.0$ Hz	3.0, /dt?/ complex mult.	1.6-2.7 complex mult.

Table 2. ¹H-NMR spectra of some dinucleotide analogues /all spectra in PyH-d₆, in ppm downfield from TMS/



	R ⁵	R ^{5'}	C ^{6'}	C ^{5'}	C ⁵	C ^{5'}	N-C	N-C'	C-C-C	C ² , C ^{2'}	C ⁴ , C ^{4'}
Ura[1(CH ₂) ₃]6-azaUra	-	-	77.0	47.5	55.3	47.5	45.3	43.3	20.9	151.6 151.3	168.7 167.6
Thy[1(CH ₂) ₃]6-azaUra	-	21.7	81.2	45.5	66.3	45.5	46.6	46.6	23.1	150.6 149.8	166.6 165.7
<u>2a</u>	-	8.1 27.7	78.3	48.3	64.5	48.3	46.7	46.7	23.1	150.3 149.5	165.7 165.5
<u>2b</u>	-	13.8 16.9 36.9	78.7	47.8	64.9	47.8	46.8	46.8	23.1	150.2 149.4	165.8 165.4
<u>2c</u>	reverses to <u>1c</u> within time of NMR spectrum performance										

 Table 3. ¹³C-NMR spectra of some dinucleotide analogues /all spectra in DMSO-d₆, in ppm downfield from TMS/

In both rotamers the pattern of N-CH₂ groups can be described as AA'BB', but only one pair of protons is effected by the stereoisomerization whereas another pair is not influenced by these changes. To prove such suggestion we run out the ¹H-NMR spectra in elevated temperature / up to 75°C/, where the complex pattern of splittings simplified. The hypothesis of existence of equilibrium between two rotamers is supported by ¹³C-NMR spectra. The additional set of signals, especially in carbonyl region, C⁶ and C^{6'}, and N-C-C-N appeared when spectra have been run at room temperature. The low solubility of 2a, 2b and 2c in solvents suitable for low temperature NMR study makes such experiments impossible to run. Such conclusions support strongly our predictions concerning cis-syn configuration of obtained photo-dimers. In case of trans-syn geometry this rotameric equilibrium does not occur. The anti configuration is not taken for the consideration because of length of trimethylene chain. This work was supported by the Polish Academy of Sciences.

REFERENCES

- 1.K.Golankiewicz,J.Jankowska and H.Koroniak, Heterocycles, in press.
- 2.D.T.Browne,J.Eisinger and N.J.Leonard, J.Amer.Chem.Soc.,1968, 90, 7302.
- 3.K.Golankiewicz and L.Strekowski, Mol.Photochem., 1972, 4, 189.
- 4.J.Jankowska and H.Koroniak, unpublished data.

Received, 12th December, 1983