


A CONVENIENT SYNTHESIS OF NOVEL 3-AMIDINO-2-OXO-1,2-DIHYDROQUINOXALINE


Yoshihisa Kurasawa*, Yoshihisa Okamoto, and Atsushi Takada
 School of Pharmaceutical Sciences, Kitasato University,
 Shirokane, Minato-ku, Tokyo 108, Japan

Abstract — The reactions of 3-(5-oxo-1,2,4-oxadiazolin-3-yl)-2-oxo-1,2-dihydroquinoxaline (2) and 2-oxo-1,2-dihydroquinoxaline-3-amidoxime (3), derived from 3-cyano-2-oxo-1,2-dihydroquinoxaline (1), with $\text{FeSO}_4\text{-HCl}$ and Fe-HCl afforded Fe(II) complexes of 3-amidino-2-oxo-1,2-dihydroquinoxaline (4), whose treatment with NaOH gave the free ligand 4.

In a previous paper,¹ we reported the conversion of 3-cyano-2-oxo-1,2-dihydroquinoxaline (1) into 3-(5-oxo-1,2,4-oxadiazolin-3-yl)-2-oxo-1,2-dihydroquinoxaline (2) via 2-oxo-1,2-dihydroquinoxaline-3-amidoxime (3). However, there have been no reports on the conversion of 1 into 3-amidino-2-oxo-1,2-dihydroquinoxaline (4) (Scheme 1). Since further reduction of N-O bond of 2 or 3 would result in the formation of 4, accomplishment of this process would provide an access to a route of nitrile to amidine, presumably bearing comparison with the Pinner's amidine synthesis.² In



Scheme 1



Scheme 2

Table I. IR and ¹H-NMR Spectral Data for 4, 5a, and 5b.

Compound	IR ν (KBr) (cm ⁻¹)	¹ H-NMR (solvent) δ (ppm)
3-AQ (4)	3420 1700	(CF ₃ COOH): 9.28(s, 1H, NH), ^a 8.40-7.33(m, 4H, aromatic)
Fe(3-AQ) ₂ Cl ₂ ·H ₂ O (5a)	3570 1700 3400 3300	(DMSO-d ₆): 9.64(br.s, 4H, NH), 8.00-7.17(m, 4H, aromatic), 3.50(br.) ^b
Fe(3-AQ) ₂ SO ₄ ·H ₂ O (5b)	3450 1705 3320 1685 3170	(DMSO-d ₆): 9.57(br.s, 4H, NH), 8.00-7.33(m, 4H, aromatic), 4.23(br.) ^b

a: Three other NH protons disappeared, presumably due to CF₃COOH. b: A signal due to H₂O.

Table II. Mass Spectral Data for 4, 5a, and 5b.

Compound	m/z	Ion species	Formula	Calcd.	Found	Relat. abundance
<u>4</u>	188	[M] ⁺	C ₉ H ₈ N ₄ O	188.070	188.068	100
	171	[M-NH ₃] ⁺	C ₉ H ₅ N ₃ O	171.043	171.044	24.0
	160	[M-CO] ⁺	C ₈ H ₈ N ₄	160.075	160.073	46.8
<u>5a</u>	same as <u>4</u> *				188.071	100
					171.042	15.7
					160.073	32.0
<u>5b</u>	same as <u>4</u> *				188.068	100
					171.043	37.9
					160.073	32.1

* 5a and 5b exhibited M⁺ of the free ligand 4 due to thermal dissociation in the inlet system of the mass spectrometer.

this paper, we describe a facile synthesis of 4 via 2 or 3 from 1. Although the methods have already been reported for the conversion of 1,2,4-oxadiazoline ring and amidoxime into amidine using reducing agents such as P-HI³ and Fe(CO)₅,⁴ respectively, we have found that the reduction of N-O bond of 2 and 3 is conveniently achieved with an inexpensive and easily available agent FeSO₄-HCl⁵ and Fe-HCl to give 4 (3-AQ).

Refluxing of 3 (4 g) with Fe (2 g) in c.HCl (50 ml) and AcOH (150 ml) provided the Fe(3-AQ)₂Cl₂·H₂O complex (5a) (4 g, 88.3%) as yellow needles (from H₂O-EtOH) of mp 269-270 °C (dec.) [Anal. Calcd for C₁₈H₁₈Cl₂FeN₈O₃ (5a): C, 41.48; H, 3.48; Cl, 13.61; N, 21.50. Found: C, 41.62; H, 3.83; Cl, 13.65; N, 21.71.]. The reaction of 3 (1 g) with FeSO₄ (5 g) in c.HCl (10 ml) and AcOH (90 ml) produced the Fe(3-AQ)₂·SO₄·H₂O complex (5b) (600 mg, 50.4%) as yellow needles (from H₂O-EtOH) of mp 190-193 °C [Anal. Calcd for C₁₈H₁₈FeN₈O₇S (5b): C, 39.57; H, 3.32; N, 20.51; S, 5.87. Found: C, 39.83; H, 3.75; N, 20.44; S, 5.82.]. Moreover, similar reactions of 2-oxo-1,2-dihydroquinoxaline-3-amidoxime (6) (1 g) with Fe (1 g) and FeSO₄ (5 g) in c. HCl (10 ml) and AcOH (90 ml) gave 5a (720 mg, 56.3%) and 5b (850 mg, 63.4%), respectively. Treatment of 5a and 5b (1 g) with 10% NaOH (10 ml) in H₂O (100 ml)

furnished the free ligand 3-AQ (4) (640 mg, 83.3% from 5a; 480 mg, 69.7% from 5b) as yellow needles (trituration with EtOH-H₂O) of mp 310-312 °C [Anal. Calcd for C₉H₈N₄O (4): C, 57.44; H, 4.29; N, 29.77. Found: C, 57.21; H, 4.29; N, 29.47.].

The structural assignment of 4, 5a, and 5b (Scheme 2) was based on the microanalytical and spectral data (Tables I and II). Especially, the IR spectral data indicated that the NH absorption bands are quite different among 4, 5a, and 5b, presumably due to the presence or absence of a chelation, while the chelation hardly exerts an influence on the C=O absorption bands of 4, 5a, and 5b, suggesting no participation of the C=O group in the chelation.⁶ Mass spectra of 4, 5a, and 5b showed similar fragmentations.

REFERENCES AND FOOTNOTES

1. Y. Kurasawa, M. Ichikawa, A. Sakakura, and A. Takada, Chem. Pharm. Bull., submitted.
2. A. W. Dox, "Organic Syntheses," Coll. Vol. I, ed. by A. H. Blatt, John Wiley and Sons, Inc., New York, 1932, p. 5.
3. C. Behr, "The Chemistry of Heterocyclic Compounds," 1962, 17, 256, and references cited therein.
4. A. Dondoni, J. C. S. Chem. Comm., 1975, 761.
5. L. I. Smith and J. W. Opie, "Organic Syntheses," Coll. Vol. III, 1955, p. 56.
6. G. S. Sanyal, A. B. Modak, and A. K. Mudi, Ind. J. Chem., 1981, 20A, 510; M. Noji, Y. Kidani, and H. Koike, Bull. Chem. Soc. Japan, 1975, 48, 2274.

Received, 13th February, 1984