

THE REACTION OF 1-METHYL-2-VINYLIINDOLES WITH 4-PHENYL-1,2,4-
TRIAZOLINE-3,5-DIONE

Rosa Teresa Sanchis-Llorca,^a José Sepúlveda-Arques,^{a*} Elena
Zaballos-García,^a and R. Alan Jones^{b*}

^a Facultad de Farmacia, Departamento de Química Orgánica,
Universidad de Valencia, 46010-Valencia, Spain

^b School of Chemical Sciences, University of East Anglia,
Norwich NR4 7TJ, UK

Abstract - 1-Methyl-2-vinyliindoles ($1 \rightleftharpoons 2$) react readily with
4-phenyl-1,2,4-triazoline-3,5-dione at 20°C to yield the
Michael-type 1:1 adducts (3). No evidence was found for the
formation of any cycloadducts.

2-Vinylpyrroles react cleanly and in high yield at the free 5-position of the pyrrole ring with diethyl azodicarboxylate (DEAD) to yield Michael-type adducts¹ in a manner analogous to that reported for the corresponding reaction with simple 1-alkylpyrroles² or, when the 5-position is blocked or sterically hindered, they produce pyrrolo[3,2-c]pyridazine derivatives or dihydropyrrolyl-1,3,4-oxadiazines.¹ In contrast, the reactions of simple alkylindoles³ and of 2- and 3-vinyliindoles⁴ with DEAD are extremely complex and produce many unstable products in low yield. We now report that 1-methyl-2-vinyliindoles ($1 \rightleftharpoons 2$) generally react readily at 20°C with 4-phenyl-1,2,4-triazoline-3,5-dione (PTAD) to produce stable crystalline 1-(1-methyl-2-vinylindol-3-yl)-4-phenyl-1,2,4-triazolidine-3,5-diones (3) in high yield. However, the instability of 1-methyl-2-vinyliindole⁵ is such that only polymeric material was obtained from the reaction with PTAD, and the relatively low yield of (3a) from 2-(1-methylindol-2-yl)propene can be rationalised in terms of its susceptibility to dimerise readily at relatively low temperatures.⁵ No evidence was found for the formation of $\pi 2 + \pi 4$ cycloadducts, which is in keeping with the reported Michael-type addition of 1-methyl-2-vinyliindoles with dimethyl acetylene-dicarboxylate and their failure to yield dihydrocarbazoles.⁶ These observations are

2-(1-Methylindol-2-yl)propene⁶ gave 1-(1-methyl-2-(propen-2-yl)indol-3-yl)-4-phenyl-1,2,4-triazolidine-3,5-dione (0.70 g, 50%), m p 125 - 128°C, Rf 0.12 (Found: C, 69.1; H, 5.2, N, 16.2 C₂₀H₁₈N₄O₂ requires C, 69.4, H, 5.2, N, 16.2%). δ_{H} 2.05 (3H, s), 3.60 (3H, s), 5.20 (1H, brs), 5.40 (1H, brs), and 7.10 - 7.40 (10H, m); δ_{C} 22.8 (q), 30.6 (q), 107.1 (s), 109.8 (d), 117.6 (d), 120.8 (d), 121.6 (t), 122.7 (d), 123.9 (s), 125.6 (d), 127.9 (d), 128.9 (d), 131.7 (s), 133.8 (s), 135.6 (s), 141.4 (s), 151.6 (s), and 153.1 (s).

3,3-Dimethyl-2-(1-methylindol-2-yl)but-1-ene⁶ gave 1-(1-methyl-2-(3,3-dimethylbut-1-en-2-yl)indol-3-yl)-4-phenyl-1,2,4-triazolidine-3,5-dione (0.207 g, 82%), m p 104 - 107°C, Rf 0.12 (Found: C, 70.4; H, 6.2; N, 14.0, M⁺ 388.1869 C₂₃H₂₄N₄O₂ requires C, 71.1; H, 6.2; N, 14.4%, M⁺ 388.1898). δ_{H} 1.10 (9H, s), 3.50 (3H, s), 5.18 (1H, d, J = 1Hz), 5.54 (1H, d, J = 1Hz), and 7.00 - 7.55 (9H, m); δ_{C} 31.3 (q), 37.3 (s), 108.4 (s), 109.9 (d), 117.8 (s), 120.6 (t), 120.9 (d), 122.6 (d), 123.5 (s), 125.6 (d), 128.0 (d), 129.0 (d), 131.8 (s), 135.6 (s), 140.5 (s), 148.0 (s), 151.2 (s), and 153.2 (s).

1-(1-Methylindol-2-yl)-1-phenylethene⁶ gave 1-(1-methyl-2-(1-phenylethen-1-yl)indol-3-yl)-4-phenyl-1,2,4-triazolidine-3,5-dione (0.236 g, 90%), m p 169 - 171°C, Rf 0.16 (Found: C, 73.1; H, 4.9; N, 13.5 C₂₅H₂₀N₄O₂ requires C, 73.5; H, 4.9; N, 13.7%). δ_{H} 3.23 (3H, s), 5.55 (1H, d, J = 1Hz), 5.83 (1H, d, J = 1Hz) and 7.00 - 7.50 (14H, m); δ_{C} 30.9 (q), 109.4 (s), 109.9 (d), 117.9 (d), 121.0 (d), 121.1 (t), 123.9 (s), 125.6 (d), 126.6 (d), 127.9 (d), 128.4 (d), 128.6 (d), 128.9 (d), 131.6 (s), 135.8 (s), 137.6 (s), 138.7 (s), 139.4 (s), 151.5 (s), and 153.1 (s).

ACKNOWLEDGEMENTS

We are grateful for financial support from C.A.I.C.Y.T. (project 1499/82) and the British Council. We also thank Mr John Egles of the Institute for Food Research Norwich for the measurement of the high resolution mass spectra.

REFERENCES

1. T. Aznar-Saliente, R.A. Jones, R.T. Sanchis-Llorca, and J. Sepúlveda Arques, J. Chem. Research (S), 1985, 12; (M) 0232 - 0240.
2. W. Flitsch and J. Heinrich, Tetrahedron Lett., 1980, 21, 3673; C.K. Lee, S.J. Lim, and C.S. Hahan, J. Org. Chem., 1980, 45, 1692.

3. M. Collona and A. Monte, Gazz. Chim. Ital., 1962, **92**, 1401; H. Pleininger and D. Wild, Chem. Ber., 1966, **99**, 3063.
4. (a) E. Zaballos-Garcia, Ph.D. Thesis, University of Valencia, 1986;
(b) J.M. Weinmann, Ph.D. Thesis, University of Minnesota, 1964.
5. F.E. Ziegler, E.B. Spitzner, and C.K. Wilkins, J. Org. Chem., 1971, **36**, 1759.
6. R.A. Jones, P. Martinez Fresneda, T. Aznar-Saliente, and J. Sepúlveda Arques, Tetrahedron, 1984, **40**, 4837.
7. J.H. Hall, L. Kaler, and R. Herring, J. Org. Chem., 1984, **49**, 2579.

Received, 11th August, 1986