

A CONVENIENT SYNTHETIC APPROACH TO 4-SUBSTITUTED INDOLES¹

Fumio Yamada and Masanori Somei*


Faculty of Pharmaceutical Sciences, Kanazawa University

13-1 Takara-machi, Kanazawa 920, Japan

Abstract ——— A simple and practical synthetic method for 4-halogenoindoles, 4-indolecarbaldehyde, 4-cyanoindole, and 4-methoxycarbonylindole is elaborated.

4-Halogenoindoles,² 4-indolecarbaldehyde,³ 4-cyanoindole,^{2b,4} and 4-alkoxycarbonylindoles^{3c,5} are useful building blocks for the construction of natural alkaloids having 4-substituted indole nucleus. However, they are still not readily available, though much efforts have been paid on their syntheses. In this communication, we wish to describe a simple and practical synthetic method for these building blocks. We have already established an experimental procedure for the synthesis of (3-formylindol-4-yl)thallium bis-trifluoroacetate (2) starting from 3-indolecarbaldehyde (1) in 100-200g scale.⁶ Now, we have found that the thallium compound (2) can be used as a common intermediate for the syntheses of above mentioned building blocks. Thus, carbonylation of 2 in methanol⁷ in the presence of a catalytic amount of palladium acetate afforded methyl 3-formyl-4-indolecarboxylate (3) in 54% yield. Cyanation of 2 with cuprous cyanide⁸ in refluxing N,N-dimethylformamide produced 4-cyano-3-indolecarbaldehyde (4) in 53% yield. Preparation of 4-halogeno-3-indolecarbaldehydes (5a-c) was carried out by the reaction of 2 with appropriate halogenating reagents according to our reported procedure.^{2a}

Since 4-substituted indoles having no substituent at the 3-position are sometimes needed, we next tried to get them utilizing above mentioned readily accessible compounds, 3, 4, and 5a-c. Direct decarbonylation of 5a by treatment with either acids or bases were unsuccessful. Therefore, various oxidative reagents were examined on 5a as a representative substrate with an aim to obtain the corresponding 4-iodo-3-indolecarboxylic acid (8a) but all attempts⁹ ended in failure. Finally, oxidation with sodium chlorite in the presence of 2-methyl-2-butene¹⁰ was found to work well



and 5a was converted to 8a in 73% yield. Under similar reaction conditions, methyl 3-formyl-4-indolecarboxylate (3), 4-cyano- (4), 4-bromo- (5b), and 4-chloro-3-indolecarbaldehyde (5c) were successfully transformed to the corresponding carboxylic acids, 6, 7, 8b, and 8c, in 77%, 90%, 86%, and 93% yields, respectively. Treatment of 8a with ethereal diazomethane afforded methyl 4-iodo-3-indolecarboxylate (14a) in quantitative yield, although direct conversion of 5a to 14a using manganese dioxide and sodium cyanide in methanol¹¹ proceeded in only 24% yield. Based upon these results, when the sodium chlorite oxidation reaction was carried out and followed by treatment with ethereal diazomethane, one pot syntheses of methyl 4-halogeno-3-indolecarboxylates (14a, b, c¹²) was realized in 78%, 81%, and 93% yields, respectively. These compounds were identical with the authentic samples prepared from methyl 3-indolecarboxylate (16) via (3-methoxycarbonylindol-4-yl)-thallium bis-trifluoroacetate (15) and were easily led to the corresponding 4-halogenoindoles (11a-c) in high yields according to the reported alkaline hydrolysis reaction.^{2a}

Decarboxylation of 4-substituted-3-indolecarboxylic acids were readily attained simply by heating them in pyridine. Under the reaction condition, 6, 7, and 8a afforded the expected 4-substituted indoles, 9, 10, and 11a, in 99%, 90%, and 89% yields, respectively. Subsequent reduction¹³ of 4-cyanoindole (10) with diisobutylaluminum hydride produced 4-indolecarbaldehyde (13) in 62% yield. Since interconversion^{3,4,5} among 9, 4-hydroxymethylindole¹⁴ (12), and 13 are well established, various simple 4-substituted indoles (3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, and 14) became now readily available from 3-formylindole (1) in hundreds gram scale.

REFERENCES AND NOTES

1. This report is part XXXVIII of a series entitled "The Chemistry of Indoles". Part XXXVII: M. Somei, F. Yamada, and Y. Makita, *Heterocycles*, in press, 26.
2. a) M. Somei, K. Kizu, M. Kunitomo, and F. Yamada, *Chem. Pharm. Bull.*, **33**, 3696 (1985); b) M. Somei and M. Tsuchiya, *Chem. Pharm. Bull.*, **29**, 3145 (1981); c) M. Matsumoto, Y. Ishida, and N. Hatanaka, *Heterocycles*, **24**, 1667 (1986); d) P.J. Harrington and L.S. Hegedus, *J. Org. Chem.*, **49**, 2657 (1984).
3. a) N. Hatanaka, N. Watanabe, and M. Matsumoto, *Heterocycles*, **24**, 1987 (1986); b) M. Somei, F. Yamada, Y. Karasawa, and C. Kaneko, *Chemistry Lett.*, 1981, 615; M. Somei, Y. Karasawa, and C. Kaneko, *ibid.*, 1980, 813; c) A.P. Kozikowski, H. Ishida, and Y.-Y. Chen, *J. Org. Chem.*, **45**, 3350 (1980).
4. a) N. Hatanaka and M. Matsumoto, *Heterocycles*, **24**, 1963 (1986); b) R.D. Clark and C.B. Repke, *Heterocycles*, **22**, 195 (1984); c) R.D. Clark, *J. Heterocyclic*

- Chem., 20, 1393 (1983); d) G.S. Ponticello and J.J. Baldwin, J. Org. Chem., 44, 4003 (1979); e) F. Troxler, A. Harnisch, G. Bormann, F. Seeman, and L. Szabo, Helv. Chim. Acta., 51, 1616 (1968); f) F.C. Uhre, J. Am. Chem. Soc., 71, 761 (1949)
5. a) T. Sakamoto, Y. Kondo, and H. Yamanaka, Heterocycles, 22, 1347 (1984);
b) M. Somei, Y. Karasawa, T. Shoda, and C. Kaneko, Chem. Pharm. Bull., 29, 249 (1981); c) L.I. Kruse, Heterocycles, 16, 1119 (1981); d) M. Somei and M. Natsume, Tetrahedron Lett., 1973, 2451; e) T. Watanabe, F. Hamaguchi, and S. Ohki, Chem. Pharm. Bull., 20, 2123 (1972).
6. M. Somei and F. Yamada, Chem. Pharm. Bull., 32, 5064 (1984); M. Somei, Y. Makita, and F. Yamada, ibid., 34, 948 (1986).
7. R.C. Rarock and C.A. Fellows, J. Am. Chem. Soc., 104, 1900 (1982).
8. E.C. Taylor and A.H. Katz, Tetrahedron Lett., 25, 5473 (1984).
9. Various oxidation conditions were examined using CrO₃, PCC, and KMnO₄, but all attempts to obtain 8a were unsuccessful.
10. B.S. Bal, W.E. Childers, Jr., and H.W. Pinnick, Tetrahedron, 37, 2091 (1981).
11. E.J. Corey, N.N. Gilman, and B.E. Ganem, J. Am. Chem. Soc., 90, 5616 (1968).
12. Melting points of 14b and 14c, reported in our previous paper,^{2a} should be revised to 113.0-114.0°C (recrystallized from MeOH-H₂O) and 153.0-155.0°C (recrystallized from CH₂Cl₂-n-hexane), respectively.
13. H. Plieninger, M. Höbel, and V. Liede, Chem. Ber., 96, 1618 (1963); H. Plieninger, ibid., 88, 370 (1955).
14. M. Somei and T. Shoda, Heterocycles, 17, 417 (1982) and references cited therein.

Received, 24th December, 1986