

HETEROCYCLES, Vol. 89, No. 5, 2014, pp. 1125 - 1157. © 2014 The Japan Institute of Heterocyclic Chemistry
Received, 20th December, 2013, Accepted, 25th December, 2013, Published online, 10th January, 2014
DOI: 10.3987/REV-13-790

HETEROCYCLIZATION WITH SOME HETEROCYCLIC DIAMINES: SYNTHETIC APPROACHES FOR NITROGEN BRIDGEHEAD HETEROCYCLIC SYSTEMS

Magdy A. Ibrahim* and Nasser M. El-Gohary

Department of Chemistry, Faculty of Education, Ain Shams University, Roxy,
11711, Cairo-Egypt

*E-mail: magdy_ahmed1977@yahoo.com

Abstract – This review represents the methods developed for the synthesis of a variety of nitrogen bridgehead heterocyclic systems such as [1,2,4]triazolo[1,5-*a*]pyridines, pyrido[1,2-*b*][1,2,4]triazines, pyrido[1,2-*b*][1,2,4]-triazepines, pyrazolo [1,5-*b*][1,2,4]triazole, pyrazolo[1,5-*b*][1,2,4]triazine, pyrazolo[1,5-*b*][1,2,4]triazepine, [1,2,4]triazolo[3,4-*b*][1,2,4]triazine, [1,2,4]triazolo[4,3-*b*][1,2,4]triazepines, pyrimido[3,4-*b*][1,2,4]triazine from the heterocyclization of some heterocyclic diamines with a variety of electrophilic reagents.

CONTENTS

1. Introduction
2. Heterocyclization with diaminopyridones
 - 2.1. Synthesis of *o*-diaminopyridones
 - 2.2. Synthetic approaches for [1,2,4]triazolo[1,5-*a*]pyridines
 - 2.3. Synthetic approaches for pyrido[1,2-*b*][1,2,4]triazines
 - 2.4. Synthetic approaches for pyrido[1,2-*b*][1,2,4]triazepines
3. Heterocyclization with diaminoimidazoles
4. Heterocyclization with diaminotriazoles
5. Heterocyclization with diaminothiazoles
6. Heterocyclization with diaminopyrimidines
7. Heterocyclization with diaminotriazine

1. INTRODUCTION

o-Diamines are very active substrates for building of various heterocyclic systems,¹⁻⁵ and are largely used in formation of complexes.⁶⁻¹⁰ In symmetrical diamines, the product will be the same irrespective of which amine participates first in the reaction. In the case of unsymmetrical diamines, the substituents influence the initial participation of a particular amino group in the reaction, resulting in chemoselective products. The electron withdrawing/donating nature of substituents in diamine influences the nucleophilicity of the amino groups. The present review article concise on the utilities of heterocyclic diamines in heterocyclic synthesis *via* reactions with a large numbers of electrophilic reagents.


2. Heterocyclization with diaminopyridones

Polyfunctional pyridines are highly reactive reagents that have been used extensively in heterocyclic synthesis,¹¹⁻¹³ and possess biological as well as pharmacological activities.¹⁴⁻¹⁶ [1,2,4]Triazolo[1,5-*a*]-pyridines, pyrido[1,2-*b*][1,2,4]triazines and pyrido[1,2-*b*][1,2,4]triazepines are also interesting compounds due to their pronounced biological importance. *o*-Diaminopyridone derivatives are widely used in the synthesis of a variety of nitrogen bridgehead triazolo[1,5-*a*]pyridines, pyrido[1,2-*b*][1,2,4]-triazines and pyrido[1,2-*b*][1,2,4]triazepines *via* heterocyclization of *o*-diaminopyridones with some mono electrophilic reagents, α,β -bifunctional and α,γ -bifunctional electrophiles, respectively.


2.1. Synthesis of *o*-diaminopyridones

Refluxing alcoholic solution of 2-cyanoacetohydrazide (**1**) with arylmethylidinemalononitriles **2** (X=CN) and ethyl 3-cyano-2-(aryl)-prop-2-enoates **2** (X=CO₂Et), in the presence of a few drops of piperidine as a catalyst, produced 4-(aryl)-1,6-diamino-2-oxo-1,2-dihydropyridine-3,5-dicarbonitriles **3** (X=CN) and ethyl 1,2-diamino-5-cyano-4-(aryl)-6-oxo-1,6-dihydropyridine-3-carboxylates **3** (X=CO₂Et), respectively (Scheme 1).¹⁷ This method is a rapid and common for the synthesis of *o*-diaminopyridone derivatives. The proposed mechanism for formation of compounds **3** is depicted in Scheme 2.¹⁷


Also, 3-cyano-1,6-diaminopyridone derivatives **5** possessing various alkoxy carbonyl groups were prepared directly from the reaction of 2-cyanoacetohydrazide (**1**) with dialkyl 2,3-dicyanobutenedionates **4** (Scheme 3).¹⁸ The ¹H NMR spectrum is a good tool to differentiate between the nucleophilicity of the two amino groups in the diaminopyridone derivatives **3** and **5**. The ¹H NMR spectra usually showed two exchangeable signals in the range δ 4.61-5.70 ppm and 8.30-10.78 ppm characteristic for the (*N*-NH₂) and (*C*-NH₂) protons, respectively. These results indicate the difference in nucleophilicity between the two amino groups. Thus, the hydrazide β -nitrogen (*N*-NH₂) is more nucleophilic and reacted more rapidly with the electron deficient carbon than the amino group at carbon atom (*C*-NH₂).



Scheme 1



Scheme 2




Scheme 3

2.2. Synthetic approaches for [1,2,4]triazolo[1,5-*a*]pyridines


[1,2,4]Triazolo[1,5-*a*]pyridines constitute an important class of heterocyclic systems due to their variable biological activities including antifungal,¹⁹ antimicrobial,²⁰ antitumor,^{21,22} analgesic, anti-inflammatory,²³ and antiviral activity.²⁴ Different methods are reported for the synthesis of 1,2,4-triazolo[1,5-*a*]-

pyridines,²⁵⁻³⁵ the most common one is the condensation of *o*-diaminopyridones **3** or **5** with mono electrophilic reagents. Thus, 7-(aryl)-5-oxo-1*H*-4,5-dihydro[1,2,4]triazolo[1,5-*a*]pyridine derivatives **6** were prepared by heterocyclization of 4-aryl-1,6-diaminopyridones **3** with formic acid,³⁵ ethyl formate,³⁶ or triethyl orthoformate (Scheme 4).³⁷



Scheme 4


4-Aryl-1,6-diamino-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile **3** reacted with *N*-ethoxymethylenebenzohydrazide in isopropanol to give [1,2,4]triazolo[1,5-*a*]pyridine-6,8-dicarbonitrile **6** which reacted with benzohydrazide to give triazolopyridones **7** and 7-aryl-6,8-bis(5-phenyl-4*H*-1,2,4-triazol-3-yl)-4,5-dihydro-1*H*-[1,2,4]triazolo[1,5-*a*]pyridine-5-ones **8** (Scheme 5).³⁸



Scheme 5


Refluxing diaminopyridone **3** with acetyl chloride and benzoyl chloride in boiling DMF produced

7-(4-chlorophenyl)-5-oxo-1*H*-4,5-dihydro-2-(methyl/phenyl)[1,2,4]triazolo[1,5-*a*]pyridine-6,8-dicarbonitriles **9** (Scheme 6).³⁵ Also, compounds **9** (R=Me) were prepared by the reaction of diaminopyridones **3** with acetic anhydride (Scheme 6).³⁹




Scheme 6

On the other hand, 1-acetyl-7-(6-chloro/methyl-4-oxo-4*H*-chromon-3-yl)-2-methyl-5-oxo-1,5-dihydro[1,2,4]triazolo[1,5-*a*]pyridine-6,8-dicarbonitriles **10** were prepared by heterocyclization of diaminopyridone derivatives **3** with acetic anhydride (Scheme 7).^{36, 40}




Scheme 7

2-Substituted[1,2,4]triazolo[1,5-*a*]pyridine derivatives **11** was prepared in high yield by cyclocondensation of 3-cyano-1,6-diaminopyridone derivatives **5** with carboxylic acid orthoesters such as triethyl orthoformate or triethyl orthoacetate (Scheme 8).¹⁸




Scheme 8

Diaminopyridones **3** reacted with triethyl orthoformate and triethyl orthoacetate as cyclizing and alkylating agent in acetonitrile to give the unexpected 3-ethyl[1,2,4]triazolo[1,5-*a*]pyridine derivatives **12** (Scheme 9).³⁹



Scheme 9

1,6-diaminopyridones **3** reacted with aromatic aldehydes in refluxing methanol in the presence of catalytic amount of sulfuric acid to give 7-aryl-2-(4-fluorophenyl)-5-oxo-1,5-dihydro-1*H*-[1,2,4]-triazolo[1,5-*a*]pyridine-6,8-dicarbonitrile **13** (Scheme 10).^{38,39}




Scheme 10


Cyclocondensation of 1,6-diaminopyridone **3** with 6-chloro-3-formylchromone **14** in DMF under reflux containing few drops of piperidine afforded the [1,2,4]triazolo[1,5-*a*]pyridine derivative **15**. Oxidation of the latter compound by ferric chloride in boiling dimethyl sulfoxide (DMSO) yielded 2,7-bis(6-chloro-4-oxo-4*H*-chromen-3-yl)-5-oxo-1,5-dihydro-1,2,4-triazolo[1,5-*a*]pyridine-6,8-dicarbonitrile **16**.³⁶ Compound **16** was also obtained by refluxing compound **3** with 6-chlorochromone-3-carboxylic acid **17** in phosphoryl chloride (Scheme 11).⁴⁰

7-(2-Furyl)-5-oxo-3,5-dihydro[1,2,4]triazolo[1,5-*a*]pyridine derivatives **18** were prepared by the reaction of diaminopyridones **3** with ethyl chloroformate in boiling DMF (Scheme 12).³⁹


Refluxing 1,6-diaminopyridone **3** with carbon disulfide in ethanolic KOH yielded 2-thioxo[1,2,4]-triazolo[1,5-*a*]pyridine-6,8-dicarbonitrile **19**,⁴¹ which oxidized by FeCl₃/EtOH to give 7,7-di(4-chlorophenyl)-5,5-dioxo-1,1,5,5-tetrahydro-2,2-dithio-di[1,2,4]triazolo[1,5-*a*]pyridine-6,6,8,8-tetracarbonitrile **20** (Scheme 13).³⁵



Scheme 11



Scheme 12




Scheme 13


The diaminopyridone derivative **3** reacted with carbon disulfide in ethanolic KOH solution under reflux to yield [1,2,4]triazolo[1,5-*a*]pyridine derivative **19** which reacted with chloroacetic acid, oxalyl chloride, chloroacetonitrile and bromomalonitrile to produce the corresponding thiazolo[3,2:2,3][1,2,4]triazolo[1,5-*a*]pyridines **31-34**, respectively (Scheme 14).⁴²

Also, reaction of 1,6-diaminopyridone **3** with triphenylphosphine gave 1-amino-6-(triphenylphosphoranylideneamino)-2-oxo-4-phenyl-1,2-dihydro-pyridine-3,5-dicarbonitrile-iminophosphorane **25** which reacted with carbon disulfide in dry toluene to give compound **19**. The latter compound

reacted with aromatic aldehydes, chloroacetic acid and fused sodium acetate in acetic acid/acetic anhydride to give 2-(4-substitutedbenzylidene)-3,6-dioxo-8-phenyl-3,6-dihydro-2H-thiazolo[3,2:2,3]-[1,2,4]triazolo[1,5-*a*]pyridine-7,9-dicarbonitrile **26** (Scheme 15).⁴³




Scheme 14



Scheme 15


Reaction of diaminopyridone **3** with anthranilic acid in phosphoryl chloride gave ethyl 2-(2-amino-phenyl)-6-cyano-7-(4-methoxyphenyl)-5-oxo-3,5-dihydro[1,2,4]triazolo[1,5-*a*]pyridine-8-carboxylate **27** which reacted with formic acid, acetyl chloride, chloroacetyl chloride, ethyl chloroformate and

carbonyl sulfide to give the corresponding pyrido[1,2:2,3][1,2,4]triazolo[1,5-*c*]quinazoline derivatives **28-32**, respectively (Scheme 16).⁴²



Scheme 16


2-(2-Benzoylamino-phenyl)-7-(4-chlorophenyl)-5-oxo-1*H*-[1,2,4]triazolo[1,5-*a*]pyridine-6,8-dicarbonitrile **35** was obtained from the reaction of diaminopyridone **3** and 2-phenyl-4*H*-3,1-benzoxazin-4-one **33**, via the non isolable intermediate **34** (Scheme 17).⁴⁴



Scheme 17

2.3. Synthetic approaches for pyrido[1,2-*b*][1,2,4]triazines


4-Aryl-1,6-diamino-2-oxo-1,2-dihydropyridine-3,5-dicarbonitriles **3** were efficiently used for preparation of nitrogen bridgehead pyrido[1,2-*b*][1,2,4]triazines. Thus, the isomeric structures of pyrido[1,2-*b*][1,2,4]triazine-7,9-dicarbonitrile derivatives **36** and **37** have been obtained from condensation of diaminopyridones **3** with chloroacetic acid and chloroacetyl chloride in refluxing DMF. Also, heterocyclization of diaminopyridones **3** with dibromoethane and phenacyl bromide in basic media gave pyrido[1,2-*b*][1,2,4]triazine-7,9-dicarbonitriles **38** and **39**, respectively (Scheme 18).⁴⁵




Scheme 18

Condensation of diaminopyridone **3** with 5,6-diphenyl-3-mercapto-1,2,4-triazine **40** gave *N*-(triazinyl-amino)pyridine derivative **41**. Heterocyclization of compound **41** with chloroacetic acid, chloroacetyl chloride, 1,2-dibromoethane, phenacyl bromide, diethyl oxalate, and α,β -unsaturated oxoacid **42** led to the direct formation of the pyrido[1,2-*b*][1,2,4]triazine derivatives **43-48**, respectively (Scheme 19).³⁵

On the other hand, cyclocondensation of compound **3** with α,β -unsaturated keto-acid **42** in refluxing glacial acetic acid yielded 8-aryl-2,6-dioxo-3-substituted-1,2,5,6-tetrahydropyrido[1,2-*b*][1,2,4]triazine-7,9-dicarbonitriles **49**, while its treatment with diethyl oxalate in dry dioxane and/or with oxalyl chloride in warming DMF afforded the 2,3,6-trioxo analogue **50** (Scheme 20). Hydroxymethylation of compound **50** using methanol-formaldehyde produced the 1,4-dihydroxymethylpyridotriazine derivative **51** which upon full heterocyclization by refluxing with thiosemicarbazide in glacial acetic acid led to the direct formation of 7-oxo-2-thioxo-2,3,5,6,7,11-hexahydropyrido[1',2':2,3][1,2,4]triazino[5,6-*e*]triazine-8,10-dicarbonitrile **52**. Careful hydrazinolysis of compound **52** afforded the corresponding 3-hydrazino-1,2,4-triazine derivative **53** (Scheme 20).⁴⁵




Scheme 19



Scheme 20


1,2-Dioxygen compounds also used for building of various fused heterocyclic systems. Thus, treatment compound **3** with butane-2,3-dione in glacial acetic acid afforded 2,3-dimethylpyrido[1,2-*b*][1,2,4]-triazine **54**, while the corresponding 2,3-diphenylpyridotriazine derivatives **55** were obtained from refluxing **3** with benzil in glacial acetic acid. The dihydro analogous **56** were obtained from refluxing compounds **3** with benzoin under the same reaction conditions. Oxidation of compounds **56** in methanolic ferric chloride produced compounds **55** (Scheme 21).⁴⁵ Also, some new 8-aryl-2,6-dioxo-1,2-dihydropyrido[1,2-*b*][1,2,4]triazine-7,9-dicarbonitriles **57** have been synthesized from cyclocondensation of compounds **3** with α -oxocarboxylic acids namely; glyoxalic, pyruvic, α -oxobutyric and 4-chlorostyrylglyoxalic in refluxing glacial acetic acid (Scheme 21).⁴⁵ It must be noted that this reaction occurred preferentially firstly between the N^1 -amino group (N -NH₂) and α -keto functions to form a hydrazone intermediate, which underwent cyclocondensation reaction between the other amino group at C⁶ (C -NH₂) and the hydroxyl group of the acid functions affording the target pyridotriazine derivatives **57**.



Scheme 21


The course of reactions of 1,2,-dioxygen heterocyclic compounds with aromatic heterocyclic *o*-diamines was found to depend on the reaction conditions, type of solvent (dielectric constant), and also the types of substituents (e-donor or e-acceptor) in the diamino compounds. Treatment of compound **3** with indole-2,3-dione in absolute ethanol containing a few drops of piperidine yielded the Schiff base condensate, 6-amino-4-aryl-2-oxo-1-[2-oxo-1,2-dihydro-3-indolo-3-ylidene)amino]-1,2-dihydropyridine-3,5-dicarbonitrile **58**. When that reaction was carried out in boiling alcoholic sodium hydroxide solution, it gave 3-(2-aminophenyl)pyridotriazine derivative **59**. Refluxing both **58** and **59** in glacial

acetic acid/fused sodium acetate gave the full condensation product, indolotriazinopyridine **60**, which on further acetylation by heating in acetic anhydride yielded the *N*-acetyl derivative **61** (Scheme 22).⁴⁵



Scheme 22

Reaction of diaminopyridone **3** with *N*-acetylisatine **62** in absolute ethanol containing a few drops of piperidine produced 8-aryl-2-(2-acetanilido)-3,6-dioxo-3,6-dihydro-4*H*-pyrido[1,2-*b*][1,2-4]triazine-7,9-dicarbonitrile **64**, via the non isolable intermediate **63** and not to the isomeric product **65** as illustrated in Scheme 23.⁴⁵




Scheme 23


Cyclocondensation of diaminopyridone **3** with ethyl 2-chloro-3-oxobutanoate in DMF under reflux containing a catalytic amount of piperidine produced the pyrido[1,2-*b*][1,2,4]triazine-2-carboxylate derivative **66**, which was transformed to the pyrido[1,2-*b*][1,2,4]triazino[4,5-*d*][1,2,4]triazine derivative **67** upon fusion with benzoic acid hydrazide (Scheme 25).³⁶ On the other hand, the interaction of **3** with 2,3-dichloroquinoxaline **68** under reflux led to the formation of the corresponding quinoxalino[2,3-*e*]-pyrido[1,2-*b*][1,2,4]triazine **69**. Compound **69** exists in two tautomeric forms **69A** and **69B** due to amino-imino tautomerism (Scheme 24).³⁶

Also, the reaction of compound **3** with 3-chloro-7,8-diphenyl-4*H*-1,2,4-triazino[4,3-*b*][1,2,4]triazine **70** in pyridine under reflux led to the formation of the corresponding pyrido[1,2-*b*][1,2,4]triazino-[3',2':3,4]triazino[5,6-*e*][1,2,4]triazine **71**, respectively (Scheme 25).³⁶


8-(4-Chlorophenyl)-2,6-dioxo-1*H*-pyrido[1,2-*b*][1,2,4]triazine-3,7,9-tricarbonitrile **73** was obtained from the reaction of 1,6-diaminopyridone **3** with ethyl α -cyano- α -phenylazoacetate **72** in acetic acid as shown in Scheme 26.⁴⁴



Scheme 24




Scheme 25



Scheme 26


Pyrido[1,2-*b*][1,2,4]triazin-2(1*H*)-one **75** was obtained from ring closure of diaminopyridone **3** with 3-(2,2-dichloroacetyl)-4-hydroxy-1-methylquinolin-2(1*H*)-one **74** in boiling DMF (Scheme 27).⁴⁶



Scheme 27


2.4. Synthetic approaches for pyrido[1,2-*b*][1,2,4]triazepines

Reaction of diaminopyridone derivatives with some α,γ -bifunctional electrophiles afforded pyrido[1,2-*b*][1,2,4]triazepines. When 1,6-diamino-4-(4-chlorophenyl)-2-oxo-1,2-dihydropyridine-3,5-dicarbonitrile **3** was allowed to react with diethyl malonate and ethyl ethoxymethylenecyanoacetate in boiling DMF, 9-(4-chlorophenyl)-pyrido[1,2-*b*][1,2,4]triazepine derivatives **76** and **77** were obtained (Scheme 28).⁴⁴




Scheme 28

Treatment of 1,6-diaminopyridones **3** with pentane-2,4-dione afforded 9-aryl-2,4-dimethyl-7-oxo-6,7-dihydropyrido[1,2-*b*][1,2,4]triazepine-8,10-dicarbonitriles **78**,^{38,44} which reacted with benzene diazonium chloride to give 3-phenylhydrazono-1*H*-pyrido[1,2-*b*][1,2,4]triazepine-8,10-dicarbonitrile **78**. Compound **79** was also prepared from treating diaminopyridone **3** with 3-phenylazo-2,4-pentanedione (Scheme 29).⁴⁴




Scheme 29

Pyrido[1,2-*b*][1,2,4]triazepines **80-82** were prepared by the reaction of 1,6-diaminopyridone **3** with 4-(dimethylamino)but-3-en-2-one, 2-cyano-3,3-bis-(methylthio)acrylonitrile and 2-cyano-3-(methylsulfanyl)-*N*-phenyl-3-(phenylamino)prop-2-enamide, respectively (Scheme 30).^{36,40}




Scheme 30

Condensation of diaminopyridone **3** with 5-chloro-3-methyl-1-phenylpyrazole-4-carboxaldehyde and 2-chloro-3-formylquinoline in DMF containing few drops in triethylamine afforded the heteroannulated pyrido[1,2-*b*][1,2,4]triazepines namely pyrazolo[3,4-*e*]pyrido[1,2-*b*][1,2,4]triazepine **83** and quinolino[2,3-*e*]pyrido[1,2-*b*][1,2,4]triazepine **84**, respectively (Scheme 31).⁴⁰



Scheme 31


2-Aminopyrido[1,2-*b*][1,2,4]triazepine-3,8,10-tricarbonitrile **85** was prepared by reaction of 1,6-diaminopyridone **3** and (4-chlorobenzylidene)malononitrile in boiling DMF and piperidine (Scheme 32).^{36,40}




Scheme 32

Condensation of diaminopyridone **3** with ethyl 2-cyano-3-(4-methoxyphenyl)prop-2-enoate under the same reaction conditions in boiling DMF containing two drops of triethylamine yielded ethyl 2-amino-8,10-dicyano-4-(4-methoxyphenyl)-9-(6-methyl-4-oxo-4*H*-chromen-3-yl)-7-oxo-5,7-dihydro-pyrido[1,2-*b*][1,2,4]triazepine-3-carboxylate **86** (Scheme 33).⁴⁰

Treatment of 1,6-diaminopyridone **3** with dehydroacetic acid and dimethyl acetylene-dicarboxylate afforded pyrano[2,3-*e*]pyrido[1,2-*b*][1,2,4]triazepine **87** and methyl pyrido[1,2-*b*][1,2,4]triazepine-4-carboxylate **88**, respectively (Scheme 34).⁴⁴




Scheme 33




Scheme 34

9-Aryl-4-methyl-2,4-diphenyl-7-oxo-3,4,6,7-tetrahydro-5*H*-pyrido[1,2-*b*][1,2,4]triazepine-8,10-dicarbonitrile **89** was prepared by the reaction of 1,6-diaminopyridone **3** with acetophenone in methanol in the presence of catalytic amount of sulfuric acid (Scheme 35).³⁸




Scheme 35

9-(Aryl)-3-(2-hydroxybenzoyl)-7-oxo-6*H*-pyrido[1,2-*b*][1,2,4]triazepine-8,10-dicarbonitrile **90** was prepared by the reaction of diaminopyridone **3** and chromone-3-carbaldehyde in dry pyridine (Scheme 36).⁴⁴




Scheme 36

Treatment of diaminopyridone **3** with chromone-3-carbonitrile in DMF under reflux gave 2-amino-3-(2-hydroxybenzoyl)-7-oxo-9-(aryl)-5*H*-pyrido[1,2-*b*][1,2,4]triazepine-8,10-dicarbonitrile **91** as described by Abdel-megid,^{40,44} while gave the Schiff base **92** as published by Sosnovskikh and Moshkin (Scheme 37).⁴⁷




Scheme 37

Reaction of 1,6-diaminopyridone **3** with 6-ethyl-3-[(4-oxo-4*H*-chromen-3-yl)methylidene]pyrano[3,2-*c*]-quinoline-2,4,5-(3*H*,6*H*)-trione **93** afforded the pyrido[1,2-*b*][1,2,4]triazepines **94**, bearing chromone and quinolinone nuclei (Scheme 38). The reactions proceeds initially *via* α -pyrone ring opening by the more nucleophilic amino group (*N*-NH₂) followed by ring closure to produce the desired products.⁴⁸



Scheme 38


The RORC reactions of chromone derivative **95** with 1,6-diaminopyridone **3** gave pyrido[1,2-*b*][1,2,4]-triazepine derivative **96** (Scheme 39).⁴⁹



Scheme 39


3. Heterocyclization with diaminoimidazoles

Condensation of 1,5-diamino-3-*tert*-butylpyrazole **97** with carbon disulfide, α , β -dicarbonyl compounds (diacetyl, benzil), ethyl pyruvate and acetylacetone gave the corresponding pyrazolo[1,5-*b*][1,2,4]triazole **98**, pyrazolo[1,5-*b*][1,2,4]triazine **99** and **110** and pyrazolo[1,5-*b*][1,2,4]triazepine **101** ring systems, respectively (Scheme 40).⁵⁰




Scheme 40

Cyclization of 1,2-diamino-4-arylimidazoles **102** with 1,3-disubstituted propenones produced imidazo[1,5-*b*]pyridazines **103**. While reaction of 1,2-diamino-4-arylimidazoles **102** with acetylacetone gave a mixture of 7-amino-5-(4-bromophenyl)-2,4-dimethylimidazo[1,5-*b*]pyridazine **104** and 5-(4-bromophenyl)-2,4-dimethyl-7-[(4-oxo-2-pent-2-en-yl)amino]imidazo[1,5-*b*]pyridazines **105** (Scheme 41), compound **104** and **105** were separated by crystallization process.⁵¹




Scheme 41

Ethyl 3-amino[1,2,4]triazino[2,3-*a*]benzimidazole-2-carboxylate **108** was obtained selectively by the reaction of 1,2-diaminobenzimidazole **106** with diethyl (*E*)-2,3-dicyanobutenedioate **107** in boiling acetonitrile (Scheme 42).⁵²



Scheme 42


On the other hand, 1,2,4-triazino[2,3-*f*]xanthines **110** and **111** were prepared from the reaction of 5,6-diaminotheophylline **109** with the acyclic and cyclic dicarbonyl compounds, respectively (Scheme 43).⁵³



Scheme 43


4. Heterocyclization with diaminotriazoles

Cyclocondensation of 3,4-diamino-5-methyl-1,2,4-triazole hydrochloride **112** with 1-methylisatine gave 3,10-dimethyl-1,2,4-triazolo[4',3':2,3][1,2,4]triazino[5,6-*b*]indole **113** (Scheme 44).⁵⁴




Scheme 44

3,4-Diamino-1,2,4-triazole hydrobromide **114** reacted with α -dicarbonyl compounds in acetic acid to afford the corresponding [1,2,4]triazolo[3,4-*b*][1,2,4]triazine derivative **115** (Scheme 45).⁵⁵




Scheme 45

4-(Phenyl/benzyl)-2-(2-chloroethyl)-1*H*-1,3,3a,5,6-pentaza-2-phosphapentalen-2-oxide **117** was prepared by the reaction of 5-phenyl/benzyl-1,2-diamino-1,3,4-triazole **116** with 2-chloroethylphosphonyl dichloride in pyridine containing triethylamine (Scheme 46).⁵⁶




Scheme 46

Reaction of 3,4,5-triamino-4*H*-1,2,4-triazole (guanazine) **118** with cyanogen bromide did not give the hypothetical cyclization product, 7*H*-1,2,4-triazolo[4,3-*b*][1,2,4]triazole **119** but afforded an unexpected product identified as 2,3,5,6-tetraamino-9-imino-3*H*,9*H*-bis[1,2,4]triazolo[1,5-*a*:5',1'-*d*][1,3,5]triazinium bromide **120** (Scheme 47).⁵⁷




Scheme 47

Heterocyclization of 3,4,5-triamino[1,2,4]triazole **118** with 4-nitrobenzaldehyde and cyclohexanone afforded 2,3-diamino-9-(4-nitrophenyl)-5,6,7,8-tetrahydro[1,2,4]triazolo[5,1-*b*]quinazoline **121** (Scheme 48).⁵⁸




Scheme 48

In a similar manner, tetrahydro[1,2,4]triazolo[5,1-*b*]quinazolin-8-ones **122** were prepared from the reaction of 3,4,5-triamino[1,2,4]triazole **118** with aldehydes and dimedone. In case of $4\text{-BrC}_6\text{H}_4\text{CHO}$, compound **123** was isolated as by product in low yield (Scheme 49).⁵⁸



Scheme 49


Cyclocondensation of diaminotriazole **124** with β -chlorocinnamaldehyde **125** and 2-chloro-3-formylquinolines **126** in the presence of a catalytic amount of *p*-TsOH produced triazolotriazepines **127** and quinolino[3,2-*f*][1,2,4]triazolo[4,3-*b*][1,2,4]triazepines **128** (Scheme 50).^{59,60}



Scheme 50

5. Heterocyclization with diaminothiazoles


Condensation of 1-amino-2-iminonaphtho[1,2-*d*]thiazole **129** with some α -ketocarboxylic acid and their esters in methanol gave the open chain product **130**, however, when the reaction was performed in glacial acetic acid or DMF, cyclic products identified as 10-alkyl/aryl/heteroaryl/aralkyl-9*H*-naphtho[1',2':4,5]-thiazolo[3,2-*b*][1,2,4]triazin-9-ones **131** were obtained (Scheme 51).⁶¹



Scheme 51


6. Heterocyclization with diaminopyrimidines

Cyclocondensation of triaminopyrimidinethione **132** with pyruvic acid in glacial acetic acid gave pyrimido[3,4-*b*][1,2,4]triazine derivative **133** (Scheme 52).⁶²




Scheme 52

1,3-Diphenyl-7-methyl-1H-pyrazolo[3',4':4,5]pyrimido[1,6-b][1,2,4]triazepin-5(6H)-one **135** was prepared by condensation of 5-amino-1,3-diphenyl-4,5-dihydro-4-imino-1H-pyrazolo[3,4-d]pyrimidine **134** with ethyl acetoacetate. Reaction of compound **135** with diazotized aromatic amines produced the arylazo derivatives **136**. The latter compounds were also obtained from the reaction of compound **134** with ethyl 2-arylhrazono-3-oxobutanoate (Scheme 53).⁶³




Scheme 53

Furo[3,2-*e*][1,2,4]triazolo[1,5-*c*]pyrimidines **138** were prepared by reaction of 5,6-di-(2-furyl)-3*H*,4*H*-4-imino-2-methylfuro[2,3-*d*]pyrimidin-3-amine **137** with triethyl orthoformate, diethyloxalate, ethyl cyanoacetate, cyanogen bromide and carbon disulfide, respectively. Also, condensation of compound **137** with isatine gave the condensation products 12,13-di-(2-furyl)-9-methylfuro[2',3':5,6]-pyrimido[3,4-*b*][2,3-*e*]indolo[1,2,4]triazine **139** (Scheme 54).⁶⁴




Scheme 54

Cyclocondensation of compound **140** with an excess formic acid, acetyl chloride and cyanogen bromide gave the corresponding 8,10-diphenyl-11-(4-chlorophenyl)-pyrazolo[4',3':5,6]pyrano[3,2-*e*][1,2,4]-triazolo[1,5-*c*]pyrimidine derivatives **141** (Scheme 55).⁶⁵




Scheme 55

The polyfused [1,2,4]triazolo[1,5-*c*]pyrimidines **143** were prepared by reaction of 9-amino-7-(4-chlorophenyl)-8,9-dihydro-8-imino-6*H*,7*H*-[1]benzopyrano[3,4:5,6]pyrano[2,3-*d*]pyrimidin-6-ones **142** with triethyl orthoformate, acetyl chloride, benzoyl chloride, diethyl oxalate and ethyl cyanoacetate. Also, treating **142** with carbon disulfide and methyl chloroformate in alcoholic potassium hydroxide solution gave [1,2,4]triazolo[1,5-*c*]pyrimidines **144**. Also, reaction of **142** with ethyl chloroacetate afforded the 15-(methyl/4-chlorophenyl)-3,4-dihydro-2*H*,14*H*,15*H*-[1]benzopyrano[3',4':5,6]pyrano[2,3-*d*]pyrimido-[1,6-*b*][1,2,4]triazine-3,14-dione **145** via elimination of EtOH and HCl (Scheme 56).^{66,67}




Scheme 56

Condensation of ethyl 3-amino-4-imino-3,4,5,6,7,8-hexahydropyrido[4,3:4,5]thieno[2,3-*d*]pyrimidine-7-carboxylate **146** with triethyl orthoformate, acetic anhydride and phenylisothiocyanate gave the corresponding ethyl pyrido[4,3:4,5]thieno[2,3-*d*]-1,2,4-triazolo[3,2-*f*]pyrimidine-8(7*H*)-carboxylate **147**. Also, condensation of **146** with chloroacetyl chloride in DMF/TEA gave 4-oxo-2,3,4,8,10,11-hexahydropyrido[4,3:4,5]thieno[2,3-*d*]-1,2,4-triazolo[3,2-*f*]pyrimidine-9-carboxylate **148** (Scheme 57).⁶⁸




Scheme 57

During an investigation of the reactions of indole-2,3-dione with 2,3-diamino-4(3*H*)quinazolinone **149** in various solvents, a mixture of products were obtained. This reaction when carried out in ethanol containing catalytic amounts of acetic acid yielded several products obtained *via* 3-[2-amino-4-quinazolinone)imino]-2*H*-indol-2-one **150**. Boiling of the latter compound in ethanolic KOH gave 1,2,4-triazino[3,2-*b*]quinazoline-2,6-dione **151**, while in ethanol containing a few drops of piperidine it yielded spiro[3*H*-indol-3',2'(1*H*)] [1,2,4]triazolo[5,1-*b*]quinazoline-2,9-dione **152**. Also, boiling **150** in glacial acetic acid afforded indolo[2',3':5,6][1,2,4]triazino[3,2-*b*]quinazolin-14-one **153** (Scheme 58).⁶⁹



Scheme 58

On the other hand, treatment of *N*-acetylisatine with 2,3-diamino-4(3*H*)quinazolinone **149** led to *N*-2-(triazinoquinazolinone)phenyl acetamide **154** under the same conditions (Scheme 59).⁷⁰




Scheme 59

7. Heterocyclization with diaminotriazine

Condensation of 3,4-diamino-4,5-dihydro-1,2,4-triazin-5-ones **155** with acetone in the presence of weak

organic acid gave 1,2,3,7-tetrahydro[1,2,4]triazolo[3,2-*c*][1,2,4]triazin-7-ones **156**.⁷¹ While, the reaction of compound **155** with acetaldehyde and diethyl phosphite in THF in the presence of sodium hydride as a catalyst gave [1,2,4]triazino[4,3-*b*][1,2,4,5]triazaphosphinine derivative **157** (Scheme 60).⁷²



Scheme 60

REFERENCES

1. M. A. Ibrahim, *Heterocycles*, 2011, **83**, 2689.
2. J. W. Fronabarger, R. D. Chapman, and R. D. Gilardi, *Tetrahedron Lett.*, 2006, **47**, 7707.
3. R. S. Bhosale, S. R. Sarda, S. S. Ardhapure, W. N. Jadhav, S. R. Bhusare, and R. P. Pawar, *Tetrahedron Lett.*, 2005, **46**, 7183.
4. R. A. Timm, J. A. Bonacin, A. L. Formiga, and B. H. E. Toma, *J. Braz. Chem. Soc.*, 2008, **19**, 287.
5. J. T. Kuethe, A. Wong, and I. W. Davies, *J. Org. Chem.*, 2004, **69**, 7752.
6. M. R. Ganjali, P. Norouzi, F. Faridbod, M. Ghorbani, and M. Adib, *Anal. Chim. Acta*, 2006, **569**, 35.
7. B. Gustaffon, M. Hakansson, A. T. Hutton, J. R. Moss, and S. Jagner, *Inorg. Chim. Acta*, 2005, **358**, 1327.
8. H. Na, D. C. Lee, J. W. Lim, J. H. Choi, J. C. Byun, and Y. C. Park, *Polyhedron*, 2002, **21**, 917.
9. D. M. Boghaei and S. Mohebi, *Tetrahedron*, 2002, **58**, 5357.
10. T. Jeewoth, M. G. Bhowon, and H. L. K. Wah, *Transition Met. Chem.*, 1999, **24**, 445.
11. M. H. Elnagdi, S. A. Ghazlan, F. M. Abdel-Razik, and A. S. Maghraby, *Chem. Synop.*, 1991, **5**, 116.
12. F. A. Attaby, S. M. Eldin, and F. M. Abdel-Razik, *Phosphorus, Sulfur Silicon Relat. Elem.*, 1995, **106**, 21.
13. K. A. Asadov, R. N. Burangulova, F. H. Guseninov, R. Z. Gilmanov, and I. P. Phaljachov, *Chem. Heterocycl. Compds.*, 2003, **39**, 392.
14. M. Miletin, J. Hartl, M. Dolezal, Z. Odlerova, K. Kralova, and M. Machacek, *Molecules*, 2000, **5**, 208.
15. A. Abdel-Rahman, E. A. Bakhite, and E. A. Al-Taifi, *J. Chin. Chem. Soc.*, 2002, **49**, 223.
16. C. S. Rao, V. Venkaleswarlu, and G. Achaiyah, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 2134.
17. A. A. A. Al-Najjar, S. A. Amer, M. Riad, I. Elghamy, and M. H. Elnagdi, *J. Chem. Res.*, 1996, 296.

18. Y. Yamada, H. Yasuda, and A. Takayama, [Heterocycles](#), 1998, **48**, 1185.
19. Y. Luo and Y. Hu, [Arch. Pharm.](#), 2006, **339**, 262.
20. E. R. El-Sawy, A. H. Mandour, I. E. Islam, and H. M. Abo-Salem, *Egypt. J. Chem.*, 2008, **51**, 523.
21. X. Tao and Y. Hu, [Med. Chem.](#), 2010, **6**, 65.
22. G. Zhang and Y. Hu, [J. Heterocycl. Chem.](#), 2007, **44**, 919.
23. M. S. Mohamed, M. E. A. Zaki, N. M. Khalifa, and Y. M. Zohny, [Heterocycl. Commun.](#), 2008, **14**, 345.
24. H. Berner and H. Reinshagen, [Monatsh. Chem.](#), 1975, **106**, 1059.
25. M. A. Barys, E. A. El Rady, and F. M. Abd El Latif, [J. Heterocycl. Chem.](#), 2008, **45**, 773.
26. A. H. Mandour, O. A. Fathalla, and W. M. Basyouni, *Modelling, Measurement Control C*, 2000, **61**, 53.
27. E. Huntsman and J. Balsells, [Eur. J. Org. Chem.](#), 2005, **17**, 3761.
28. G. L. Rusinov, V. L. Mikhailov, and O. N. Chupakhin, [Russ. Chem. Bull.](#), 1999, **48**, 1391.
29. A. H. M. Hussein, *Z. Naturforsch. B: Chem. Sci.*, 1998, **53**, 488.
30. A. H. M. Hussein, [Heteroatom Chem.](#), 1997, **8**, 1.
31. A. M. Hussein, S. M. Sherif, and A. A. Atalla, [Monatsh. Chem.](#), 1996, **127**, 1153.
32. A. E. H. Ismail, *Pharmazie*, 1996, **51**, 982.
33. C. Yamazaki, Y. Miyamoto, and H. Sakima, [J. Chem. Soc., Perkin Trans. 1](#), 1994, **7**, 825.
34. M. J. Callejo, P. Lafuente, N. Martin-Leon, M. Quinteiro, C. Seoane, and J. L. Soto, [J. Chem. Soc., Perkin Trans. 1](#), 1990, **6**, 1687.
35. W. R. Abdel-Monem, *Chem. Pap.*, 2004, **58**, 276.
36. T. E. Ali and M. A. Ibrahim, [J. Braz. Chem. Soc.](#), 2010, **21**, 1007.
37. W. R. Abdel-Monem, [Eur. J. Chem.](#), 2010, **3**, 168.
38. A. A. Harb, *Chem. Pap.*, 2004, **58**, 260.
39. W. M. Basyouni, *Acta Chem. Slov.*, 2003, **50**, 223.
40. M. Abdel-Megid, M. A. Ibrahim, Y. Gabr, N. M. El-Gohary, and E. A. Mohamed, [J. Heterocycl. Chem.](#), 2013, **50**, 615.
41. M. A. Ibrahim, R. M. Abdel-Rahman, A. M. Abdel-Halim, S. S. Ibrahim, and H. A. Allimony, [ARKIVOC](#), 2008, **xvi**, 202.
42. A. M. El-Kazak and M. A. Ibrahim, [ARKIVOC](#), 2013, **iii**, 282.
43. M. Suresh, S. B. Jonnalagadda, and C. V. Rao, *Oriental J. Chem.*, 2011, **27**, 127.
44. M. Abdel-Megid, [Chem. Heterocycl. Compds.](#), 2009, **45**, 1523.
45. M. A. Ibrahim, R. M. Abdel-Rahman, A. M. Abdel-Halim, S. S. Ibrahim, and H. A. Allimony, [J. Braz. Chem. Soc.](#), 2009, **20**, 1275.

46. H. M. Hassanin and S. M. El-edfawy, [Heterocycles](#), 2012, **85**, 2421.
47. V. Y. Sosnovskikh and V. S. Moshkin, [Chem. Heterocycl. Compds.](#), 2012, **48**, 139.
48. H. M. Hassanin, M. A. Ibrahim, and Y. A. Alnamer, *Turkish J. Chem.*, 2012, **36**, 682.
49. M. A. Ibrahim, M. Abdel-Megid, and N. M. El-Gohary, [J. Braz. Chem. Soc.](#), 2011, **21**, 1130.
50. A. J. Blake, D. Clarke, W. Mares, and H. McNab, [Org. Biomol. Chem.](#), 2003, **1**, 4268.
51. N. N. Kolos, V. D. Orlov, B. V. Paponov, and O. V. Shishkin, [Chem. Heterocycl. Compds.](#), 1999, **35**, 1207.
52. Y. Yamada, H. Yasuda, Y. Yoshihara, and K. Yoshizawa, [J. Heterocycl. Chem.](#), 1999, **36**, 1317.
53. V. P. Gnidgets, I. Kherson, M. V. Povstyanoi, P. B. Kapov, and I. I. Grandberg, *Ukr. Khim. Zh.*, 1988, **54**(5), 531; *Chem. Abstr.*, 1989, **110**, 75450v.
54. M. A. E. Shaban, A. Z. Nasr, and A. E. A. Morgaan, [Farmaco](#), 1999, **54**, 800.
55. J. Daunis and H. Loper, [J. Org. Chem.](#), 1977, **42**, 1018.
56. P. V. G. Reddy, Y. B. R. Kiran, C. S. Reddy, and C. D. Reddy, [Chem. Pharm. Bull.](#), 2004, **52**, 307.
57. J. W. Fronabarger, R. D. Chapman, and R. D. Gilardi, [Tetrahedron Lett.](#), 2006, **47**, 7707.
58. V. V. Lipson, S. M. Desenko, V. V. Borodina, M. G. Shirobobjkova, O. V. Shishkin, and R. J. Zubatyuk, [Russ. Chem. Bull.](#), 2006, **55**, 1224.
59. M. Gupta, [J. Heterocycl. Chem.](#), 2007, **44**, 1023.
60. M. Gupta, S. Paul, and R. Gupta, *Indian J. Chem.*, 2010, **49B**, 475.
61. L. Kang-Chain, S. Bi-Jane, and L. Chuen-Hsiang, [J. Heterocycl. Chem.](#), 1993, **30**, 1331.
62. T. G. Kokjhaova, L. N. Dianoova, N. V. Volhova, D. S. Anisimova, G. M. Anoshina, and G. V. Svechikova, *Khim-Farm. Zh*, 1988, **22**, 805; (*Chem. Abstr.*, 1988, **110**, 114803J).
63. A. S. Shawali and T. A. Farghaly, [Tetrahedron](#), 2009, **65**, 644.
64. N. A. Hassan, [Molecules](#), 2000, **5**, 826.
65. M. E. Zaki, [Molecules](#), 1998, **3**, 71.
66. A. M. El-Agrody, M. S. Abd El-Latif, N. A. El-Hady, A. H. Fakery, and A. H. Bedair, [Molecules](#), 2001, **6**, 519.
67. A. H. Bedair, N. A. El-Hady, M. S. Abd El-Latif, A. H. Fakery, and A. M. El-Agrody, [Farmaco](#), 2000, **55**, 708.
68. E. K. Ahmed, M. Ameen, and F. F. Abdel-Latif, [Phosphorus, Sulfur Silicon Relat. Elem.](#), 2006, **181**, 497.
69. K. C. Joshi, A. Dandia, and S. Khanna, *Indian J. Chem.*, 1992, **31B**, 105.
70. A. Dandia, S. Sanan, S. Bhagat, and K. Arya, 6th International Electronic Conference on Synthetic Org. Chem. (ECSOC-6), 2002.
71. D. Wilfried, D. Karlfried, and E. Ludwig, (Bayer A.-G.) Ger. Offen. 1,965,937; (*Chem. Abstr.*, 1971,

75, 129842d).

72. T. E. Ali, [*Eur. J. Med. Chem.*, 2009, **44**, 4539](#).



Magdy Ahmed Mohamed Ibrahim was born in Cairo, Egypt, in 1977. He received his B.Sc. degree in physics and chemistry (1999) and his M.Sc. degree in organic chemistry (2004). Also, he received his Ph.D. degree (2007) from the organic division, Department of Chemistry, Faculty of Education, Ain Shams University. He won the award for the best research article in heterocyclic chemistry field at the Egyptian universities and research centers in 2010. In 2012 he was promoted to assistant professor in organic chemistry. His research is focused on the synthesis of new heterocyclic compounds containing chromone, pyridine, quinolin-2(1*H*)-one and 1,2,4-triazine. In addition, He has published about 35 scientific papers in international scientific journals.



Nasser Mohamed Ahmed El-Gohary was born in Giza, Egypt, in 1980. He received his B.Sc. degree in physics and chemistry (2001) and his M.Sc. degree in organic chemistry (2009). Also, he received his Ph.D. degree (2012) from the organic division, Department of Chemistry, Faculty of Education, Ain Shams University. His research is focused on the synthesis of new heterocyclic compounds containing chromone nucleus.