

ALKATRIENYL SULFOXIDES AND SULFONES. PART VI.  
CHELETROPIC ADDITION OF SULFUR DIOXIDE TO 1- AND  
3-VINYLLALLENYL SULFOXIDES AND SULFONES<sup>1,2</sup>

Valerij Ch. Christov\* and Ivaylo K. Ivanov


Department of Chemistry, University of Shumen, BG-9700 Shumen, Bulgaria  
E-Mail: vchristo@shu-bg.net

**Abstract** – The reaction of 1- and 3-vinylallenyl sulfoxides and sulfones with sulfur dioxide in 1,2-dichloroethane at reflux proceeds as a cheletropic addition leading to the formation of 2-isopropylidene-3-sulfinyl(sulfonyl)- or 2-sulfinyl(sulfonyl)methylidene-substituted 2,5-dihydrothiophene 1,1-dioxides in very good yields.

Conjugated dienes are known to undergo cheletropic additions<sup>3</sup> with sulfur dioxide to generate the corresponding 2,5-dihydrothiophene 1,1-dioxides (sulfolenes),<sup>4</sup> or to generate polymers (polysulfones).<sup>5</sup> At low temperature and in the presence of a protic or Lewis acid catalyst, simple acyclic alkyl-substituted 1,3-dienes that can adopt the *s-cis* conformation add reversibly to SO<sub>2</sub> by hetero-Diels-Alder additions to generate the corresponding 3,6-dihydro-1,2-oxathiine 2-oxides (sultines).<sup>6,7</sup> The latter are unstable above -50 °C and undergo fast cycloreversion to liberate the starting 1,3-dienes and SO<sub>2</sub>, which can then undergo the expected cheletropic addition at higher temperature. The competition between the hetero-Diels-Alder and the cheletropic additions of sulfur dioxide strongly depends on the nature of the substituents of the 1,3-dienes.<sup>8</sup> Reactions of sulfur dioxide with conjugated diallenes take place at room temperature with formation of 2,5-bis(alkylidene)-2,5-dihydrothiophene 1,1-dioxides.<sup>9</sup> Reactions of sulfur dioxide with vinylallenes, in particular, with vinylallenyl sulfoxides and sulfones have not been investigated to date. During our previous works concerning electrophile-induced cyclization reactions of alkatrienyl sulfoxides and sulfones,<sup>10a-10d</sup> we were able to show that 1- and 3-vinylallenyl sulfoxides and sulfones are readily accessible by [2,3] sigmatropic rearrangement of the corresponding 1- and 3-vinylpropargylic sulfenates and sulfates, formed in the reactions of the corresponding  $\alpha$ -alkynols with sulfonyl or sulfinyl chlorides.<sup>10a-10d</sup> In view of the advantages of vinylallenes as diene component in Diels-Alder reaction,<sup>10e</sup> we therefore initiated a study of their use in cheletropic addition. The results of this work are presented here.


We initiated this study with the cheletropic addition of sulfur dioxide to 5-methyl-3-(phenylsulfinyl)hexa-1,3,4-triene (**1a**). The reaction conditions have been optimized in order to obtain better yields. The reaction in benzene, toluene, xylene, CCl<sub>4</sub>, CH<sub>2</sub>Cl<sub>2</sub>, and CHCl<sub>3</sub> as a solvent resulted in low product yields and mainly recovered the starting material or polymeric residue. According to results in the screening solvents, 1,2-dichloroethane was found to be the best. Thus, optimization of the reaction conditions was conducted in ClCH<sub>2</sub>CH<sub>2</sub>Cl and finally it was found that when sulfur dioxide was bubbled through a stirred boiling solution of the vinylallene (**1a**), fast darkening of the reaction mixture took place and after 3.5 h no characteristic band for the allenic bond in the IR spectrum of the mixture was observed. The resulting 2,5-dihydrothiophene 1,1-dioxide (sulfolene) (**3a**) was isolated by preparative TLC in 78% yield. Note that at room temperature the two reactants still interacted (much more slowly) with the formation of the compound (**3a**).

To establish the generality of this method, the cheletropic addition of SO<sub>2</sub> to the corresponding 1-vinylallenyl sulfone (**1b**) was examined under the optimized conditions and 3-(methylsulfonyl)-2,5-dihydrothiophene 1,1-dioxide (**4b**) was isolated in 79% yield after bubbling with SO<sub>2</sub> and heating at reflux for 4 h (Scheme 1).



Scheme 1

Interestingly, this protocol can also be successfully applied to the corresponding cheletropic addition of SO<sub>2</sub> to the 3-vinylallenic starting materials. We have carried out the reaction of the 1-phenylsulfinyl- and 1-methylsulfonyl-substituted penta-1,2,4-trienes (**2a**) and (**2b**) with sulfur dioxide leading to (*E*)-2-phenylsulfinyl- or (*E*)-2-methylsulfonyl-methylidene-2,5-dihydrothiophene 1,1-dioxides (**4a**) and (**4b**) (Scheme 2). Although it was anticipated that the olefinic proton of the (*E*)-isomer would be observed



Scheme 2

downfield from the corresponding proton of the (*Z*)-isomer,<sup>13</sup> with the chemical shift value<sup>12</sup> ( $\delta = 7.32$  ppm for **4a** and 7.35 ppm for **4b**) alone we cannot determine whether dihydrothiophenes **4** is the (*E*)- or (*Z*)-isomer. On the other hand, in the <sup>13</sup>C NMR spectra the chemical shift value of the two carbon atoms of the exocyclic double bond of the (*E*)-isomer would be observed upfield from the corresponding carbons of the (*Z*)-isomer.<sup>13</sup> The obtained chemical shift value<sup>12</sup> for these carbons ( $\delta = 136.2$  and 155.5 ppm for **4a** and 137.3 and 149.2 ppm for **4b**) suggests that the olefinic proton is situated *cis* to the sulfonic group in the ring. Thus, the structure of **4** was determined to be (*E*).

Reaction times longer than 4 hours decrease the yields due to polymerization of the starting vinylallenic materials or decomposition of the products. The only product obtained in each case was the 2,5-dihydrothiophene 1,1-dioxides (**3a,b**) and (**4a,b**) – in all cases no traces of the corresponding 3,6-dihydro-1,2-oxathiine 2-oxides (sultines) as products of the hetero-Diels-Alder reaction or any other products were detected. Structural assignments of the new dihydrothiophenic compounds (**3**) and (**4**) are based on <sup>1</sup>H and <sup>13</sup>C NMR spectral data, IR spectra as well as elemental analyses.<sup>11,12</sup> In particular, on one hand, the protons on C-5 of the ring in <sup>1</sup>H NMR spectra resonate at  $\delta = 3.79$ -4.18 ppm as two double doublets and the proton on C-4 appears as triplet, thus accounting for the depicted regiochemistry of the cycloaddition. On the other hand, the <sup>13</sup>C NMR spectra of these products (**3**) and (**4**) also showed a complex signals of the sp<sup>2</sup> carbons ( $\delta = 120.7$ -155.5 ppm) and the sp<sup>3</sup> C-5 ( $\delta = 47.2$ -48.4 ppm), characteristic for newly-formed 2,5-dihydrothiophenes.

In conclusion, we have developed an efficient synthesis of 2- and 3-sulfinyl(sulfonyl)-substituted 2,5-dihydrothiophene 1,1-dioxides by cheletropic addition of sulfur dioxide to 1- and 3-vinylallenyl sulfoxides and sulfones. This study reveals the potential of the cycloaddition reactions of vinylallenyl sulfoxides and sulfones to selectively construct exocyclic double bond on six-<sup>2</sup> or five-membered rings. Further expansion and applications of this methodology are in progress and will be reported in due course.

## ACKNOWLEDGEMENTS

Support from the Research Fund of the University of Shumen (Projects No. 12 / 2003 and No. 16 / 2004) is acknowledged. The first author would like to thank two anonymous referees for their useful comments.

## REFERENCES AND NOTES

1. Dedicated to Professor Dr. Toru Minami of Kyushu Institute of Technology, Kitakyushu, Japan on the occasion of his 65<sup>th</sup> birthday.
2. For Part V, see: V. Ch. Christov and I. K. Ivanov, *Synth. Commun.*, 2004, **34**, issue 21, in press.
3. (a) R. B. Woodward and R. Hoffmann, 'The Conservation of Orbital Symmetry', Academic Press, Inc., New York, 1970; (b) S. D. Turk and R. L. Coob, in '1,4-Cycloaddition Reactions', ed. by J. Hamer, Academic Press, Inc., New York, 1967, p.13; (c) M. J. S. Dewar, 1984, *J. Am. Chem. Soc.*, 1984, **106**, 209 and proceeding papers.
4. (a) G. De Bruin, *Koninkl. Ned. Akad. Wetenschap. Proc.*, 1914, **17**, 585; (b) E. Eigenberger, *J. Prakt.*

- Chem.*, 1930, **127**, 307; (c) H. J. Backer and J. Strating, *Rec. Trav. Chim. Pays-Bas*, 1934, **53**, 523; (d) E. de R. van Zuydewijn and J. Boeseken, *Rec. Trav. Chim. Pays-Bas*, 1934, **53**, 673; (e) H. Staudinger and B. Ritzenthaler, *Ber.*, 1935, **68**, 455; (f) H. J. Backer and J. Strating, *Rec. Trav. Chim. Pays-Bas*, 1943, **62**, 815.
- See for example: D. Masilamani, E. H. Manahan, J. Vitrone, and M. M. Rogic, *J. Org. Chem.*, 1983, **48**, 4918.
  - For the hetero-Diels-Alder addition of sulfur dioxide to 1,3-dienes, see: (a) B. Deguin and P. Vogel, *J. Am. Chem. Soc.*, 1992, **114**, 9210; (b) B. Deguin and P. Vogel, *Tetrahedron Lett.*, 1993, **34**, 6269; (c) F. Monnat, P. Vogel, R. Meana, and J. A. Sordo, *Angew. Chem., Int. Ed.*, 2003, **42**, 3924; (d) E. Roversi, R. Scoppelliti, E. Solari, R. Estoppey, P. Vogel, P. Brana, B. Mendendez, and J. A. Sordo, *Chem. Eur. J.*, 2002, **8**, 1336; (e) D. Markovic, E. Roversi, R. Scoppelliti, P. Vogel, and R. Meana, *Chem. Eur. J.*, 2003, **9**, 4911.
  - For the competition between hetero-Diels-Alder addition and cheletropic addition of sulfur dioxide to 1,3-dienes, see: (a) F. Monnat, P. Vogel, and J. A. Sordo, *Helv. Chim. Acta*, 2002, **85**, 712; (b) E. Roversi, F. Monnat, P. Vogel, K. Schenk, and P. Roversi, *Helv. Chim. Acta*, 2002, **85**, 733; (c) E. Roversi and P. Vogel, *Helv. Chim. Acta*, 2002, **85**, 761.
  - For mechanisms of the hetero-Diels-Alder and cheletropic additions of sulfur dioxide to 1,3-dienes, see: (a) D. Suarez, E. Iglesias, T. L. Sordo, and J. A. Sordo, *J. Phys. Org. Chem.*, 1996, **9**, 17; (b) F. Monnat, P. Vogel, V. M. Rayon, and J. A. Sordo, *J. Org. Chem.*, 2002, **67**, 1882.
  - (a) K. Kleveland and L. Skattebol, *J. Chem. Soc., Chem. Commun.*, 1973, 432; (b) K. Kleveland and L. Skattebol, *Acta Chem. Scand.*, 1975, **B 29**, 827.
  - Our previous reports on alkatrienyl sulfoxides and sulfones, see: (a) Part I, see: V. Ch. Christov and I. K. Ivanov, *Phosphorus, Sulfur, Silicon*, 2002, **177**, 2445; (b) Part II, see: V. Ch. Christov and I. K. Ivanov, *Sulfur Lett.*, 2002, **25**, 191; (c) Part III, see: V. Ch. Christov and I. K. Ivanov, *Heterocyclic Commun.*, 2003, **9**, 629. (d) For Part IV, see: V. Ch. Christov and I. K. Ivanov, *Phosphorus, Sulfur, Silicon*, 2004, in press; (e) See: Ref.<sup>2</sup>
  - (a) *3-Phenylsulfinyl-2-isopropylidene-2,5-dihydrothiophene 1,1-dioxide (3a)*: Yellow oil, 78 % yield. TLC: ethyl acetate : hexane = 1 : 1. *Anal.* Calcd for C<sub>13</sub>H<sub>14</sub>O<sub>3</sub>S<sub>2</sub>: C 55.29, H 5.00, S 22.71; Found, C 55.47, H 4.94, S 22.63. IR (film): 1046, 1143, 1288, 1599-1640. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 250 MHz): δ = 2.07 (s, 3H, Me<sub>a</sub>), 2.21 (s, 3H, Me<sub>b</sub>), 3.72, 3.87 (2xddd, <sup>3</sup>J<sub>HH</sub> 2.9 Hz, <sup>2</sup>J<sub>HH</sub> 12.0 Hz, 2H, CH<sub>2</sub>), 7.2 (t, <sup>3</sup>J<sub>HH</sub> 2.9 Hz, 1H, =C-H), 7.53-8.21 (m, 5H, Ph). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 50 MHz): δ = 18.4, 20.9, 47.3, 126.1, 129.5, 130.4, 136.2, 142.9, 145.8, 146.5, 153.3. (b) *3-Methylsulfonyl-2-isopropylidene-2,5-dihydrothiophene 1,1-dioxide (3b)*: Yellow oil, 79 % yield. TLC: ethyl acetate : heptane = 1 : 2. *Anal.* Calcd for C<sub>8</sub>H<sub>12</sub>O<sub>4</sub>S<sub>2</sub>: C 40.66, H 5.12, S 27.14; Found, C 40.61, H 5.03, S 26.98. IR (film): 1141, 1326, 1600-1648. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 250 MHz): δ = 2.02 (s, 3H, Me<sub>a</sub>), 2.10 (s, 3H, Me<sub>b</sub>), 3.07 (s, 3H, SO<sub>2</sub>Me), 3.77, 3.98 (2xddd, <sup>3</sup>J<sub>HH</sub> 2.9 Hz, <sup>2</sup>J<sub>HH</sub> 12.1 Hz, 2H, CH<sub>2</sub>), 7.23 (t, <sup>3</sup>J<sub>HH</sub> 2.9 Hz, 1H, =C-H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 50 MHz): δ = 22.2, 24.4, 36.7, 47.7, 126.3, 130.4, 141.6, 153.8.
  - (a) *(E)-2-Phenylsulfinylmethylidene-3-methyl-2,5-dihydrothiophene 1,1-dioxide (4a)*: Yellow oil, 72 % yield. TLC: ethyl acetate : heptane = 1 : 1. *Anal.* Calcd for C<sub>12</sub>H<sub>12</sub>O<sub>3</sub>S<sub>2</sub>: C 53.71, H 4.51, S 23.90; Found, C 53.83, H 4.48, S 24.08. IR (film): 1038 (S=O), 1122 (ν SO<sub>2</sub>), 1306 (ν SO<sub>2</sub>), 1581-1684 (C=C). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 250 MHz): δ = 2.24 (s, 3H, Me), 4.11, 4.24 (2xddd, <sup>3</sup>J<sub>HH</sub> 3.1 Hz, <sup>2</sup>J<sub>HH</sub> 12.7 Hz, 2H, CH<sub>2</sub>), 6.18 (t, <sup>3</sup>J<sub>HH</sub> 3.1 Hz, 1H, =C<sup>4</sup>-H), 7.32 (s, 1H, =C-H), 7.51-8.45 (m, 5H, Ph). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 50 MHz): δ = 21.5, 47.2, 120.7, 125.3, 127.6, 130.1, 136.2, 139.8, 150.0, 155.5. (b) *(E)-2-Methylsulfonylmethylidene-3-methyl-2,5-dihydrothiophene 1,1-dioxide (4b)*: Yellow oil, 76 % yield. TLC: ethyl acetate : heptane = 5 : 1. *Anal.* Calcd for C<sub>7</sub>H<sub>10</sub>O<sub>4</sub>S<sub>2</sub>: C 37.82, H 4.53, S 28.85; Found, C 37.73, H 4.59, S 28.67. IR (film): 1126 (ν SO<sub>2</sub>), 1297 (ν SO<sub>2</sub>), 1590-1640 (C=C). <sup>1</sup>H NMR (CDCl<sub>3</sub>, 250 MHz): δ = 2.27 (s, 3H, Me), 2.75 (s, 3H, SO<sub>2</sub>Me), 3.93, 4.13 (2xddd, <sup>3</sup>J<sub>HH</sub> 3.2 Hz, <sup>2</sup>J<sub>HH</sub> 11.9 Hz, 2H, CH<sub>2</sub>), 6.01 (t, <sup>3</sup>J<sub>HH</sub> 3.2 Hz, 1H, =C<sup>4</sup>-H), 7.35 (s, 1H, =C-H). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 50 MHz): δ = 21.2, 43.7, 48.4, 129.9, 137.3, 143.8, 149.2.
  - Advanced Chemistry Development, Inc. (ACD / Labs) (<http://www.acdlabs.com>).