

HETEROCYCLES, Vol. 68, No. 11, 2006, pp. 2343 - 2347. © The Japan Institute of Heterocyclic Chemistry
Received, 7th August, 2006, Accepted, 20th September, 2006, Published online, 22nd September, 2006. COM-06-10859

***N,N,N',N'*-TETRABROMOBENZENE-1,3-DISOLFONAMIDE (TBBDS) AS AN EFFICIENT PROMOTER FOR ONE-POT CONVERSION OF *N*-ARYLGLYCINES TO SYDNONES IN THE PRESENCE OF NaNO₂/Ac₂O UNDER NEUTRAL CONDITIONS**

Davood Azarifar,*^a Hassan Ghasemnejad-Bosra,^a Ramin Ghorbani-Vaghei,^a and Mahmood Tajbaksh^b

^aDepartment of Chemistry, School of Science, Bu-Ali Sina University, Postal Code 65174, Hamadan, Iran. ^bDepartment of Chemistry, Faculty of Science, Mazandaran University, Babolsar, Iran. *Corresponding author; Fax +98(811)8272404; E-mail: azarifar@basu.ac.ir


Abstract– *N,N,N',N'*-Tetrabromobenzene-1,3-disulfonamide (TBBDS)-promoted one-pot conversion of various *N*-arylglycines to sydnones using a combination of NaNO₂ and Ac₂O has been achieved efficiently through *N*-nitrosation followed by cyclization in high yields (85-95%) under mild and neutral conditions.

Sydnones (*cf.* **2**) are unique members of the class of heterocyclic compounds known as mesoionic.¹ Sydnones were first prepared by Earl and Mackney in 1935,² and the greatest interest in them, ever since, stems from their biological activity as antibacterial,³ antitumour,⁴ antimalarial,⁵ anti-inflammatory,⁶ and antihypertensive⁷ agents. Sydnones also undergo a variety of transformations including electrophilic aromatic substitution (at the 4-position),⁸ 1,3-dipolar cycloaddition reactions to form pyrazoles or related species,⁹ and cleavage to hydrazines² or heterocycles¹⁰ when treated with HCl.

Sydnones are intrinsically neutral substances that are normally prepared by dehydrative cyclization of *N*-nitrosamino acids.¹¹ *N*-Nitrosamino acids used in the synthesis of sydnones are themselves prepared from *N*-nitrosation of amino acids. *N*-Nitrosation is a well-known reaction in organic synthesis¹² that is usually accomplished by nitrous acid generated from the treatment of sodium nitrite with an aqueous mineral acid.¹³

In continuation of our research on various transformations by halogenating agents¹⁴⁻¹⁶ and sydnones,¹⁷⁻¹⁹ and also in order to avoid the drawbacks generally caused by the use of strong acidic media in nitrosation reactions, we wish, herein, to report on the *N,N,N',N'*-tetrabromobenzene-1,3-disulfonamide (TBBDS) as

a more robust and efficient promoter for one-pot conversion of *N*-arylglycines to sydrones under neutral conditions. In this work we have observed that *N,N,N',N'*-tetrabromobenzene-1,3-disulfonamide (TBBDS) can efficiently enhance the conversion of the *N*-arylglycines (**1a-j**) to the sydrones (**2a-j**) using sodium nitrite and acetic anhydride in CH₂Cl₂ in satisfactory yields (85-95 %) (**Scheme 1**, **Table 1**). As shown in the table, the reactions occur satisfactorily within 5-8 h at 0-5 °C. The experimental results indicated that the most effective conversion occurs if the molar *N*-arylglycines/TBBDS ratio is maintained at 4. Longer reaction times are required when lesser amounts of TBBDS are employed. It is important to note that no sydrones were afforded when the reactions were carried out without using any TBBDS in the reaction. In accordance with the previously reported action of acetic anhydride in the cyclization of *N*-nitrosoglycines to sydrones,² and also the crucial role of the acetyl hypobromite, generated from the reaction of acetic anhydride with hypobromous acid during the reaction, in *N*-bromination of sulfonamide,²⁰ we propose a possible mechanism for these reactions as shown in **Scheme 2**, in which the cyclization of the intermediate *N*-nitrosoglycines is, probably, activated by TBBDS.




Scheme 1

Table 1 Conversion of the *N*-arylglycines (**1a-j**) to the sydrones (**2a-j**) with a combination of NaNO₂/Ac₂O in CH₂Cl₂ promoted by TBBDS.

Entry	Product ^a	Ar	Time (h)	Yield (%) ^b	mp (°C)	
					Found	Reported ^{20,21}
1	2a	<i>o</i> -CH ₃ C ₆ H ₄	7.0	92	98	97
2	2b	<i>p</i> -CH ₃ C ₆ H ₄	6.0	90	143	145
3	2c	<i>o</i> -CH ₃ OC ₆ H ₄	6.0	93	96	97
4	2d	<i>p</i> -CH ₃ OC ₆ H ₄	5.5	89	126	125
5	2e	<i>p</i> -NO ₂ C ₆ H ₄	6.0	88	147	148
6	2f	<i>o</i> -NO ₂ C ₆ H ₄	8.0	85	184	184
7	2g	<i>p</i> -ClC ₆ H ₄	7.0	91	114	113
8	2h	2,4-Cl ₂ C ₆ H ₃	7.8	85	94	96
9	2i	<i>p</i> -BrC ₆ H ₄	6.2	93	138	137
10	2j	C ₆ H ₅	5.0	95	134	135

^aAll the isolated products were characterized on the basis of their physical properties and ¹H NMR, ¹³C NMR and IR spectra and by direct comparison with literature data.^{2,8,21,22}

^bIsolated yields.



Scheme 2

EXPERIMENTAL

Chemicals were obtained from Merck and Fluka Chemical Companies. IR spectra were recorded using a Shimadzu 435-U-04 spectrophotometer (KBr pellets) and ^1H NMR and ^{13}C NMR spectra were obtained in CDCl_3 using 90 MHz JEOL FT NMR spectrometer. All melting points were determined on a Büchi 530 melting point apparatus, and reported uncorrected.

Conversion of the *N*-Arylglycines (1a-j) to the Sydnone (2a-j) with $\text{NaNO}_2/\text{Ac}_2\text{O}$ using TBBDS; General Procedure

To a magnetically stirred solution of *N*-arylglycine (**1**) (2 mmol) in CH_2Cl_2 (40 mL), was added *N,N,N',N'*-tetrabromobenzene-1,3-disulfonamide (TBBDS) (0.5 mmol), NaNO_2 (0.17 g, 2.5 mmol) and Ac_2O (0.31 g, 3 mmol) at 0-5 °C. After the complete conversion of the substrate in 5-8 h (Table 1) as monitored by TLC using AcOEt /hexane mixture (1:1), the reaction mixture was poured into water (5 mL), and then solid NaHCO_3 was added cautiously with stirring to remove the remaining glycines. The resulting mixture was filtered, the filtrate was extracted with CH_2Cl_2 , and then the organic layer was separated, dried and evaporated in *vacuo* to leave the solid product (**2**), which was further purified by

recrystallization from EtOH. These products were characterized on the basis of their physical properties and also their IR and NMR spectra with direct comparison with literature data.^{2,8,21,22}

ACKNOWLEDGEMENT

We wish to thank the University of Bu-Ali Sina, Hamadan, Iran, for financial support to carry out this research.

REFERENCES

1. M. Ohta and H. Kato, In *Nonbenzeneoid Aromatics* Vol. 1, ed. by J. P. Snyder; Academic Press: New York, 1969, p. 146; F. H. C. Stewart, *Chem. Rev.*, 1964, **64**, 129.
2. J. C. Earl and A. W. Mackney, *J. Chem. Soc.*, 1935, 899.
3. P. Brooks and J. Walker, *J. Chem. Soc.*, 1959, 4409; E. Ackermann, *Pharmazie*, 1967, **22**, 537 (*Chem. Abstr.*, 1968, **68**, 28, 171).
4. L. B. Kier and E. B. Roche, *J. Pharm. Sci.*, 1967, **56**, 149.
5. D. J. McCaustland, W. H. Burton, and C. C. Cheng, *J. Heterocycl. Chem.*, 1971, **8**, 89.
6. G. Pala, A. Mantegani, G. Coppi, and R. Genova, *Chim. Ther.*, 1969, **4**, 31; (*Chem. Abstr.*, 1969, **71**, 3328); T. Kamiya, Y. Saito, and T. Teraji, Fujisawa Pharm. Co., Ltd., Japanes Patent 7232073, 1972 (*Chem. Abstr.*, 1973, **78**, 42592).
7. Y. Saito and T. Kamiya, Takeda Chem. Ind. Ltd., Japanese Patent 7010510, 1970 (*Chem. Abstr.*, 1970, **73**, 14853).
8. K. Turnbull, *J. Heterocycl. Chem.*, 1985, **22**, 965.
9. R. Huisgen, R. Grashey, H. Gotthardt, and R. Schmidt, *Angew. Chem.*, 1962, **74**, 29.
10. C. R. Gelvin and K. Turnbull, *Helv. Chim. Acta*, 1992, **75**, 1931.
11. C. J. Thoman and D. J. Voaden, *Org. Synth., Coll. Vol. V*; Wiley: New York, 1973, 962.
12. D. L. H. Williams, In *'The Chemistry of Functional Groups, The Chemistry of Amino, Nitroso, Nitro and Related Groups'*, Supplement F₂; ed. by S. Patai; Wiley: Chichester, 1996, pp. 665-682.
13. R. L. Shriner, T. L. Reynold, C. Fuson, D. Y. Curtin, and T. C. Morrill, *The Systematic Identification of Organic Compounds*, 6th ed., Wiley: New York, 1980, pp. 220-223.
14. D. Azarifar, M. A. Zolfigol, and B. Maleki, *Synthesis*, 2004, 1744.
15. D. Azarifar, M. A. Zolfigol, and B. Maleki, *Bull. Korean Chem. Soc.*, 2004, **25**, 23 (*Chem. Abstr.*, 2004, **141**, 1029).
16. D. Azarifar, H. Ghasemnejad-Bosra, and F. Ramazanian-Lehmali, *Mendeleev Commun.* 2005, 209.
17. D. Azarifar, H. Ghasemnejad-Bosra, M. A. Zolfigol, and M. Tajbaksh, *Heterocycles*, 2006, **68**, 175.
18. D. Azarifar and H. Ghasemnejad-Bosra, *Synthesis*, 2006, 1123.

19. D. Azarifar, H. Ghasemnejad-Bosra, and M. Tajbaksh, *J. Heterocycl. Chem.*, 2006, in press.
20. T. R. Beebe and J. W. Wolfe, *J. Org. Chem.*, 1970, **35**, 2056.
21. R. A. Eade and J. C. Earl, *J. Chem. Soc.*, 1946, 591.
22. W. Baker, W. D. Ollis, and V. D. Poole, *J. Chem. Soc.*, 1949, 307; W. Baker, W. D. Ollis, and V. D. Poole, *J. Chem. Soc.*, 1950, 1542.