

HETEROCYCLES, Vol. 68, No. 11, 2006, pp. 2253 - 2257. © The Japan Institute of Heterocyclic Chemistry
 Received, 28th August, 2006, Accepted, 2nd October, 2006, Published online, 3rd October, 2006. COM-06-10871

MICROWAVE-ASSISTED CYCLOADDITION REACTION OF AZIDES TO *N*-SUBSTITUTED 2-AZABICYCLO[2.2.1]HEPT-5-EN-3-ONES


Minoru Ishikura,^{a*} Miyako Hasunuma,^a Koji Yamada,^a and Reiko Yanada^b

^a Faculty of Pharmaceutical Sciences, Health Sciences University of Hokkaido, Ishikari-Tobetsu, Hokkaido 061-0293, Japan. E-mail: ishikura@hoku-iryu-u.ac.jp

^b Faculty of Pharmaceutical Sciences, Hiroshima International University, 5-1-1 Hirokoshingai, Kure, Hiroshima 737-0112, Japan

Abstract – The reaction of *N*-substituted 2-azabicyclo[2.2.1]hept-5-en-3-ones (ABH) (**1**) with azides (**2,4**) could be effected under microwave conditions, producing aziridine-fused ABH (**3**) and triazolines (**5,6**) in good yields.

Due to its attractive chemical diversity which is dependent on the increased ground state energy resulting from the great ring-strain in bicyclo[2.2.1]heptene system containing the amide group and the double bond, our interest was drawn to the synthetic exploration of 2-azabicyclo[2.2.1]hept-5-en-3-one (ABH) (**1**; R=H) as an important class of synthetic building block.¹ In connection with our project to prepare novel carbocyclic nucleosides,² we have previously disclosed the stereoselective addition of azides to **1a**.³ In contrast to the known facile reaction of norbornene with benzoyl azide,⁴ the attempted addition reaction of azides to **1a** under standard conditions was sluggish.⁵ Eventually, the high-pressure was found to promote the 1,3-dipolar cycloaddition reaction to give triazolines, but it required a longer reaction time and caution in treating under high-pressure conditions. We found that the addition reaction time could be significantly reduced when the reaction was performed under microwave



Scheme 1

conditions, and herein present our preliminary results.

Despite the recent remarkable progress in microwave-assisted protocols in synthetic organic chemistry,⁶ there have been few reports on microwave-assisted addition reaction of azides to double bond.⁷


Initially, a mixture of *N*-Boc ABH (**1a**) and electron-poor **2a** (2 equiv) were irradiated in a microwave reactor at 100 to 140 °C for 30 min with or without solvent,⁸ resulting in the formation of aziridine-fused ABH (**3a**) through nitrene-addition⁹ in low yields, along with substantial amounts of unchanged **1a** (Scheme 1 and Table 1). On the other hand, *N*-TBDMS ABH (**1b**) proved to be much more reactive toward **2a**. Microwave-irradiation of **1b** with **2a** at 140 °C for 30 min allowed the isolation of **3b** in 63% yield. Similarly, **1b** reacted with **2b** at 140 °C for 30 min to provide **3d** in 61% yield, as compared with the observed lower reactivity of **1a** toward **2b**.

Table 1 Microwave-assisted reaction of **1** with azides (**2**)

1	2	Conditions		Yield (%) ^a of 3
1a	2a	140 °C	30 min, in DMF	15 (3a) ^{2c}
1a	2a	100 °C	30 min, in toluene	25 (3a) ^{2c}
1a	2a	120 °C	30 min, without solvent	36 (3a) ^{2c}
1b	2a	140 °C	30 min, in DMF	47 (3b) ¹⁰
1b	2a	100 °C	30 min, in toluene	58 (3b) ¹⁰
1b	2a	140 °C	30 min, without solvent	63 (3b) ¹⁰
1a	2b	140 °C	60 min, in DMF	10 (3c) ³
1a	2b	140 °C	30 min, without solvent	17 (3c) ³
1b	2b	140 °C	60 min, in DMF	33 (3d) ¹⁰
1b	2b	140 °C	30 min, without solvent	61 (3d) ¹⁰

^a Yields based on **1**

The reaction of **1** could also be performed with azides (**4**). This reaction proceeded to completion within 10 min, yielding a pair of regioisomeric triazolines (**5** and **6**) in good yields by way of 1,3-dipolar



Scheme 2

cycloaddition reaction (Scheme 2 and Table 2). After the reaction was complete, **5** and **6** were separated by SiO₂ column chromatography. The structures of **5** and **6** were assigned based on a comparison of chemical shifts of the H-1 and H-4 protons in ¹H-NMR spectrum, as reported previously.³ The reactivity of **1a** toward **4** compares favorably with that of **1b**, and the reaction was best undertaken without solvent under microwave-conditions at 100 °C for 10 min.

Table 2 Microwave-assisted reaction of **1** with azides (**4**)

1	4	Conditions	Yield (%) ^a of 5	Yield (%) ^a of 6
1a	4a	140 °C, 10 min, in DMF	22 (5a) ³	39 (6a) ³
1a	4a	100 °C, 10 min, without solvent	23 (5a) ³	49 (6a) ³
1b	4a	140 °C, 10 min, in DMF	21 (5b) ¹⁰	45 (6b) ¹⁰
1b	4a	100 °C, 10 min, without solvent	24 (5b) ¹⁰	55 (6b) ¹⁰
1a	4b	140 °C, 30 min, in DMF	23 (5c) ¹⁰	36 (6c) ¹⁰
1a	4b	100 °C, 10 min, without solvent	28 (5c) ¹⁰	53 (6c) ¹⁰
1b	4b	140 °C, 30 min, in DMF	27 (5d) ¹⁰	36 (6d) ¹⁰
1b	4b	100 °C, 10 min, without solvent	31 (5d) ¹⁰	52 (6d) ¹⁰

^a Yields based on **1**

In summary, we have disclosed an alternative protocol for the reaction of ABH (**1**) with azides (**2**, **4**) using microwave-irradiation, providing aziridine-fused ABH (**3**) and triazolines (**5**, **6**), respectively.

ACKNOWLEDGEMENTS

This was supported in part by “Academic Frontier” Project for Private University: matching fund subsidy from Ministry of Education, Culture, Sports, Science and Technology, 2002-2006, in part by a Grant-in Aid for High Technology Research Program from Ministry of Education, Culture, Sports, Science and Technology of Japan, and in part by a Grant-in Aid for Scientific Research (No. 18590011) from the Ministry of Education, Culture, Sports, Science and Technology of Japan. The authors thank Kuraray Co., Ltd. for providing the ABH.

REFERENCES

1. M. K. Gurjar, S. Bera, R. R. Joshi, and R. A. Joshi, *Heterocycles*, 2003, **60**, 2293; M. Ishikura, M. Hasunuma, and M. Saijo, *Heterocycles*, 2004, **63**, 5; M. Ishikura, M. Hasunuma, and K. Yanada, *Heterocycles*, 2005, **65**, 2587; R. J. Brea, M. Amorin, L. Castedo, and J. R. Granja, *Angew. Chem. Int. Ed.*, 2005, **44**, 5710.
2. a) N. Katagiri, Y. Yamatoya, and M. Ishikura, *Tetrahedron Lett.*, 1999, **40**, 9069; b) M. Ishikura, A. Murakami, and N. Katagiri, *Org. Biomol. Chem.*, 2003, **1**, 452; c) M. Ishikura, K. Matsumoto, M. Hasunuma, and N. Katagiri, *Heterocycles*, 2003, **60**, 2737; d) M. Ishikura, K. Matsumoto, and A. Murakami, *Heterocycles*, 2004, **64**, 241.
3. M. Ishikura, S. Kudo, A. Hino, N. Ohnuki, and N. Katagiri, *Heterocycles*, 2000, **53**, 1499.

4. R. Huisgen, L. Möbius, G. Müller, H. Stangl, G. Szeimies, and J. M. Vernon, *Chem. Ber.*, 1965, **98**, 3992.
5. There is one example of cycloaddition reaction of phenyl azide to **1** (R=H) at 90 °C in CH₂Cl₂; J. R. Malpass, D. Belkacemi, G. A. Griffith, and M. D. Robertson, *ARKIVOC*, 2002, (vi), 164.
6. P. Lidstrom, J. Tierney, B. Wathey, and J. Westman, *Tetrahedron*, 2001, **57**, 9225; M. Larhed, C. Moberg, and A. Hallberg, *Acc. Chem. Res.*, 2002, **35**, 717; C. O. Kappa, *Angew. Chem. Int. Ed.*, 2004, **43**, 6250.
7. F. Louërât, K. Bougrin, A. Loupy, A. M. O. Retana, J. Pagalday, and F. Palacios, *Heterocycles*, 1998, **48**, 161.
8. Microwave irradiation was carried out using Green-Motif I (IMCR-25003) microwave reactor (IDX Corporation). To a glass tube were added **1b** (1 mmol) and **2a** (2 mmol), and the vessel was placed into the cavity of microwave reactor. Microwave irradiation at 300 W was performed, the temperature being ramped from rt to 140 °C. The mixture was let stand for 30 min. After cooling, the mixture was separated by medium pressure chromatography (SiO₂) with hexane-AcOEt.
9. *N*-Acyl azide as a good nitrene precursor: W. Lwowski, "Azides and Nitrenes—Reactivity and Utility," ed. by E. F. Scriven, Academic Press, London, 1984, pp. 205-246.
10. **3b**: ¹H-NMR (CDCl₃) δ: 0.21 (s, 3H), 0.32 (s, 3H), 0.89 (s, 9H), 1.52 (d, 1H, *J*=9.7 Hz), 1.86 (d, 1H, *J*=9.7 Hz), 2.45 (s, 3H), 2.84 (s, 1H), 3.28 (d, 1H, *J*=6.3 Hz), 3.43 (d, 1H, *J*=6.3 Hz), 3.94 (s, 1H), 7.35 (d, 2H, *J*=8.6 Hz), 7.80 (d, 2H, *J*=8.6 Hz). ¹³C-NMR (CDCl₃) δ: -5.3, -4.9, 18.9, 21.7, 26.3, 33.0, 40.0, 44.1, 47.2, 58.4, 127.9, 129.9, 134.8, 145.0, 181.9. **3d**: ¹H-NMR (CDCl₃) δ: 0.19 (s, 3H), 0.28 (s, 3H), 0.91 (s, 9H), 1.54 (d, 1H, *J*=9.7 Hz), 1.86 (d, 1H, *J*=9.7 Hz), 2.82 (s, 1H), 3.25 (s, 1H), 3.28 (s, 1H), 3.86 (s, 1H), 7.18-7.21 (m, 6H), 7.32-7.35 (m, 4H). ¹³C-NMR (CDCl₃) δ: -5.2, -4.8, 18.9, 26.4, 32.6, 37.6, 42.1, 47.3, 58.7, 120.3, 120.4, 125.5, 129.9, 182.6. **5b**: ¹H-NMR (CDCl₃) δ: 0.33 (s, 3H), 0.36 (s, 3H), 0.95 (s, 9H), 1.39 (d, 1H, *J*=10.9 Hz), 1.94 (d, 1H, *J*=10.9 Hz), 3.23 (s, 1H), 4.15 (s, 1H), 4.20 (d, 1H, *J*=9.2 Hz), 5.20 (d, 1H, *J*=9.2 Hz), 7.08 (t, 1H, *J*=7.4 Hz), 7.23-7.28 (m, 2H), 7.36-7.39 (m, 2H). ¹³C-NMR (CDCl₃) δ: -5.2, -4.8, 19.1, 26.3, 35.7, 51.4, 61.4, 62.3, 82.5, 114.0, 123.0, 129.8, 139.7, 181.0. **5c**: ¹H-NMR (CDCl₃) δ: 1.36 (td, 1H, *J*=1.7, 11.5 Hz), 1.43 (s, 9H), 1.85 (td, 1H, *J*=1.7, 10.9 Hz), 3.27 (s, 1H), 3.73 (d, 1H, *J*=9.2 Hz), 4.17 (s, 1H), 4.81 (d, 1H, *J*=14.9 Hz), 4.91 (d, 1H, *J*=14.9 Hz), 5.02 (d, 1H, *J*=9.2 Hz), 7.30-7.40 (m, 5H). ¹³C-NMR (CDCl₃) δ: 28.1, 31.8, 52.1, 53.5, 61.5, 62.2, 83.1, 83.6, 128.4, 128.5, 129.1, 135.9, 148.7, 172.0. **5d**: ¹H-NMR (CDCl₃) δ: -0.12 (s, 3H), 0.12 (s, 3H), 0.80 (s, 9H), 1.38 (d, 1H, *J*=10.3 Hz), 1.80 (d, 1H, *J*=10.3 Hz), 3.09 (s, 1H), 3.35 (s, 1H), 3.46 (d, 1H, *J*=9.7 Hz), 4.64 (d, 1H, *J*=14.3 Hz), 4.95 (d, 1H, *J*=14.3 Hz), 4.98 (d, 1H, *J*=9.7 Hz), 7.31-7.40 (m, 5H). ¹³C-NMR (CDCl₃) δ: -5.4, -5.5,

18.8, 26.2, 35.9, 51.7, 53.8, 62.3, 64.5, 83.2, 128.5, 128.7, 129.1, 135.9, 181.5. **6b**: $^1\text{H-NMR}$ (CDCl_3) δ : 0.32 (s, 3H), 0.34 (s, 3H), 0.96 (s, 9), 1.39 (d, 1H, $J=10.9$ Hz), 1.92 (d, 1H, $J=10.9$ Hz), 3.13 (s, 1H), 4.35 (s, 1H), 4.38 (d, 1H, $J=9.2$ Hz), 5.06 (d, 1H, $J=9.2$ Hz), 7.06-77.09 (m, 1H), 7.35-7.39 (m, 4H). $^{13}\text{C-NMR}$ (CDCl_3) δ : -5.1, -5.08, 18.9, 26.4, 35.8, 51.1, 57.0, 62.6, 88.5, 114.3, 123.1, 129.7, 139.3, 180.8. **6c**: $^1\text{H-NMR}$ (CDCl_3) δ : 1.36 (d, 1H, $J=10.9$ Hz), 1.50 (s, 9H), 1.90 (d, 1H, $J=10.9$ Hz), 2.74 (s, 1H), 3.77 (d, 1H, $J=9.2$ Hz), 4.72 (d, 1H, $J=14.9$ Hz), 4.85 (s, 1H), 4.96 (d, 1H, $J=14.9$ Hz), 5.01 (d, 1H, $J=9.2$ Hz), 7.25-7.37 (m, 5H). $^{13}\text{C-NMR}$ (CDCl_3) δ : 28.1, 32.2, 52.1, 53.0, 58.8, 62.1, 83.9, 86.7, 128.3, 128.5, 129.1, 135.4, 148.6, 172.1. **6d**: $^1\text{H-NMR}$ (CDCl_3) δ : 0.23 (s, 3H), 0.25 (s, 3H), 0.90 (s, 9H), 1.38 (d, 1H, $J=10.3$ Hz), 1.86 (d, 1H, $J=10.3$ Hz), 2.60 (s, 1H), 3.74 (d, 1H, $J=9.3$ Hz), 4.21 (s, 1H), 4.70 (d, 1H, $J=14.3$ Hz), 4.83 (d, 1H, $J=9.3$ Hz), 4.95 (d, 1H, $J=14.3$ Hz), 7.27-7.37 (m, 5H). $^{13}\text{C-NMR}$ (CDCl_3) δ : -5.2, -5.3, 18.9, 26.3, 36.3, 51.6, 52.7, 58.5, 62.7, 88.8, 128.3, 129.0, 135.8, 181.3.