

SYNTHESIS OF (\pm)-3-(1',1'-DIMETHYLALLYL)DECURSINOL AND 3-(1',1'-DIMETHYLALLYL)-XANTHYLETIN

Rosario Hernández Galán, Guillermo M. Massanet, Enrique Pando, Francisco Rodríguez Luis, and Javier Salvá


Departamento de Química Orgánica, Facultad de Ciencias, Universidad de Cádiz, Apartado 40, Puerto Real (Cádiz), Spain


Abstract -- (\pm)-3-(1',1'-Dimethylallyl)decursinol (1) and 3-(1',1'-dimethylallyl)-xanthyletin (2) were synthesized from umbelliferone (3) in each 6.9% overall yield.

As a part of our continuing efforts toward synthesis of 3-(1',1'-dimethylallyl)coumarin derivatives¹ we describe herein the synthesis of (\pm)-3-(1',1'-dimethylallyl)decursinol (1) and 3-(1',1'-dimethylallyl)xanthyletin (2), the latter is a natural product previously isolated from *Boenninghausenia albiflora*² (Rutaceae). Because of their biological activity^{3,4} and because their small quantities from natural sources, we undertook their synthesis in order to obtain sufficient quantities for study their further pharmacological effects.

1 and 2 were obtained each in 6.9% overall yield from umbelliferone (3). The key step involves sigmatropic rearrangements in dimethylaniline (DMA), of 6-(3',3'-dimethylallyl)-7-(3",3"-dimethylallyloxy)coumarin (4) to gravelliferone (5, 26%) and 6-(3',3'-dimethylallyl)-8,8,9-trimethyl-8,9-dihydroangelicin (6, 57%), as reported previously¹. Simultaneously this reaction has been described⁵ using diethylaniline as solvent but the yields in the C-3 prenylated compound (5, 20%), and in 6 (8%) were not improved.

Treatment of 5 with MCPBA in CHCl_3 led to (\pm)-3-(1',1'-dimethylallyl)decursinol (1, 64%). When 1 was treated with p-toluensulphonylchloride (TsCl) afforded the p-toluensulphonate ester 7 (49.8%) accompanied by the starting product 1 (48.1%). It has been reported⁶ that the best way to prepare pyranocoumarins by dehydration of the corresponding alcohols is by refluxing their p-toluensulphonate ester in collidine. So refluxing 7 in collidine the elimination product 3-(1',1'-dimethylallyl)xanthyletin (2) was obtained in quantitative yield.





EXPERIMENTAL

Melting points were determined in a Kofler block Reichert-Jung apparatus and are uncorrected. Ir spectra were recorded in a Perkin-Elmer 257, values being given in cm^{-1} . Uv were registered in a Shimadzu MPS-2000, dissolved in MeOH, and are given in nm. The nmr spectra were recorded with a Hitachi-Perkin-Elmer H-24B 60 MHz instrument or in a Bruker Spectrospin 200 MHz using TMS as internal reference; chemical shifts are given in δ and coupling constants in Hz. Mass spectra were measured with a VG 12250. Thin Layer Chromatography was done on MN Alugram SIL G/UV 254 plates, 0.25 mm thick. Merck silica gel (0.06-0.2 mm) was used in column chromatography and elution was carried out with mixtures of hexane and ethyl acetate.

Gravelliciferrone (5) was obtained from umbelliferone (3), through 8-iodoumbelliferone, 7-(1',1'-dimethylpropyloxy)-8-iodocoumarin, 7-(1',1'-dimethylallyloxy)-8-iodocoumarin, demethyl-suberosin and 6-(3',3'-dimethylallyl)-7-(3',3'-dimethylallyloxy)coumarin, in a 10.3% overall yield, as reported previously¹.

(±)-3-(1',1'-Dimethylallyl)decursinol (1): Gravelliciferrone (5) (36 mg) was dissolved in 5 ml of CHCl_3 previously acidified (2 drops HCl conc. in 10 ml CHCl_3) and the solution was cooled at 0°C. MCPBA (25 mg) dissolved in 5 ml of the acidified CHCl_3 was added. After agitating for 6 h the reaction mixture was monitored by TLC, showing that the starting material had disappeared. After evaporating the solvent the reaction mixture was chromatographed on a silica gel column separating 24 mg (64%) of 1 (overall yield 6.9%), mp 181-183°C (CHCl_3). $(\alpha)_D^{25}$ 0° (c, 0.12; CHCl_3), uv λ_{max} 330, 258, ir (KBr) 3340, 1700, 1630, 1570, 1500, 1275, H-nmr (200 MHz, CDCl_3) 7.44 (s, H-4); 7.13 (s, H-5); 6.72 (s, H-8); 6.13 (dd, J=17 and 11, H-2'); 5.06 (d, J=11, H-3'cis); 5.04 (d, J=17, H-3'trans); 3.83 (t, J=6, H-3"); 3.08 (dd, J=17 and 6, H-4"); 2.79 (dd, J=17 and 6, H-4"); 1.44 (s, $\text{Me}_2\text{-C-1'}$); 1.36 (s, Me-C-2"); 1.33 (s, Me-C-2"), ms m/z 314 (7); 299 (7); 243 (8); 69 (86); 57 (100).

(±)-3-(1',1'-Dimethylallyl)decursinol p-Toluensulphonate Ester (7): To 1 (27 mg) dissolved in 3 ml of dry pyridine, 20 mg of p-toluensulphonyl chloride in 3 ml of pyridine was added. The mixture was stirred at room temperature for 7 days after which it was taken to dryness and separation over silica gel column afforded: 7 (16 mg, 49.8%) and 1 (13 mg, 48.1%) 7, oil, $(\alpha)_D^{25}$ 0° (c, 0.12; CHCl₃), uv λ_{\max} 326, ir (KBr) 1730, 1620, 1570, 1500, 1370, 1285, 1180, H-nmr (200 MHz, CDCl₃) 7.57 (d, J=8, H-2'''' and H-6'''); 7.41 (s, H-4); 7.31 (d, J=8, H-3'''' and H-5'''); 7.03 (s, H-5); 6.67 (s, H-8); 6.12 (dd, J=17 and 11, H-2'); 5.05 (d, J=11, H-3'cis); 5.03 (d, J=17, H-3'trans); 4.63 (t, J=6, H-3''); 3.13 (dd, J=18 and 6, H-4''); 2.99 (dd, J=18 and 6, H-4''); 2.43 (s, Me-C-4'''); 1.43 (s, Me₂-C-2'); 1.24 (s, Me-C-2''); 1.20 (s, Me-C-2''), ms m/z 468 (9.5); 453 (4.5); 296 (10); 281 (46); 69 (100).

3-(1',1'-Dimethylallyl)xanthyletin (2): 7 (16 mg) was dissolved in collidine and refluxed (200°C) for 24 h. The reaction mixture was monitored by TLC and showed total absence of starting material. Et₂O and 5% HCl were added. The organic layer was neutralized with saturated NaHCO₃ solution, washed with brain and dried over anhydrous Na₂SO₄. Evaporation of the solvent furnished 10 mg (quantitative yield) of 3-(1',1'-dimethylallyl)xanthyletin (2) (overall yield 6.9%), mp, 93-95°C (hexane/AcOEt) (lit² 98-100°C, petrol/acetone), uv λ_{\max} 347, 304, 267, ir (KBr) 1710, 1620, 1600, 1500, 1270, H-nmr (200 MHz, CDCl₃) 7.45 (s, H-4); 7.00 (s, H-5); 6.66 (s, H-8); 6.33 (d, J=10, H-4''); 6.13 (dd, J=17 and 11, H-2'); 5.64 (d, J=10, H-3''); 5.06 (d, J=11, H-3'cis); 5.04 (d, J=17, H-3'trans); 1.44 (s, Me₂-C-1' and Me₂-C-2''), ms m/z 296 (22), 281 (100).

ACKNOWLEDGEMENT

Thanks to Organic Chemistry Department (University of Málaga) where H-nmr spectra of compounds 1, 2 and 7 were recorded.

Financial support from Comision Asesora de Investigación Científica y Técnica (CAICYT) is gratefully acknowledged (n2 PR84-0247).

REFERENCES

1. G. M. Massanet, E. Pando, F. Rodríguez Luis, and J. Salvá, *Heterocycles*, **26**(6) 1541 (1987).
2. M. N. S. Nayar, M. K. Bhan, and V. George, *Phytochemistry*, **12**, 2073 (1973).
3. M. Gosálvez, R. G. Cañero, and M. Blanco, *Europ. J. Cancer*, **12**, 1003 (1976).
4. A. G. González, V. Darias, G. Alonso, J. N. Boada, and F. Rodríguez Luis, *Planta Medica*, **31**, 351 (1977).
5. N. Cairns, L. M. Harwood, and D. P. Astles, *J. Chem. Soc. Chem. Commun.*, 400 (1987).
6. S. N. Shanbag, C. K. Mesta, M. L. Maheshwari, S. K. Paknikar, and S. C. Bhattacharyya, *Tetrahedron*, **20**, 2605 (1964).
7. J. Reisch, K. Szendrei, E. Minker, and I. Novak, *Experientia*, **24**, 992 (1968).

Received, 24th November, 1987