

THE REACTIVITY OF 2-FORMYLTHIOPHENE, 1-METHYL-2-FORMYLPYRROLE,  
 BENZALDEHYDE AND 2-FORMYLFURANE N,N-DIMETHYLHYDRAZONES WITH  
 DIMETHYL ACETYLENEDICARBOXYLATE

M. Eugenia Gonzalez, Pedro Sancho, Concepción Soriano,  
 Rafael Ballesteros, Belén Abarca, and José Sepulveda.

Dep. Química Orgánica, Facultad Farmacia, 46010-Valencia, Spain


Abstract- N,N-Dimethylhydrazones derived from 2-formylthiophene, benzaldehyde, and 1-methyl-2-formylpyrrole react with dimethyl acetylenedicarboxylate (DMAD), giving the corresponding nitriles. The reaction with furfural N,N-dimethylhydrazone takes place via the intermediacy of a Diels-Alder adduct with the furan ring.

It has been shown recently<sup>1</sup> that although 1-aza-1,3-dienes are reluctant to participate in Diels-Alder reactions, these systems with an appropriate substitution can react with electron-poor dienophiles affording the corresponding cycloadducts. Looking at this new possibility of synthesis of heterocycles and as an extension of our studies on the reactivity, as 1,3-dienes, of five membered heterocyclic compounds bearing olefinic substituents,<sup>2</sup> we have prepared the N,N-dimethylhydrazones 1, 2, 3, and 4 and we have studied their reactivity with DMAD.

The reaction of the N,N-dimethylhydrazone 1 with DMAD was carried out in chloroform or toluene at reflux temperature affording 2-thienylnitrile 5 and another compound identified as the aminodiester 6 by spectroscopic analysis.<sup>3</sup> No Diels-Alder cycloaddition compounds were found in the crude mixture.

The same reaction with benzaldehyde N,N-dimethylhydrazone 2 gave the corresponding nitrile.


When the hydrazone **3** in chloroform was mixed with DMAD at room temperature, the Michael addition reaction took place and the adducts **7a** and **7b**, from attack at  $C_2$ , could be isolated with good yields. The same reaction with two moles of DMAD at reflux temperature afforded a mixture of the E/Z isomers **8a** and **8b**, that could be separated chromatographically. Here again the aminoester **6** was isolated from the reaction mixture.



Hydrazones have the well-known capacity to react with electrophilic reagents at either of the three nucleophilic centers, the amino nitrogen, the imino nitrogen and the imino carbon.<sup>4</sup> However when  $\alpha,\beta$ -unsaturated *N,N*-dimethylhydrazones react with electron-poor dienophiles giving cycloaddition compounds<sup>4</sup>, the mechanism of the reaction involves the initial attack of the  $C_2$  to the electrophile, because of the reversed polarity of these 1-amino-1-aza-1,3-diene systems.

Our experiments with the *N,N*-dimethylhydrazones **1**, **2**, and **3**, have shown that these compounds react with DMAD by the amino nitrogen and afford nitriles as a

result of the elimination of the aminodiester 6 through a six membered intermediate 9.



9

Although many methods are described for the synthesis of nitriles from aldehyde hydrazones as starting materials,<sup>15</sup> the acetylenic esters were never used as reagents for this conversion, so the present work provides a new route for the synthesis of these compounds.

In complete contrast with the hydrazones 1, 2, and 3, the 2-formylfuran *N,N*-dimethylhydrazone 4 reacts with DMAD in chloroform at reflux temperature to give a major compound 10 through a Diels-Alder cycloaddition to the furan ring in a similar way as it has been reported by Potts<sup>16</sup> in the reaction with maleic anhydride. Small amounts of 2-furylcarbonitrile 11 and the aminodiester 6 were isolated as a result of a competing reaction of the dimethylhydrazone group. Nevertheless, it was not possible to detect any amount of the corresponding nitrile 12.

Reactions with another dienophiles are in progress.

#### EXPERIMENTAL

<sup>1</sup>H and <sup>13</sup>C nmr spectra were measured for ca. 40% solutions in CDCl<sub>3</sub> using a Bruker WP-80. Ir were determined in CCl<sub>4</sub> with a Perkin-Elmer 843 spectrometer. Melting points are uncorrected and were determined with a Kofler hot-stage apparatus. Combustion analyses were performed at the Instituto de Quimica Bio-Organica of the CSIC (Barcelona).

#### Reaction of *N,N*-Dimethylhydrazones with DMAD. General Procedure.-

DMAD (0.005 mol) in chloroform or toluene (10 ml) was added at room temperature to a solution of the corresponding *N,N*-dimethylhydrazone (0.005 mol) in chloroform or toluene (10 ml). Then the mixture was heated at reflux temperature for

30-40 h until disappearance of the starting material. Evaporation of the solvent gave a crude product which was purified by column chromatography.

**2-Formylthiophene N,N-dimethylhydrazone:** Elution of the crude product (1.4 g) with hexane-ethyl acetate (5:1) gave 2-thienylcarbonitrile **5** (0.45 g, 82%) and diethyl 2-dimethylaminomaleate **6** (0.7 g).

**Benzaldehyde N,N-dimethylhydrazone:** Column chromatography of the crude product (1.3 g) with hexane-ethyl acetate (5:1) gave benzonitrile (0.33 g, 66%) and the adduct **6** (0.6g).

**1-Methyl-2-formylpyrrole N,N-dimethylhydrazone:**

a) **At room temperature.** The reaction of DMAD (0.72 g, 0.005 mol) with the hydrazone **3** (0.75 g, 0.005 mol) in chloroform (20 ml) took place at room temperature when the mixture was stirred for 3 h. Evaporation of the solvent *in vacuo* and column chromatography with hexane-ethyl acetate (4:1) gave the adduct **7b** (0.45 g, 30%) as an oil. Anal. Calc for  $C_{14}H_{19}N_3O_4$ : C, 57.33; H, 6.48; N, 14.33. Found: C, 57.29; H, 6.57; N, 14.27. Ir  $1720\text{ cm}^{-1}$ .  $^1\text{H-nmr}$  7.1 (s, 1H), 6.75 (s, 1H), 6.0 (d, J=3Hz, 1H), 5.9 (d, J=3Hz, 1H), 3.75 (s, 3H), 3.6 (s, 6H), 2.85 (s, 6H). Further elution gave the isomer **7a** (0.42 g, 27.5%), mp 112-113 °C (yellow needles, recrystallized from hexane); Found: C, 57.52; H, 6.78; N, 14.38. Ir 1715,  $1702\text{ cm}^{-1}$ .  $^1\text{H-nmr}$  6.95 (s, 1H), 6.2 (d, J=4Hz, 1H), 6.0 (d, J=4Hz, 1H), 5.75 (s, 1H), 3.8 (s, 3H), 3.75 (s, 3H), 3.65 (s, 3H), 2.85 (s, 6H, NMe<sub>2</sub>).

b) **At reflux temperature.** The reaction of DMAD (1.5 g, 0.1mol) with the hydrazone **3** in chloroform (20 ml) at reflux temperature and working up as above gave a syrup which was chromatographed using hexane-ethyl acetate (9:1). The first compound eluted was the nitrile **8b** (0.28 g, 21.5%) as an oil. Anal. Calc for  $C_{12}H_{12}N_2O_4$  C, 58.06 H, 4.83; N, 11.29. Found: C, 58.19; H, 4.81; N, 11.05. Ir 2220,  $1720\text{ cm}^{-1}$ .  $^1\text{H-nmr}$  7.2 (s, 1H), 6.75 (d, J=4Hz, 1H), 6.1 (d, J=4Hz, 1H) 3.85 (s, 3H), 3.7 (s, 3H), 3.6 (s, 3H). Further elution gave the isomer **8a** (0.2 g, 15.4%), mp 102-103 °C (colorless needles, recrystallized from hexane-dichloromethane); Found: C, 57.9; H, 4.7; N, 11.08. Ir 2220,  $1730\text{ cm}^{-1}$ .  $^1\text{H-nmr}$  6.7 (d, J=4Hz, 1H), 6.3 (d, J=4Hz, 1H), 6.1 (s, 1H), 3.8 (s, 3H), 3.75 (s, 6H).

**Furfural N,N-dimethylhydrazone:** Chromatography of the crude product obtained after evaporation of the solvent (1.4 g), using hexane-ethyl acetate (5:1) as an eluant gave 2-furylcarbonitrile **11** (0.034 g.). The second compound eluted was the adduct **10** (0.467 g, 35%), mp 60-61 °C (yellow needles recrystallized

from hexane). Anal. Calc for  $C_{15}H_{12}N_2O_5$ : C, 55.86; H, 5.66; N, 9.81. Found: C, 55.71; H, 5.71; N, 10.0. Ir  $1733\text{ cm}^{-1}$ .  $^1\text{H-nmr}$  11-10.5 (bs, 1H), 7.75 (d,  $J=9\text{Hz}$ , 1H), 6.75 (d,  $J=9\text{Hz}$ , 1H), 6.7 (s, 1H), 3.7 (s, 6H), 2.8 (s, 6H).  $^{13}\text{C-nmr}$  169.2 (s), 168.7 (s), 160.7 (s), 132.4 (d), 131.9 (s), 127.2 (d), 126.2 (s), 119.2 (d), 108.8 (s), 52.6 (q), 52.0(q), 42.3 (q). Further elution gave the starting material **4** (0.26 g) and then the amino compound **6** (0.53g).

#### ACKNOWLEDGEMENTS

We wish to thank the CAICYT for financial support (project PB-85-0205)

#### REFERENCES

1. B. Serckx-Poncin, A. M. Hesbain-Frisque, and L. Ghosez, Tetrahedron Lett., 1982, **23**, 3261.
2. R. A. Jones, M. T. P. Marriott, W. P. Rosenthal, and J. Sepulveda, J. Org. Chem., 1980, **45**, 4515; R. A. Jones, P. Martinez, T. Aznar, and J. Sepulveda, Tetrahedron, 1984, **40**, 4837; R. A. Jones, T. Aznar, and J. Sepulveda, J. Chem. Soc. Perkin Trans I, 1984, 2541; B. Abarca, R. Ballesteros, E. Enriquez and G. Jones, Tetrahedron, 1985, **41**, 2435; Ibid., 1987, **43**, 269; M. Medico, and J. Sepulveda, Tetrahedron, 1986, **42**, 6683.
3. R. Huisgen, K. Herbig, A. Siegl, and H. Huber, Chem. Ber., 1966, **99**, 2526.
4. P. Bouchet, J. Elguero, and R. Jaquier, Bull. Soc. Chim. France, 1967, 4716; H. Suschitzky, R. E. Walrond, and R. Hull, J. Chem. Soc. Perkin I, 1977, 47; B. B. Snider, R. S. E. Conn, and S. Sealson, J. Org. Chem., 1979, **44**, 218; G. Le Fevre and J. Hamelin, Tetrahedron, 1980, **36**, 887; M. N. Gudi, J. C. Hiriyakkanavar, and M. V. George, Indian J. Chem., 1971, **9**, 743; M. K. Saxena, M. N. Gudi, and M. V. George, Tetrahedron, 1973, **29**, 101; R. Baumes, R. Jaquier, and G. Tarrago, Bull. Soc. Chim. France, 1974, **11**, 2547; S. F. Nelsen, J. Org. Chem., 1969, **34**, 2248.
5. R. F. Smith, J. A. Albrigh, and A. M. Waring, J. Org. Chem., 1966, **31**, 4100; R. F. Smith and K. J. Mc-Grath, J. Org. Chem., 1976, **41**, 395; R. W. Binkley, Tetrahedron Lett., 1970, 2085; F. Yoneda and T. Nagamatsu, Bull. Chem. Soc. Japan, 1975, **48**, 1484; A. R. Katritzky, and P. Molina, J. Chem. Soc. Perkin Trans I, 1979, 1957; T. Cuvigny, J. F. Le Borgne, M. Larcheveque, and H. Normant, Syntheses, 1976, 237.
6. K.T.Potts and E. B. Walsh, J. Org. Chem., 1984, **49**, 4099.

Received, 7th January, 1988