

A LEWIS ACID MEDIATED INTRAMOLECULAR [2+2] CYCLOADDITION OF 3-(9-METHYLUNDECA-7,9-DIENYL)-9-METHYLENE-2,8-DIOXABICYCLO-[4.3.0]NONANE-5,7-DIONE

Kei Takeda,* Akihiko Shimotani, and Eiichi Yoshii*


Faculty of Pharmaceutical Sciences, Toyama Medical and Pharmaceutical University
Sugitani 2630, Toyama 930-01, Japan

Kentaro Yamaguchi


School of Pharmaceutical Sciences, Showa University, 1-5-8 Hatanodai, Shinagawa-ku,
Tokyo 142, Japan

Abstract - Treatment of 3-(9-methylundeca-7,9-dienyl)-9-methylene-2,8-dioxabicyclo[4.3.0]nonane-5,7-dione (**3**) with SnCl_4 in CH_2Cl_2 at -80°C produced intramolecular [2+2] cycloadducts (**6a-d**) rather than intramolecular Diels-Alder adducts.


In a project directed toward the synthesis of the antibiotic okilactomycin (**1**),¹ we have been interested in utilizing an intramolecular Diels-Alder technique for a simultaneous construction of the macro-ring as well as the spiro lactone system having the correct stereochemistries. Conceptually, the same strategy has previously been employed in our total synthesis of chlorothricolide.² This report describes a model cycloaddition study performed with compound (**3**), resulting in formation of the intramolecular [2+2] cycloadducts (**6a-d**), rather than the desired Diels-Alder adduct (**2**).



The substrate (**3**) was prepared as outlined in Scheme 1, in which formation of the 2,8-dioxabicyclononane-dione skeleton was effected by a two-step acid-catalyzed cyclization of γ,δ -unsaturated α -dioxolanylcarbonyl- β -ketoester (**5**) which was obtained according to the Gelin's procedure.⁴


Scheme 1³

Since **3** proved to be thermally unstable, readily undergoing polymerization, the cycloaddition was carried out at low temperature (-80°C) in CH₂Cl₂ (8 mM) in the presence of SnCl₄ (1 equiv) for 90 min. Under the conditions, there were produced four diastereomeric [2+2] cycloadducts (**6a-d**) in a ratio of *ca.* 3.5:1:1:1 (*ca.* 30% yield⁷) (Scheme 2).

Scheme 2³

The unprecedented cyclobutane structures fused to a 13-membered ring were confirmed by a single crystal X-ray analysis⁸ on **6a** and **6b**. The structures of **6c** and **6d**, which unfortunately could not be obtained as suitable crystals for X-ray analysis, were determined by comparison of their ¹H nmr spectra with those of **6a** and **6b**. The unexpected formation of [2+2] cycloadducts can be interpreted, as analyzed by molecular models, by an

easy access of the C(10') olefinic center and the exo-methylene group of the butenolide without severe distortion of the molecule leading to a stabilized dipolar intermediate (7). On the other hand, it seems to be relatively difficult to align the diene and the dienophile in the proper geometry for a concerted [4+2] cycloaddition. There is no reported example⁹ where cyclobutane ring is formed in intramolecular [2+2] cycloaddition accompanying by formation of macrocyclic ring. Further studies are underway to define the scope and limitations of this type of intramolecular [2+2] cycloaddition.



REFERENCES

1. H. Imai, K. Suzuki, M. Morioka, Y. Numasaki, S. Kadota, K. Nagai, T. Sato, M. Iwanami, and T. Saito, *J. Antibiot.*, 1987, **40**, 1475; H. Imai, H. Kaniwa, T. Tokunaga, S. Fujita, T. Furuya, H. Matsumoto, and M. Shimizu, *ibid.*, 1987, **40**, 1483.
2. K. Takeda, Y. Igarashi, K. Okazaki, and E. Yoshii, *J. Org. Chem.*, 1990, **55**, 3431.
3. All compounds are racemic.
4. P. Pollet and S. Gelin, *Synthesis*, 1978, 142; B. Chantegrel, A-I. Nadi, and S. Gelin, *ibid.*, 1983, 948; S. Gelin and R. Gelin, *Bull. Soc. Chim. Fr.*, 1968, 288.
5. P. J. Kocienski, B. Lythgoe, and I. Waterhouse, *J. Chem. Soc., Perkin Trans. I*, 1980, 1045.
6. A. Tanaka and K. Yamashita, *Agric. Biol. Chem.*, 1980, **44**, 199.
7. The main side reactions were polymerization of compound (3).
8. **6a** : FW = 330.42, Monoclinic P2₁/a, a = 15.870(1), b = 8.061(1), c = 14.879(1)Å, β = 112.46(1)°, V = 1759.1(2)Å³, Z = 4, D_c = 1.248 Mgm⁻³, R = 0.048 for 2608 reflections. **6b** : FW = 330.42, Monoclinic A2/a, a = 28.875(4), b = 6.820(2), c = 19.572(2)Å, β = 108.56(1)°, V = 3653(1)Å³, Z = 8, D_c = 1.201 Mgm⁻³, R = 0.068 for 2724 reflections Detailed crystallographic analyses will be published elsewhere.
9. For non-photochemical [2+2] cycloaddition; see P. G. Gassman and A. C. Lottes, *Tetrahedron Lett.*, 1992, **33**, 157, and references cited therein.

Received, 8th July, 1992