

CONFORMATIONAL PREFERENCE IN BENZYLOXY- AND SILOXY-SUBSTITUTED THIANES, THIANE 1-OXIDES, AND THIANE 1,1-DIOXIDES

Yoshimitsu Nagao^{a*}, Michimasa Goto^b, Kiyoshi Kida^a, and Motoo Shiro^c


^aFaculty of Pharmaceutical Sciences, The University of Tokushima, Sho-machi, Tokushima 770, Japan

^bChemical Instrument Center, Nagoya University, Chikusa, Nagoya 464-01, Japan

^cRigaku Corporation, 3-9-12, Matsubara-cho, Akishima, Tokyo 196, Japan

Abstract- Conformer ratios in 4-benzyloxy- and 4-siloxy-substituted thianes, *cis*- and *trans*-thiane 1-oxides, thiane 1,1-dioxides, and dihydrothiines were revealed on the basis of their low temperature ¹H nmr analyses. Extreme benzyloxy- and siloxy-axial (or -pseudoaxial) conformer preferences in *trans*-thiane 1-oxides, thiane 1,1-dioxides, and dihydrothiines were clearly demonstrated.

Previously, we reported axial (*ax*) conformer preferences (>70% at 25 °C) of alkoxy and siloxy groups in the chair-type cyclohexanones (**1**) and (**2**), glutaric anhydride (**3**; X = O), and glutarimide (**3**; X = NCH₂Ph) not



only in their CDCl_3 solution but also in the crystalline state.¹ These prochiral compounds (1) and (3) should be attractive to us from the viewpoint of a new design for the asymmetric induction.²

Significant attention was earlier directed toward the conformational analyses of substituted thianes and their 1-oxides and 1,1-dioxides. Martin and Ubel reported a conformational preference for the forms having sulfoxide oxygen-*ax* over those with sulfoxide oxygen-equatorial (*eq*) in the *cis*- and *trans*-4-chloro-, 4-hydroxy-, or 4-*p*-toluenesulfonyloxythiane 1-oxides.³ Ōki and Nogami disclosed interesting evidence of a conformational preference for the forms having Br-*eq* in the 4-bromothiane and its *cis*-1-oxide and for the forms having Br-*ax* in the *trans*-4-bromothiane 1-oxide and its 1,1-dioxide.⁴ These results mentioned above prompted us to investigate the systematic conformational analyses of the titled compounds (4-7) and dihydrothiines (8).




Figure 1. Conformational equilibrium modes in 4-RO-substituted thianes, thiane 1-oxides, thiane 1,1-dioxides, and dihydrothiines

The low-temperature ^1H nmr spectra of 4-RO-substituted thianes (4a-d),⁵ *cis*-thiane 1-oxides (5a-c),⁵ *trans*-thiane 1-oxides (6a-c),⁵ thiane 1,1-dioxides (7a-c),⁵ and 4-RO-dihydrothiine (8c)⁶ gave rise to two sets of 4-H

or 2-H peaks which correspond to the RO-*eq* and RO-*ax* conformers based on the result from their ring interconversion as shown in Figure 1 and Table I.

Table I. Conformer ratios in RO-substituted thianes (**4a-d**), *cis*-thiane 1-oxides (**5a-c**), *trans*-thiane 1-oxides (**6a-c**), thiane 1,1-dioxides (**7a-c**), and dihydrothiure (**8c**) based on ^1H nmr analysis (400 MHz, CD_2Cl_2).





Compound	Conformer Ratio ^{a)}	δ (ppm) of 4-H		T_c ^{b)} /°C
	RO- <i>eq</i> : RO- <i>ax</i>	RO- <i>eq</i>	RO- <i>ax</i>	
4a	76 : 24	3.23	3.77	-60
4b	69 : 31 ^{c)}	3.46	4.09	-55
4c	47 : 53	3.42	4.03	-50
4d	89 : 11	3.37	4.02	-60
5a	55 : 45	3.45	3.71	-40
5b	34 : 66	3.65	3.97	-40
5c	33 : 67	3.60	3.89	-40
6a	11 : 89	3.40	3.81	-50
6b	4 : 96	3.76	4.10	-50
6c	2 : 98	3.70	4.07	-50
7a	5 : 95	3.59	3.84	-60
7b	2 : 98	3.82	4.10	-70
7c	3 : 97	3.74	4.05	-60
8c	2 : 98 ^{d)}	3.16 ^{e)}	3.57 ^{e)}	-75

a) Unless otherwise, determined at -80 °C. b) Coalescence temp. of 4-H peaks. c) Determined at -90 °C d) Determined at -95 °C. e) δ (ppm) Value of 2-H_{ax}.

^1H nmr (200 MHz, CDCl_3 , room temperature) bandwidths (W) of 4-H peaks of **4a-d** exhibited fairly large values (**4a**: 27.6 Hz, **4b**: 25.1 Hz, **4c**: 24.0 Hz, and **4d**: 29.0 Hz), which suggested their RO-*eq* conformer preferences.¹ Then, we examined their 400 MHz ^1H nmr analyses at -80 °C and could reveal the details of their conformational ratios (*eq* vs *ax*) in CD_2Cl_2 solution as shown in Table I. Interestingly, the order of RO-*eq* conformer preferences among siloxy derivatives (**4b-d**) is shown to be **4d**>**4b**>**4c**. This order must be contrary to that of the bulky size of siloxy groups. Eventually, the order (**4d**>**4a**>**4b**>**4c** and **5a**>**5b**>**5c**) seemed to be in proportion to the basicity (electron density) of the oxygen atom of 4-RO groups. Hence, basicity order of the oxygen atom of the related benzyloxy and siloxy derivatives⁷ was tentatively determined on the basis of up-field shift ($\Delta\delta$ ppm) of ^{119}Sn -peak of Me_2SnCl_2 in the presence of the corresponding cyclohexanol ether as shown in

Table 2. Interestingly, the basicity order ($\text{Me}_3\text{Si-O} > \text{PhCH}_2\text{-O} > t\text{-Bu}(\text{Me})_2\text{Si-O} > t\text{-Bu}(\text{Ph})_2\text{Si-O}$) obtained by our method is perfectly consistent with the order of the RO-*eq* conformer preferences in the compounds (4) and (5).

Table 2. Up-field shift ($\Delta\delta$ ppm) of ^{119}Sn -peak (149 MHz, ^{119}Sn nmr) of Me_2SnCl_2 in the presence of ethers in CDCl_3 at -50°C .

Run ^{a)}	Conditions ^{a)}	Chemical Shift of ^{119}Sn (δ ppm) ^{b)}	$\Delta\delta$ ppm ^{c)}
1)	Me_2SnCl_2 (0.3 mmol)/ CDCl_3 (2.5 ml)	143.42	—
2)	Me_2SnCl_2 + 	120.09	-23.33
3)	Me_2SnCl_2 + 	131.51	-11.91
4)	Me_2SnCl_2 + 	138.45	-4.97
5)	Me_2SnCl_2 + 	142.92	-0.50

a) In Runs 2-5, a mixture of Me_2SnCl_2 (0.3 mmol) and each ether (0.3 mmol) was determined in CDCl_3 (2.5 ml). b) Each chemical shift is indicated in the ppm value relative to the $\text{Me}_4^{119}\text{Sn}$ signal.

c) $\Delta\delta$ ppm = δ ppm ($\text{Me}_2^{119}\text{SnCl}_2$ + ether) - δ ppm ($\text{Me}_2^{119}\text{SnCl}_2$)

In the cases of *cis*-4-RO-thiane 1-oxides (**5a-c**), their RO-*ax* conformer preferences proved to be 45% (**5a**), 66% (**5b**), and 67% (**5c**), respectively, on the basis of their 400 MHz ^1H nmr analyses at -80°C . Each RO-*ax* conformer preference extent of the thiane 1-oxides **5a-c** should be evidently larger than that of the corresponding thianes (**4a-c**) exhibiting the same order aspect **5c** > **5b** > **5a** as **4c** > **4b** > **4a**. In the cases of *trans*-4-RO-thiane 1-oxides (**6a-c**) and 1,1-dioxides (**7a-c**), the extreme 4-RO-*ax* conformer preference extent of 4-benzyloxy derivatives (**6a** and **7a**) seems to be little lower than that of the corresponding 4-siloxy ones (**6b,c** and **7b,c**). Tentative ^1H nmr (200 MHz, CDCl_3 , room temperature) analyses of 4-RO-dihydrothiines (**8a,c**) and 1,1-dioxide of **8c** provided the fairly small W values [18.3 Hz (**8a**), 19.7 Hz (**8c**), and 18.6 Hz (1,1-dioxide of **8c**)] due to their 4-H peaks, which should mean their 4-RO-*pseudoax* conformer preferences as we anticipated.⁸ The dynamic ^1H nmr analysis of **8c** at -95°C definitely clarified its conformational ratio.

Finally, crystalline compounds (**6c** and **7c**) were submitted to the X-ray analysis.⁹ Perspective views of their crystallographic structures are depicted in Figure 2.⁹ The thiane 1-oxide and 1,1-dioxide rings adopt a chair form and the siloxy group, in fact, occupies the *ax* site in their molecules.




Figure 2. Perspectives view of the crystallographic structures of compounds **6c** and **7c**.

REFERENCES AND NOTES

1. Y. Nagao, M. Goto, M. Ochiai, and M. Shiro, *Chem. Lett.*, 1990, 1503.
2. Y. Nagao, Y. Hagiwara, Y. Hasegawa, M. Ochiai, T. Inoue, M. Shiro, and E. Fujita, *Chem. Lett.*, 1988, 381; S. Hanessian, D. Delorme, S. Beaudoin, and Y. Leblanc, *J. Am. Chem. Soc.*, 1984, **106**, 5754; S. E. Denmark and C.-T. Chen, *J. Am. Chem. Soc.*, 1992, **114**, 10674; D. Sato, H. Kawasaki, I. Shimada, Y. Arata, K. Okamura, T. Date and K. Koga, *J. Am. Chem. Soc.*, 1992, **114**, 761.
3. J. C. Martin and J. J. Ubel, *J. Am. Chem. Soc.*, 1964, **86**, 2936.
4. M. Ōki and N. Nogami, The 32nd National Meeting of the Chemical Society of Japan, Tokyo, April 1975, Abstr., p. 1087; N. Nogami, S. Sato, M. Ōki, and Y. Saito, The 36th National Meeting of the Chemical Society of Japan, Higashi Osaka, April 1977, Abstr., p. 592.
5. Reduction of commercially available 4-kethothiane with LiAlH_4 in Et_2O followed by conventional benzylation or silylation employing each corresponding halide gave 4-benzyloxy- or 4-siloxythianes (**4a-d**) in a good yield, respectively. Their *cis*- and *trans*-thiane 1-oxides (**5a-c** and **6a-c**) and thiane 1,1-dioxides (**7a-c**) were prepared by oxidation with 1 or 2 mol equiv of *m*-chloroperoxybenzoic acid in CH_2Cl_2 . Chromatographic separation of the resultant mixture of *cis*- and *trans*-thiane 1-oxides on a silica gel plate

gave each pure isomer. Assignment of the *cis* or *trans* structure should be done by their ^1H nmr (200 MHz, CDCl_3 , room temperature) analyses [$W(4\text{-H}) = 24.3\text{-}25.6$ Hz for the *cis*-isomers or $W(4\text{-H}) = 18.5 - 21.0$ Hz for the *trans*-ones; The 3-H peaks of the *cis*-isomers are recognized in the higher magnetic field than those of the corresponding *trans*-ones.] in comparison with the same analyses of the authentic *trans*-(**6c**) and *cis*-(**5c**).

6. 4-Benzyloxy- and 4-siloxydihydrothiines (**8a,c**) were readily prepared by refluxing a mixture of the corresponding *cis*- and *trans*-thiane 1-oxides with Ac_2O in benzene.
7. B. Sternbach and A. G. MacDiarmid, *J. Am. Chem. Soc.*, 1961, **83**, 3384; R. West, L. S. Wilson, and D. L. Powell, *J. Organomet. Chem.*, 1979, **178**, 5; S. Shambayati, J. F. Blake, S. G. Wierschke, W. L. Jorgensen, and S. L. Schreiber, *J. Am. Chem. Soc.*, 1990, **112**, 697; Y. Nagao, M. Goto, and M. Ochiai, *Chem. Lett.*, 1990, 1507; K. Maruoka, J. Sato, and H. Yamamoto, *J. Am. Chem. Soc.*, 1991, **113**, 5449.
8. A. Quédraogo, M. T. P. Viet, J. K. Saunders, and J. Lessard, *Can. J. Chem.*, 1987, **65**, 1761; S. E. Denmark, M. S. Dappen, N. L. Sear, and R. T. Jacobs, *J. Am. Chem. Soc.*, 1990, **112**, 3466
9. The crystallographic data of compounds **6c** and **7c** are as follows **6c**: $\text{C}_{21}\text{H}_{28}\text{O}_2\text{SSi}$, $M = 372.60$, triclinic, $P1(\#2)$, $a = 10.240(1)\text{\AA}$, $b = 12.003(3)\text{\AA}$, $c = 9.991(2)\text{\AA}$, $\alpha = 111.44(2)^\circ$, $\beta = 107.04(1)^\circ$, $\gamma = 101.32(1)^\circ$, $V = 1027.2(4)\text{\AA}^3$, $z = 2$, $D_{\text{calc}} = 1.205\text{ g/cm}^3$, $R = 0.066$, **7c**: $\text{C}_{21}\text{H}_{28}\text{O}_3\text{SSi}$, $M = 388.597$, tetragonal, $P4_2/n$, $a = 20.316(3)\text{\AA}$, $c = 10.302(2)\text{\AA}$, $V = 4252(1)\text{\AA}^3$, $z = 8$, $D_{\text{calc}} = 1.214\text{ g/cm}^3$, $R = 0.066$.

Received, 17th November, 1994