

Pteridines, CIV.¹ REGIOSELECTIVE ALKOXYLATION OF PTERIDINES AT THE 6-POSITION BY *N*-BROMOSUCCINIMIDE AND ALCOHOL


Takashi Sugimoto*^{a)} and Wolfgang Pfeleiderer^{b)}

^{a)}Department of Natural Science Informatics, School of Informatics and Sciences, Nagoya University, Chikusa-ku, Nagoya 464-01, Japan and ^{b)}Faculty of Chemistry, University of Konstanz, D-78464 Konstanz, Germany

Abstract— Reaction of 1,3-dimethylumazine with *N*-bromosuccinimide or bromine in an appropriate alcohol afforded 6-alkoxy-1,3-dimethylumazine. *N*-Iodosuccinimide or iodine also brought about the same reaction. This procedure was applicable to other pteridine compounds, such as 2-dimethylamino-4(3*H*)-oxopterin. The mechanism of this alkylation reaction is discussed.

Substitution of a hydrogen at the pteridine ring by a nucleophilic reagent like alcohols or amines has not been recorded in the literature.² This is attributed to the competitive and more avid covalent addition of such reagent to the pteridine nucleus.² Indeed, almost all alkoxypteridines were synthesized either from the corresponding chloropteridines by nucleophilic replacement or from appropriate synthons bearing the alkoxy substituent. We describe here a new method to introduce an alkoxy group directly to the pteridine nucleus by the action of *N*-bromosuccinimide or bromine in an appropriate alcohol.

Reaction of 1,3-dimethylumazine (**1**) with *N*-bromosuccinimide (NBS; 2 eq.) in boiling methanol gave a monomethoxy derivative of **1** in 77% yield. The structure of the product was confirmed to be 6-methoxy-1,3-dimethylumazine (**2a**) by a direct comparison with the authentic 6-methoxy-³ and 7-methoxy-1,3-dimethylumazines⁴ by tlc on a silica gel plate and mixture melting point test. When the



Scheme 1. Formation of 6-alkoxy- and 6,7-dialkoxy-1,3-dimethylmazines (**2** and **3**) from **1** by NBS (or Br₂) in alcohol.

reaction was carried out by employing 6 molar eq. of NBS at room temperature for 7 days, 6,7-dimethoxy-1,3-dimethylmazine (**3a**)⁵ was also formed together with the 6-methoxy compound. No 7-methoxy-1,3-dimethylmazine could be detected.

When ethanol was used as the reaction medium in place of methanol, the corresponding 6-ethoxy-1,3-dimethylmazine (**2b**) and 6,7-diethoxy-1,3-dimethylmazine (**3b**) were obtained. A similar reaction in 1-propanol gave 1,3-dimethyl-6-propoxylmazine (**2c**). However, **1** did not react with NBS in 2-propanol and 2-methyl-2-propanol. In the above reactions, development of a color typical to bromine was observed, indicating that bromine might also bring about the alkoxylation reaction. Indeed, **1** and bromine in methanol produced the 6-methoxy- and 6,7-dimethoxy-1,3-dimethylmazines.


The regioselective formation of the 6-alkoxy-1,3-dimethylmazine (**2**) is best explained by a mechanism shown in Scheme 1. Namely, interaction of **1** with NBS or bromine at the 5,6-double bond to form the bromonium compound (**4**) seems to be the most probable initial step of the reaction. The cation (**4**) easily reacts with alcohol at the 6-position to give the ring opened product (**5**), which on subsequent elimination of hydrogen bromide forms the 6-alkoxy-1,3-dimethylmazines (**2**). An AM1 calculation on 1,3-dimethylmazine (**1**) supports this assumption: among the four possible reaction sites, the carbon at the 6-position has the highest HOMO as shown in Table 1. Electrophilic bromine first attacks this position to form the initial intermediate (**4**). An alternative path from **1** to **2** via an

Table 1. The atomic charges and HOMOs of 1,3-dimethylumazine and 2-dimethylamino-4(3*H*)-oxopteridine obtained by AM1 calculation.

position	1,3-Dimethylumazine		2-Dimethylamino-4(3 <i>H</i>)-oxopteridine	
	atomic charge*	HOMO	atomic charge*	HOMO
N-5	-0.059	0.066	-0.069	0.088
C-6	0.050	0.418	0.040	0.385
C-7	0.160	0.208	0.138	0.128
N-8	-0.197	0.198	-0.150	0.205

*Atomic charges with hydrogens summed into heavy atoms.

initial addition of an alcohol across the 5,6-double bond and subsequent oxidation of the adduct by NBS (or bromine) seems not plausible; no reaction of the substrate (**1**) was detected on a prolonged heating with manganese dioxide in methanol. The following three sets of reaction conditions were examined with an expectation to get 6-bromo- or 6-acetoxy-1,3-dimethylumazine: (A) **1** and bromine in carbon tetrachloride, (B) **1** and NBS in the presence of tetrabutylammonium bromide in chloroform, and (C) **1** and NBS in the presence of sodium acetate in acetic acid. Under these three conditions, no change of **1** was observed. However, addition of methanol to (A), or ethanol to (B) resulted in the production of 6-methoxy- or 6-ethoxy-1,3-dimethylumazine, respectively. Probably, alcohol contributes to decrease the activation energy leading to **4**. *N*-Iodosuccinimide and iodine also reacted with **1** in methanol, giving the 6-methoxy compound (**2a**), though the reaction proceeded much slower than with NBS and bromine. In an oxidation by peroxyacetic acid or hydrogen peroxide in an appropriate acid, **1** produces the *N*-oxide at the 5-position selectively,⁶ even though the atomic charge at the N-5 position is lower than that at the N-8 position. This regioselectivity owes much to steric hindrance by the methyl group at the 1-position, which prevents hydrogen peroxide or peroxyacetic acid to approach to the N-8 position.⁷ Indeed, those pteridines carrying no substituent at the 1-position, such as lumazine and 2-amino-4(3*H*)-oxopteridine, produce the *N*-oxides at the 8-position.⁶ If the present alkoxylation reaction proceeds through a mechanism similar to that for *N*-oxidation, 2-dimethylamino-4(3*H*)-oxopteridine (**6**) is expected to produce a 7-alkoxy derivative through an initial bromination on the more electron rich nitrogen atom at the 8-position, addition of alcohol to the 7,8-double bond, and subsequent elimination of hydrogen bromide. In order to prove the reaction mechanism shown in Scheme 1, we examined the



Scheme 2. Formation of 6-alkoxy- and 6-bromo-2-dimethylamino-4(3*H*)-oxopteridine (**7** and **8**) from **6** by NBS and alcohol.

reaction of 2-dimethylamino-4(3*H*)-oxopteridine (**6**) with NBS in alcohol. 2-Dimethylamino-4(3*H*)-oxopteridine (**6**) reacted much faster than 1,3-dimethylumazine (**1**) with NBS in methanol, giving 6-methoxy-2-dimethylamino-4(3*H*)-oxopteridine (**7b**) and 6-bromo-2-dimethylamino-4(3*H*)-oxopteridine (**8**). The same substrate with NBS in ethanol gave 6-ethoxy-2-dimethylamino-4(3*H*)-oxopteridine (**7c**) and the bromo compound (**8**). The position of the introduced alkoxy group was confirmed to be the 6-position by a direct comparison of its hydrolysis product with the authentic 2-dimethylamino-4,6(3*H*,5*H*)-dioxopteridine⁸ (**7a**) on a cellulose plate. The bromo compound (**8**), on heating with potassium *t*-butoxide in ethanol, gave the 6-ethoxy derivative (**7c**).

Here again, the substitution on **6** took place at the 6-position. It seems that a reaction path analogous to that in Scheme 1 proceeds also with **6**; addition of bromine across the 5,6-double bond to form the bromonium intermediate analogous to **4**, attack by alcohol to open to the three-member ring, and subsequent elimination of hydrogen bromide. Loss of the proton on C-6 of the bromonium intermediate leads to the 6-bromopteridine (**8**). Evidently, the 6-alkoxypteridines (**7**) were not formed *via* the 6-bromopteridine (**8**), because the latter was stable in hot methanol.

EXPERIMENTAL

6-Methoxy- and 6,7-dimethoxy-1,3-dimethylumazines (**2a** and **3a**)

(a) A solution of 1,3-dimethylumazine (**1**) (1.92 g, 10 mmol) and NBS (3.6 g, 20 mmol) in methanol (80 ml) was heated under reflux for 24 h. The solution was evaporated to dryness on a rotary evaporator. The residue was mixed with hot water (80 ml) and chilled. Filtration gave practically pure 6-

methoxy-1,3-dimethylumazine (**2a**) (1.72 g, 77%). Recrystallization from toluene gave colorless needles, mp 216-217 °C (lit.,³ 224 °C) (Anal. Calcd for C₉H₁₀N₄O₃: C, 48.65; H, 4.54; N, 25.22. Found: C, 48.76; H, 4.58; N, 22.25). ¹H Nmr (CDCl₃): δ 8.32 (1H, s, 7-H), 4.10 (3H, s, O-CH₃), 3.70 (3H, s, N-CH₃), and 3.52 (3H, s, N-CH₃); λ_{max} nm (log ε) in methanol: 245 (4.20), 337 (sh, 3.69), 354 (3.86), and 364 (sh, 3.81).

(b) A solution of **1** (0.48 g, 2.5 mmol) and NBS (2.5 g, 14 mmol) in methanol (100 ml) was stirred at room temperature for 7 days. The solution, after evaporation, extraction of the residue with chloroform (2 x 30 ml), and chromatography on a silica gel column (2 x 25 cm) eluted with a mixture of toluene and ethyl acetate (1:1, v/v), gave **2a** (0.15 g, 27%) and 6,7-dimethoxy-1,3-dimethylumazine (**3a**) (0.155 g, 25%) as colourless prisms, mp 217-218 °C (from toluene) (lit.,⁵ 226 °C) (Anal. Calcd for C₁₀H₁₂N₄O₄: C, 47.62; H, 4.80; N, 22.21. Found: C, 47.54; H, 4.75; N, 21.91). ¹H Nmr (CDCl₃): δ 4.16 (3H, s, O-CH₃), 4.15 (3H, s, O-CH₃), 3.66 (3H, s, N-CH₃), and 3.50 (3H, s, N-CH₃); λ_{max} nm (log ε) in methanol: 234 (sh, 4.04), 264 (3.98), 318 (sh, 3.92), 333 (4.05), and 342 (sh, 4.04).

Compound (**1**) (0.19 g, 1 mmol) and *N*-iodosuccinimide (0.45 g, 2 mmol) in methanol (20 ml) at room temperature for 3 weeks gave **2a** (40 mg, 18%).

(c) A solution of bromine (0.64 g, 4 mmol) in methanol (2 ml) was added to a boiling solution of **1** (0.48 g, 2.5 mmol) in methanol (20 ml). After refluxing for 1 h, the same amount of bromine was added and refluxing was continued for an additional 1 h. Then the solution was evaporated to dryness. The residue, on extraction, chromatography, and crystallization as above, gave **2a** (0.26 g, 47%) and **3a** (70 mg, 11%).

Compound (**1**) (0.48 g, 2.5 mmol) and iodine (1.65 g, 6.5 mmol) in methanol (100 ml) under reflux for 24 h gave **2a** (70 mg, 13%).

(d) A solution of **1** (0.48 g, 2.5 mmol) and bromine (3 g, 18 mmol) in tetrachloromethane (250 ml) was stirred at room temperature for 3 days. No reaction of the substrate was observed on tlc analysis. Then, methanol (20 ml) was added to the solution and stirred for 3 days. Evaporation and column chromatography as above, followed by crystallization from toluene, gave **2a** (0.19 g, 34%).

6-Ethoxy- and 6,7-diethoxy-1,3-dimethylumazines (2b and 3b)

(a) A solution of **1** (0.48 g, 2.5 mmol) and NBS (3 g, 17 mmol) in ethanol (150 ml) was stirred at room temperature for 4 days. Evaporation to dryness, extraction with chloroform, chromatography on a silica gel column, and crystallization from toluene as above gave 6-ethoxy-1,3-dimethylumazine (**2b**) as

colorless needles (0.27 g, 50%), mp 190-191 °C (Anal. Calcd for C₁₀H₁₂N₄O₃: C, 50.84; H, 5.12; N, 23.71. Found: C, 50.91; H, 5.14; N, 23.52) and 6,7-diethoxy-1,3-dimethylumazine (**3b**) as colorless needles (0.14 g, 20%), mp 180.5-181.5 °C (Anal. Calcd for C₁₂H₁₆N₄O₄: C, 51.42; H, 5.75; N, 19.99. Found: C, 51.35; H, 5.75; N, 19.82): **2b**: ¹H Nmr (CDCl₃): δ 8.30 (1H, s, 7-H), 4.53 (2H, q, *J* = 7.1 Hz, CH₂), 3.65 (3H, s, *N*-CH₃), 3.52 (3H, s, *N*-CH₃), and 1.44 (3H, t, *J* = 7.1 Hz, CH₃); λ_{max} nm (log ε) in methanol: 245 (4.19), 355 (3.84), and 363 (sh, 3.80). **3b**: ¹H Nmr (CDCl₃): 4.28 (2H, q, *J* = 7.1 Hz, CH₂), 4.47 (2H, q, *J* = 7.1 Hz, CH₂), 3.62 (3H, s, *N*-CH₃), 3.47 (3H, s, *N*-CH₃), 1.50 (3H, t, *J* = 7.1 Hz, CH₃), and 1.45 (3H, t, *J* = 7.1 Hz, CH₃); λ_{max} nm (log ε) in methanol: 236 (sh, 4.02), 265 (3.97), 320 (sh, 3.92), 334 (4.05), and 346 (sh, 3.96).

(b) A solution of **1** (0.48 g, 2.5 mmol), NBS (1.8 g, 10 mmol), and tetrabutylammonium bromide (1 g, 3 mmol) in chloroform (30 ml) was stirred at room temperature for 4 days. No reaction of the substrate was observed on tlc analysis. Ethanol (20 ml) was added and the solution was allowed to stand at room temperature for 7 days. The Product (**2b**) (0.155 g, 26%) was isolated from the reaction solution in an analogous way including chromatography and crystallization from toluene as above.

Compound (**1**) (0.48 g, 2.5 mmol) and bromine (3.2 g, 10 mmol) in ethanol (30 ml) under refluxing for 1 h gave **2b** (70 mg, 10%).

1,3-Dimethyl-6-propoxylumazine (2c)

A solution of **1** (0.48 g, 2.5 mmol) and NBS (4 g, 22 mmol) in 1-propanol (150 ml) was stirred at room temperature for 2 days. Evaporation to dryness, extraction with chloroform, chromatography on a silica gel column, and crystallization from toluene as above gave 1,3-dimethyl-6-propoxylumazine (**2c**) as colorless needles (0.10 g, 16%), mp 134-135 °C (Anal. Calcd for C₁₁H₁₄N₄O₃: C, 52.79; H, 5.64; N, 22.39. Found: C, 52.78; H, 5.69; N, 22.32). ¹H Nmr (CDCl₃): δ 8.29 (1H, s, 7-H), 4.41 (2H, q, *J* = 6.7 Hz, *O*-CH₂-), 3.68 (3H, s, *N*-CH₃), 3.50 (3H, s, *N*-CH₃), 1.92-1.77 (2H, m, CH₂), and 1.04 (3H, t, *J* = 7.4 Hz, CH₃); λ_{max} nm (log ε) in methanol: 245 (4.20), 355 (3.85), and 363 (sh, 3.81).

6-Methoxy- and 6-bromo-2-dimethylamino-4(3H)-oxopteridines (7b and 8)

A solution of 2-dimethylamino-4(3H)-oxopteridine (0.48 g, 2.5 mmol) and NBS (0.58 g, 3.2 mmol) in methanol (90 ml) was stirred at room temperature for 2 days. Silica gel (5 g) was added to the solution and the mixture was evaporated to dryness on a rotary evaporator. The solid residue was placed on the top of a silica gel column (2 x 20 cm) and eluted first with a mixture of acetonitrile and toluene (2:1, v/v). Evaporation and crystallization from methanol gave yellow needles (0.125 g, 19%) of 6-bromo-2-

dimethylamino-4(3*H*)-oxopteridine (**8**), mp 270-272 °C (decomp). (Anal. Calcd for C₈H₈N₅OBr: C, 35.57; H, 2.98; N, 25.93. Found: C, 35.56; H, 3.00; N, 25.80). ¹H Nmr (CDCl₃): δ 8.72 (1H, s, 7-H) and 3.33 [6H, s, N(CH₃)₂]; λ_{max} nm (log ε) in methanol: 222(sh, 4.04), 249 (3.93), 292(4.36), 370 (sh, 3.64), and 386 (3.68). The column was then eluted with a mixture of chloroform and methanol (9:1, v/v). Evaporation of the eluate and crystallization of the residue from methanol gave ivory needles (0.123 g, 22%) of 6-methoxy-2-dimethylamino-4(3*H*)-oxopteridine (**7b**), mp 235-237 °C (Anal. Calcd for C₉H₁₁N₅O₂: C, 48.86; H, 5.01; N, 31.66. Found: C, 48.85; H, 5.05; N, 31.46). ¹H Nmr (CDCl₃): δ 8.41 (1H, s, 7-H), 4.05 (3H, s, O-CH₃), and 3.27 [6H, s, N(CH₃)₂]; λ_{max} nm (log ε) in methanol: 229 (4.11), 249 (sh, 4.02), 283 (4.29), 348 (sh, 3.58), 367 (3.68), and 390 (sh, 3.56).

6-Ethoxy- and 6-bromo-2-dimethylamino-4(3H)-oxopteridines (7c and 8)

A solution of **6** (0.48 g, 2.5 mmol) and NBS (0.50 g, 2.8 mmol) in ethanol (150 ml) was stirred at room temperature for 4 days. Silica gel (5 g) was added to the solution and the mixture was evaporated to dryness on a rotary evaporator. The solid residue was placed on the top of a silica gel column (2 x 20 cm) and eluted with a mixture of chloroform and methanol (9:1, v/v). Evaporation of the first blue fluorescent eluate and crystallization of the residue from methanol gave yellow needles (75 mg, 11%) of **8**. Evaporation of the second blue fluorescent eluate and crystallization of the residue from methanol gave ivory needles (0.21 g, 36%) of 6-ethoxy-2-dimethylamino-4(3*H*)-oxopteridine (**7c**), which decomposed above 270 °C without melting (Anal. Calcd for C₁₀H₁₃N₅O₂: C, 51.06; H, 5.56; N, 29.77. Found: C, 51.00; H, 5.69; N, 29.58). ¹H Nmr (DMSO-*d*₆): δ 8.41 (1H, s, 7-H), 4.34 (2H, q, *J* = 7.1 Hz, CH₂), 3.09 [6H, s, N(CH₃)₂], and 1.35 (3H, t, *J* = 7.1 Hz, CH₃); λ_{max} / nm (log ε) in methanol: 230 (4.13), 248 (sh, 4.04), 283 (4.30), 349 (sh, 3.61), 367 (3.69), and 390 (sh, 3.58).

Hydrolysis of 6-ethoxy-2-dimethylamino-4(3H)-oxopteridine

A solution of 6-ethoxy-2-dimethylamino-4(3*H*)-oxopteridine (40 mg, 0.17 mmol) in 0.5 M hydrochloric acid (10 ml) was heated under gentle reflux for 45 min. The solution was treated with activated charcoal, neutralized with 1.4% ammonia, and evaporated to dryness. The residue was submitted to chromatography twice on a silica gel column (2 x 25 cm) by eluting with a mixture of chloroform and methanol (5:1, v/v). Evaporation of the eluate gave a solid, which was dissolved in hot ethanol and concentrated to about 1 ml. to give yellowish brown powder (8 mg, 23%) of 2-dimethylamino-4,6(3*H*,5*H*)-dioxopteridine (**7a**).

Conversion of 6-bromo-2-dimethylamino-4(3H)-oxopteridine into 6-ethoxy-2-dimethylamino-4(3H)-oxopteridine

A few milligrams of 6-bromo-2-dimethylamino-4(3H)-oxopteridine (**8**) and t-BuOK (about 20 mg) in ethanol (20 ml) were heated under reflux overnight. After neutralization with acetic acid, the product was identified as 6-ethoxy-2-dimethylamino-4(3H)-oxopteridine (**7c**) by a direct comparison on silica gel plates.

ACKNOWLEDGMENTS

We thank Mrs. M. Bischler for measuring the uv spectra, Dr. Thomas Wagner and Mr. S. Nishimura for measuring the nmr spectra, and Mr. K. Kiguchi for the AM1 calculation. Dr. W. Hutzenlaub kindly gave us several compounds, which were essential for the structure determination of several products. Thanks are also due to the Ministry of Education Science and Culture, Japan for offering a travel grant to T. S..

REFERENCES

1. Part CIII: K. Sharma and W. Pfeleiderer, *Indian J. Heterocycl. Chem.*, 1995, in pres.
2. D. J. Brown, "Fused Pyrimidines. Part Three. Pteridines," ed. by E. C. Taylor and A. Weissberger, John Wiley & Sons, New York, 1988, pp. 7-42.
3. W. Pfeleiderer, *Chem. Ber.*, 1957, **90**, 2604.
4. W. Pfeleiderer, *Chem. Ber.*, 1957, **90**, 2588.
5. W. Pfeleiderer, *Chem. Ber.*, 1957, **90**, 2631.
6. W. Pfeleiderer and W. Hutzenlaub, *Chem. Ber.*, 1973, **106**, 3149; W. Pfeleiderer and W. Hutzenlaub, *Angew. Chem.*, 1965, **77**, 1136; H. Yamamoto, W. Pfeleiderer, and W. Hutzenlaub, *Chem. Ber.*, 1973, **106**, 3175; W. Hutzenlaub, H. Yamamoto, G. B. Barlin, and W. Pfeleiderer, *Chem. Ber.*, 1973, **106**, 3203.
7. D. J. Brown, "Fused Pyrimidines. Part Three. Pteridines," ed. by E. C. Taylor and A. Weissberger, John Wiley & Sons, New York, 1988, pp. 309-311.
8. M. Böhme, W. Hutzenlaub, W. J. Richter, E. F. Elstner, G. Huttner, J. von Seyerl, and W. Pfeleiderer, *Liebigs Ann. Chem.*, 1986, 1705.

Received, 1st December, 1994