


## A FACILE AND CONVENIENT SYNTHESIS OF 9-METHYL-3-(1H-TETRAZOL-5-YL)-4H-PYRIDO[1,2-a]PYRIMIDIN-4-ONE POTASSIUM

Atsunori Sano\* and Masami Ishihara

*Tokyo Research Laboratories, Wako Pure Chemical Industries, Ltd., 1633 Matoba, Kawagoe, Saitama 350-11, Japan*

**Abstract** — Two convenient synthetic methods of anti-allergy agent, 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one potassium (**1a**), are described. Heating of a mixture of malononitrile (**2b**), 2-amino-3-picoline (**4**), ethyl orthoformate, and sodium azide in acetic acid, followed by hydrolysis with hydrochloric acid gave 9-methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one (**1b**) which was led to **1a** by treatment with potassium hydroxide. The use of ethyl cyanoacetate (**2a**) instead of **2b** afforded ethyl 3-(3-methyl-2-pyridylamino)-2-(1H-tetrazol-5-yl)acrylate (**8**) which was also led to **1a** by treatment with potassium hydroxide.


9-Methyl-3-(1H-tetrazol-5-yl)-4H-pyrido[1,2-a]pyrimidin-4-one potassium (**1a**),<sup>1</sup> anti-allergy agent, has been clinically used for the treatment of asthma.<sup>2</sup> The synthesis of **1a** starting from ethyl cyanoacetate (**2a**) via ethyl 2-cyano-3-(3-methyl-2-pyridylamino)acrylate (**5**) was reported by Juby (Route A in Scheme 1).<sup>3</sup> It involved the formation of a tetrazole ring using a hazardous reagent, aluminum azide<sup>4</sup> and its yield is low.



Scheme 1

Recently, we developed an alternative synthesis of **1a** starting from malononitrile (**2b**) via 3-cyano-4-imino-9-methyl-4*H*-pyrido[1,2-*a*]pyrimidine (**6**) (Route B in Scheme 1).<sup>5</sup> In this synthesis, **6** was converted in acetic acid into **7** in good yield by the treatment of sodium azide, which was safer to handle than aluminum azide. However, this method required multi-step to afford **1a** from a commercially available material (**2b**) and handling of a strong sensitizing agent (**3b**).

In this note, we wish to describe two kinds of a facile and convenient preparation of **1a** using a one-pot synthesis of **1b** or **8** (Scheme 2).



Scheme 2

It was reported that the pyrido[1,2-*a*]pyrimidine skeleton of **1a** was prepared by condensation of **2b**, 2-amino-3-picoline (**4**), and ethyl orthoformate.<sup>6</sup> In addition, we found that acetic acid was an efficient solvent for conversion of a cyano group into a 1*H*-tetrazole ring with sodium azide.<sup>4</sup> Therefore, the condensation reaction in acetic acid containing sodium azide was expected to attain both pyrido[1,2-*a*]pyrimidine and 1*H*-tetrazole ring closures.

First, we investigated a reaction of **2b** and **4** with ethyl orthoformate in acetic acid containing sodium azide (Route C in Scheme 2). Heating of a mixture of **2b**, **4**, ethyl orthoformate and sodium azide in acetic acid, followed by hydrolysis with hydrochloric acid gave 9-methyl-3-(1*H*-tetrazol-5-yl)-4*H*-pyrido[1,2-*a*]pyrimidin-4-one (**1b**) in 68% yield. The yield was not influenced by a way of blend of the reagents. The structure of **1b** was confirmed by direct comparison with an authentic sample.<sup>3</sup> Treatment of **1b** with aqueous potassium hydroxide afforded **1a** in quantitative yield.

When this condensation reaction was carried out in other acidic solvents, such as formic acid or trifluoroacetic acid, many unidentified by-products were produced. Thus, the pH of the reaction mixture was seemed to be one of the critical factors both for conversion of a cyano group into 1*H*-tetrazole ring and the condensation of **2a** and **4** with ethyl orthoformate.

Next, we tried to use **2a** in place of **2b** (Route D in Scheme 2). Condensation of a mixture of **2a**, **4**, ethyl

orthoformate and sodium azide in acetic acid afforded ethyl 3-(3-methyl-2-pyridylamino)-2-(1*H*-tetrazol-5-yl)-acrylate (**8**) as crystalline powder in 62% yield.

The structure of **8** was determined on the basis of IR and <sup>1</sup>H-NMR spectral data. The IR spectrum indicated the absence of cyano group. The <sup>1</sup>H-NMR spectrum showed a doublet signal at 4.35 ppm and a triplet signal at 1.35 ppm. These signals were assigned to the protons of an ethyl group. Moreover, two doublet signals observed at 11.44 and 9.23 ppm with a coupling constant of 13.5 Hz suggested the presence of a =CH-NH-moiety.

Treatment of **8** with aqueous potassium hydroxide afforded **1a** in 95% yield while attempts of cyclization of **8** under aqueous acidic conditions, such as aqueous hydrochloric acid or sulfuric acid, were unsuccessful.

In summary, we accomplished two convenient methods for the synthesis of **1a** employing a one-pot synthesis of **1b** or **8**. These methods were able to provide **1a** in only two steps starting from **2**.

## EXPERIMENTAL

All melting points are determined on a Yanagimoto apparatus and are uncorrected. IR spectra were recorded on a JASCO IRA-2 spectrophotometer using KBr disks. <sup>1</sup>H-NMR spectra were recorded on a JEOL GSX 270 spectrometer with tetramethylsilane as an internal standard. All reagents and solvents were of commercial quality.

### 9-Methyl-3-(1*H*-tetrazol-5-yl)-4*H*-pyrido[1,2-*a*]pyrimidin-4-one (**1b**) (from **2b**)

A mixture of malononitrile (**2b**) (3.3 g, 50 mmol), 2-amino-3-picoline (**4**) (5.4 g, 50 mmol), ethyl orthoformate (8.2 g, 55 mmol) and sodium azide (3.6 g, 55 mmol) in acetic acid (50 mL) was heated at 90°C for 2 h. Concentrated hydrochloric acid (25 mL) was poured into the mixture, which was further heated at 90°C for 2 h. The mixture was cooled and diluted with water (50 mL). The resulting precipitates were collected in suction and dried to give **1b** (7.7 g, 67.5%), mp 306°C(dec.) (lit.,<sup>3</sup> 310-311°C(dec.)). All spectral data of **1b** were identical with those of an authentic sample.<sup>3</sup> IR (KBr): 3100, 1670, 1595, 1480, 1330, 1080, 900, 780 cm<sup>-1</sup>. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): 2.63 (3H, s, CH<sub>3</sub>), 7.54 (1H, t, *J*=7.0 Hz, C7-H), 8.09 (1H, d, *J*=7.0 Hz, C8-H), 9.15 (1H, d, *J*=7.0 Hz, C6-H), 9.24 (1H, s, C2-H).

### 9-Methyl-3-(1*H*-tetrazol-5-yl)-4*H*-pyrido[1,2-*a*]pyrimidin-4-one potassium (**1a**) (from **1b**)

A suspension of **1b** (4.6 g, 20 mmol), 2*N* potassium hydroxide (15 mL) in 2-propanol (20 mL) was heated at 50°C for 1 h. The mixture was cooled and diluted with 2-propanol (40 mL). The resulting precipitates were collected in suction filtration and dried to give **1a** (5.3 g, 99%) as pale yellow crystalline powder. mp 317°C(dec.). All spectral data of **1a** were identical with those of an authentic sample.<sup>3</sup> IR (KBr): 1690, 1495, 1450, 1310, 1160, 1140, 770 cm<sup>-1</sup>. <sup>1</sup>H-NMR (D<sub>2</sub>O): 2.15 (3H, s, CH<sub>3</sub>), 6.97(1H, t, *J*=7.0 Hz, C7-H), 7.38(1H, d, *J*=7.0 Hz, C8-H), 8.29 (1H, s, C2-H), 8.64(1H, d, *J*=7.0 Hz, C6-H).

### Ethyl 3-(3-methyl-2-pyridylamino)-2-(1*H*-tetrazol-5-yl)acrylate (**8**)

A mixture of ethyl cyanoacetate (**2a**) (5.7 g, 50 mmol), **4** (5.4 g, 50 mmol), ethyl orthoformate (8.2 g, 55 mmol) and sodium azide (3.6 g, 55 mmol) in acetic acid (50 mL) was heated at 90°C for 2 h. After the mixture was

cooled and diluted with water (50 mL), the resulting precipitates were collected in suction filtration and dried to give **8** (8.4 g, 61 %) as pale yellow leaflets. mp 206-210°C. IR (KBr): 3150, 1660, 1620, 1580, 1460, 1400, 1240, 1100, 770 cm<sup>-1</sup>. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): 1.35 (3H, t, *J*=8.1 Hz, CH<sub>2</sub>CH<sub>3</sub>), 2.42 (3H, s, CH<sub>3</sub>), 4.35 (2H, q, *J*=8.1 Hz, CH<sub>2</sub>CH<sub>3</sub>), 7.12 (1H, dd, *J*=8.1 and 5.4 Hz, Ar-H(C5)), 7.73 (1H, d, *J*=8.1 Hz, Ar-H(C4)), 8.26 (1H, d, *J*=5.4 Hz, Ar-H(C6)), 9.23 (1H, d, *J*=13.5 Hz, NHCH=C), 11.44 (1H, d, *J*=13.5 Hz, NHCH=C), 14.32 (1H, s, tetrazole-H). *Anal.* Calcd for C<sub>12</sub>H<sub>14</sub>N<sub>6</sub>O<sub>2</sub>: C, 52.55; H, 5.14; N, 30.64. Found: C, 52.47; H, 4.98; N, 30.42.

#### 9-Methyl-3-(1*H*-tetrazol-5-yl)-4*H*-pyrido[1,2-*a*]pyrimidin-4-one potassium (**1a**) (from **8**)

A suspension of **8** (5.5 g, 20 mmol), 2*N* potassium hydroxide (15 mL) and 2-propanol (20 mL) was refluxed for 2 h. The mixture was cooled and diluted with 2-propanol (20 mL). The resulting precipitates were collected in suction filtration and dried to give **1a** (5.2 g, 95 %) as pale yellow crystalline powder. All spectral data of **1a** were identical with those of an authentic sample as above.

#### REFERENCES AND NOTES

1. The generic name is pemirolast potassium.
2. a) J. J. Wade, C. B. Toso, C. J. Matson, and V. L. Stelzer, *J. Med. Chem.*, 1983, **26**, 608. b) P. F. Juby, T. W. Hudyma, M. Brown, J. M. Essery, and R. A. Partyka, *J. Med. Chem.*, 1982, **25**, 1145.
3. P. F. Juby, U. S. Patent 4,122,274 [*Chem. Abstr.*, 1979, **90**, 103998].
4. a) B. E. Huff, M. A. Staszak, *Tetrahedron Lett.*, 1993, **34**, 8011. b) N. I. Sax and R. J. Lewis, Sr., (eds.) "In Dangerous Properties of Industrial Materials 7th ed." Vol. 2, Van Nostrand Reinhold, New York, 1989, p. 135.
5. A. Sano, M. Ishihara, J. Yoshihara, M. Sumino, and H. Nawa, *Chem. Pharm. Bull.*, 1995, **43**, 683.
6. a) B. Podányi, I. Hermeecz, and Á. Horváth, *J. Org. Chem.*, 1986, **51**, 2988. b) Y. Okamoto, Y. Kurasawa, K. Takagi, A. Takada, and T. Ueda, *Chem. Pharm. Bull.*, 1974, **22**, 243.

Received, 10th December, 1997