

NOVEL SYNTHESIS OF DIAZETIDINE-2,4-DIONE BY RING EXPANSION OF DIAZIRIDINONE


Mitsuo Komatsu,*^a Satoshi Tamabuchi,^a Satoshi Minakata,^a and Yoshiki Ohshiro^b

^aDepartment of Applied Chemistry, Faculty of Engineering, Osaka University
Yamadaoka 2-1, Suita, Osaka 565-0871, Japan

^bResearch Institute for Science and Technology, Kinki University
Kowakae 3-4-1, Higashi-Osaka, Osaka 577-0818, Japan

Abstract—Treatment of *N,N'*-di-*tert*-butyldiaziridinone with Ni(CO)₄ under an atmosphere of carbon monoxide caused a carbonylative ring expansion to give di-*tert*-butyldiazetidinedione in good yield. In the presence of diphenylketene under the conditions, azetidinedione derivative was obtained.

Three-membered heterocycles have long been known as potential building blocks for the synthesis of numerous types of heterocycles.¹ Among such small rings, diaziridinones² would be expected to show unique ring-opening reactivity because of their highly strained three-membered structures which consist of a carbonyl carbon and two nitrogen atoms.¹⁻³ In fact, cycloaddition and addition-cyclization reactions of *N,N'*-di-*tert*-butyldiaziridinone (**1**) have already been reported by our group⁴ and by Greene and co-workers.⁵ It is also noteworthy that the carbonylative ring expansion of heterocycles has been developed by Alper's group,⁶ especially as this relates to small ring heterocycles. A carbonylation reaction of this type represents a useful method in organic synthesis, since it provides a convenient and effective one-pot procedure for ring homologation. From these points of view, we report herein the transition metal-mediated carbonylative ring expansion of a diaziridinone (**1**) leading to 1,3-diazetidione derivative. It is well known that diazetidinedione derivatives can be converted into functional materials and that some have fungicidal, herbicidal and insecticidal activities.⁷ These facts prompted us to investigate this reaction as a new procedure for the synthesis of diazetidinedione.⁸



Treatment of *N,N'*-di-*tert*-butyldiaziridinone (**1**) with an equimolar amount of Ni(CO)₄ under an atmosphere of carbon monoxide in DMF at 50 °C for 3 h afforded di-*tert*-butyldiazetidinedione (**2**) in 62% yield, along with *N,N'*-di-*tert*-butylurea (**3**) in 13% yield (Table 1, run 1). The suitable reaction


temperature and time were found to be 70 ~ 90 °C and 1 ~ 3 h, respectively (runs 2-5).⁹ Since the yield of the reaction was drastically decreased when the reaction was run under nitrogen atmosphere, CO was required for the reaction (run 6). When 0.1 equiv of Ni(CO)₄ was employed, a small amount of **2** was obtained under the conditions (run 7). The use of THF as a solvent instead of DMF was not effective for the reaction (run 8). Diazetidinedione (**2**) was not obtained in the case with Fe(CO)₅ or W(CO)₆ instead of Ni(CO)₄ (runs 9, 10). It was found that 0.1 equiv of Pd(PPh₃)₄ catalyzed the present reaction to give **2** in 35% yield (run 11).¹⁰ These results indicate that Ni(CO)₄ and Pd(PPh₃)₄ both represent viable catalysts for the insertion of CO into the nitrogen-nitrogen bond of diaziridinone (**1**). A ring expansion accompanied by the fission of the nitrogen-nitrogen bond of **1** is very rare, but one instance of which has been previously reported by us.⁴

Table 1. Carbonylation of Diaziridinone **1** Using Metal Complex under a CO Atmosphere

run	metal complex (equiv)	solv.	atmosphere	temp. (°C)	time (h)	yield (%)	
						2	3
1	Ni(CO) ₄ (1)	DMF	CO	50	3	62	13
2	Ni(CO) ₄ (1)	DMF	CO	70	3	71	13
3	Ni(CO) ₄ (1)	DMF	CO	70	1	75	15
4	Ni(CO) ₄ (1)	DMF	CO	70	10	72	17
5	Ni(CO) ₄ (1)	DMF	CO	90	3	74	15
6	Ni(CO) ₄ (1)	DMF	N ₂	70	3	37	10
7	Ni(CO) ₄ (0.1)	DMF	CO	70	3	1	12
8	Ni(CO) ₄ (1)	THF	CO	reflux	3	25	25
9	Fe(CO) ₅ (1)	DMF	CO	70	3	0	26
10	W(CO) ₅ (1)	DMF	CO	70	3	0	12
11	Pd(PPh ₃) ₄ (0.1)	DMF	CO	70	3	35	26


A possible mechanism for carbonylative ring expansion of diaziridinone (**1**) is shown in Scheme 1. The oxidative addition of diaziridinone could occur on the metal [Ni(CO)₄ or Pd(PPh₃)₃CO] with

Scheme 1. Possible Mechanism for Formation of **2** and **3**.



cleavage of the nitrogen-nitrogen bond of **1**. Subsequently, the insertion of carbon monoxide into the nitrogen-metal bond of complex (**4**) could proceed to give metallacycle (**5**), followed by reductive elimination to afford diazetidinedione (**2**). The urea derivative (**3**) would be formed by the reaction of the unreacted complex (**4**) with water either in DMF or during work up.

The above results led us to investigate the application of the present reaction system to another type of reaction, which involves a dipolarophile. When diaziridinone (**1**) was treated with diphenylketene (**6**) in the presence of $\text{Ni}(\text{CO})_4$ under an atmosphere of CO, the azetidinedione derivative (**7**) was obtained in 27% yield, along with **2** and **3**. Azetidinedione (**7**) does not represent a 1 : 1 cycloadduct of **1** with diphenylketene, but formally that of *N*-*tert*-butylisocyanate with diphenylketene. Although the reaction paths leading to **7** are not clear, possible pathways can be considered. Since compound (**7**) was not obtained by the reaction of isolated **2** with **6** under the present conditions, diphenylketene would react with either intermediate (**4**) followed by insertion of CO and reductive elimination, or intermediate (**5**) followed by reductive elimination, to give **7**. While azetidine-2,4-diones can be prepared by the carbonylation of α -lactams,¹¹ our findings demonstrate the possibility of a unique ring expansion reaction with appropriate dipolarophiles accompanied by carbonylation.



In summary, we have demonstrated here a novel synthesis of diazetidine-2,4-dione by a ring expansion reaction of *N,N'*-di-*tert*-butyldiaziridinone with carbon monoxide utilizing $\text{Ni}(\text{CO})_4$ or $\text{Pd}(\text{PPh}_3)_4$. The present reaction is the first approach to such heterocycles by CO insertion into diaziridinone involving cleavage of its N-N bond. Interestingly, the reaction proceeded catalytically in the case with $\text{Pd}(\text{PPh}_3)_4$. Moreover, a new methodology to synthesize the 4-membered rings has been developed by the reaction of diaziridinone- $\text{Ni}(\text{CO})_4$ -CO system with diphenylketene. An investigation of this reaction in the presence of other dipolarophiles is now underway.

REFERENCES AND NOTES

- For reviews, see: W. H. Pearson, B. W. Lian, S. C. Bergmeiser, K. Madavu, L. Rai, and A. Hassner, 'Comprehensive Heterocyclic Chemistry II,' Vol. 1A, ed. by A. Padwa, Pergamon Press, Oxford, 1996, pp. 1-96; A. Padwa and A. D. Woolhouse, 'Comprehensive Heterocyclic Chemistry,' Vol. 7, ed. by W. Lwowski, Pergamon Press, Oxford, 1984, pp. 47-93.
- F. D. Greene, J. C. Stowell, and W. R. Bergmark, *J. Org. Chem.*, 1969, **34**, 2254; F. D. Greene and J. C. Stowell, *J. Am. Chem. Soc.*, 1964, **86**, 3569.
- P. R. Kumar, *Indian J. Chem. B*, 1985, **24B**, 678; P. E. McGann, J. T. Groves, F. D. Greene, G. M. Stack, J. Richard, and L. M. Trefonas, *J. Org. Chem.*, 1978, **43**, 922; J. F. Liebman and A.

- Greenberg, *J. Org. Chem.*, 1974, **39**, 123; F. D. Greene, W. R. Bergmark, and J. G. Pacifici, *J. Org. Chem.*, 1969, **34**, 2263.
4. M. Komatsu, M. Kobayashi, S. Itoh, and Y. Ohshiro, *J. Org. Chem.*, 1993, **58**, 6620; M. Komatsu, Y. Kajiwara, M. Kobayashi, S. Itoh, and Y. Ohshiro, *J. Org. Chem.*, 1992, **57**, 7359; M. Komatsu, T. Yagii, and Y. Ohshiro, *Tetrahedron Lett.*, 1990, **31**, 5327; T. Hirao, T. Masunaga, Y. Ohshiro, and T. Agawa, *Synthesis*, **1983**, 477; Y. Ohshiro, M. Komatsu, Y. Yamamoto, K. Takaki, and T. Agawa, *Chem. Lett.*, **1974**, 383.
 5. C. A. Renner and F. D. Greene, *J. Org. Chem.*, 1976, **41**, 813.
 6. E. P. Marcelo and H. Alper, *J. Am. Chem. Soc.*, 1996, **118**, 111; K. Khumtaveeporn and H. Alper, *Acc. Chem. Res.*, 1995, **28**, 414.
 7. J. W. Timberlake and E. S. Elder, 'Comprehensive Heterocyclic Chemistry,' Vol. 7, ed. by W. Lwowski, Pergamon Press, Oxford, 1984, p. 484.
 8. Only one preparative method has been reported to date; see, J. C. Stowell, *J. Org. Chem.*, 1971, **36**, 3056.
 9. The general procedure is as follows: A solution of diaziridinone (**1**) (199 μL , 1 mmol) and $\text{Ni}(\text{CO})_4$ (130 μL , 1 mmol) in DMF (2.3 mL) was stirred under a carbon monoxide atmosphere at 70 °C. After 1 h, 1N HCl was added, and the resulting white precipitate was collected by filtration and purified by sublimation (80 °C, 1 mmHg) to give **2** (148.5 mg) in 75% yield. The filtrate was extracted with ether (100 mL X 3), washed with water (50 mL X 3) and dried over MgSO_4 . After filtration, the solvent was evaporated to give **3** (25.4 mg) in 15% yield.
 10. It is known that the reaction of $\text{Pd}(\text{PPh}_3)_4$ with CO affords $\text{Pd}(\text{PPh}_3)_3\text{CO}$; see, H. Alper, C. P. Perera, and F. R. Ahmed, *J. Am. Chem. Soc.*, 1981, **103**, 1289.
 11. D. Roberto and H. Alper, *Organometallics*, 1984, **3**, 1767.

Received, 1st July, 1998