

THE ISOLATION AND STRUCTURE OF PUMILIOTOXIN 341A:  
A NOVEL CYCLIC ETHER FROM THE FROG *EPIPEDOBATES*  
*TRICOLOR*

Poonam Jain,<sup>†</sup> Thomas F. Spande, H. Martin Garraffo, and John W. Daly\*








Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive  
and Kidney Diseases, National Institutes of Health, Bethesda, MD, 20892, USA

<sup>†</sup>Present address: c/o Rahul Jain, Dept. of Med. Chem., Nat. Inst. of  
Pharmaceutical Education and Research, Sector 67, SAS Nagar (Mohali) 160  
062, Punjab, India

**Abstract** - An alkaloid **341A** isolated by HPLC from skin extracts of the  
Ecuadoran poison frog *Epipedobates tricolor* was characterized by FTIR, MS and  
<sup>1</sup>H-NMR spectral analysis and assigned a spiro-fused pyranoindolizidine  
structure (**1**). It represents the first member of a small group of pumiliotoxin  
alkaloids to contain a cyclic ether moiety.

The isoprene-containing pumiliotoxin (PTX) class of alkaloids is one of the largest and most ubiquitous of the poison frog skin alkaloids.<sup>1-3</sup> These are 6-alkylideneindolizidines having a methyl and a tertiary hydroxyl group at C-8 and, in the case of the alloPTX subclass, an additional hydroxyl group at C-7. They have been detected as major or minor alkaloids in frogs from Central and South America, Australia and Madagascar; many have potent myotonic and/or cardiotoxic activity. A large variety of alkylidene side-chains have been characterized, comprising alkaloids totaling 13-21 carbons. There is another homologous class, the homopumiliotoxins (homoPTXs), comprising alkaloids of 14-21 carbons having 7-alkylidenequinolizidine structures. Approximately seventy-five members of the PTX/homoPTX classes have been reported, a small number of which lack either the tertiary hydroxyl or the tertiary methyl groups at C-8 in the PTX class or at C-9 in the homoPTX class. The largest subgroup (*ca.* 30) is comprised of 19-carbon alkaloids. Many of this group have additional hydroxyl or carbonyl groups in the side chain, the most common being pumiliotoxin A (**2**) and pumiliotoxin B (**3**), coded as **307A** and **323A**, respectively. An allopumiliotoxin coded as **323B** (**4**), is also widely encountered. Alkaloids **323A** (**3**) and **323B** (**4**) have the three hydroxyl groups apparently essential for maximum cardiotoxic activity.<sup>4</sup>

Alkaloids **339A** (**5**) and **339B** (**6**) are  $7\beta$ - and  $7\alpha$ -hydroxy-substituted alloPTXs, respectively, with the same *threo* 15,16-diol in the side-chain as **3**. Another alkaloid, **341A**, like **339A** and **339B** was shown by HRMS to be a tetra-oxygenated  $C_{19}$  alkaloid and, by MS, to be clearly of the PTX class. Alkaloid **341A** had, however, by GC-MS deuterium-exchange measurement, only three exchangeable hydrogens. Since it had no carbonyl absorption in its GC-FTIR spectrum (Figure 1), one oxygen had to be present as an ether or epoxide. The  $^1\text{H-NMR}$  analysis below locates this oxygen as being in a 6,13-ether linkage, creating a pyran ring spiro-fused to a 6,10-dihydro-7-hydroxylindolizidine nucleus, and can be visualized as arising by intramolecular cyclization of a 13,14-dihydroxy alloPTX as shown below. Uncyclized alloPTX 13,14-diols have yet to be detected in frog skin. A C-13-C-14 double bond is, however, often present in 19-carbon PTXs and alloPTXs.



## RESULTS AND DISCUSSION


A typical GC-EI-MS spectrum indicated the following molecular and fragment ions with intensities in parentheses (see EXPERIMENTAL for other spectra; these were instrument-dependent suggesting a thermal component): 341 (3), 323 (9), 298 (8), 266 (7), 254 (12), 182 (11), 114 (32), 112 (90), 84 (77), 70 (100). CI-MS with  $\text{NH}_3$  gave a protonated parent ion at  $m/z$  342, confirming the EI-MS molecular ion of  $m/z$  341. CI-MS with  $\text{ND}_3$  gave a deuterated molecular ion of  $m/z$  346 indicating three exchangeable hydrogens. EI-MS with an  $\text{ND}_3$  leak gave spectra of D-exchanged material where the following maximum numbers of deuterium-exchanged hydrogens are indicated in parentheses for the main EI fragment ions:  $m/z$  323 (2D), 298 (3D), 266 (1D), 254 (2D), 184 (2D), 182(2D), 126 (0D), 112 (1D), 84 (0D), 70 (50% 1D). The following molecular and fragment ion formulae were determined by HRMS.

Table 1. Summary of HRMS data on alkaloid **341A** (**1**)

341	$\text{M}^+$	$\text{C}_{19}\text{H}_{35}\text{NO}_4$	184	$(\text{M}^+ - \text{C}_9\text{H}_{17}\text{O}_2)$	$\text{C}_{10}\text{H}_{18}\text{NO}_2$
323	$(\text{M}^+ - \text{H}_2\text{O})$	$\text{C}_{19}\text{H}_{33}\text{NO}_3$	182	$(\text{M}^+ - \text{C}_9\text{H}_{19}\text{O}_2)$	$\text{C}_{10}\text{H}_{16}\text{NO}_2$
306	$(\text{M}^+ - \text{H}_2\text{O} - \text{OH})$	$\text{C}_{19}\text{H}_{32}\text{NO}_2$	166	$(\text{M}^+ - \text{C}_9\text{H}_{19}\text{O}_3)$	$\text{C}_{10}\text{H}_{16}\text{NO}$
298	$(\text{M}^+ - \text{C}_3\text{H}_7)^a$	$\text{C}_{16}\text{H}_{28}\text{NO}_4$	126	$(\text{M}^+ - \text{C}_{12}\text{H}_{23}\text{O}_3)$	$\text{C}_7\text{H}_{12}\text{NO}$
266	$(\text{M}^+ - \text{C}_3\text{H}_7\text{O}_2)$	$\text{C}_{16}\text{H}_{28}\text{NO}_2$	114	$(\text{M}^+ - \text{C}_{13}\text{H}_{23}\text{O}_3)$	$\text{C}_6\text{H}_{12}\text{NO}$
254	$(\text{M}^+ - \text{C}_5\text{H}_{11}\text{O})$	$\text{C}_{14}\text{H}_{24}\text{NO}_3$	112	$(\text{M}^+ - \text{C}_{12}\text{H}_{23}\text{NO}_3)$	$\text{C}_7\text{H}_{12}\text{O}$
209	$(\text{M}^+ - \text{C}_7\text{H}_{16}\text{O}_2)$	$\text{C}_{12}\text{H}_{19}\text{NO}_2$	84	$(\text{M}^+ - \text{C}_{14}\text{H}_{25}\text{O}_4)$	$\text{C}_5\text{H}_{10}\text{N}$
198	$(\text{M}^+ - \text{C}_7\text{H}_{13}\text{NO}_2)$	$\text{C}_{12}\text{H}_{22}\text{O}_2$	70	$(\text{M}^+ - \text{C}_{15}\text{H}_{27}\text{O}_4)$	$\text{C}_4\text{H}_8\text{N}$

<sup>a</sup>The error here in the best of three trials at peak matching was + 46 ppm; all other fragment ion mass measurements were within  $\pm 5.6$  ppm. The  $m/z$  298 ion profiled exactly with the total ion current of the **341A** GC peak and also, because of its D-exchange value, is not considered to be an impurity ion. EI-MS/MS on  $m/z$  298 shows major ions at  $m/z$  280 and 262, consistent with the loss of 2 molecules of  $\text{H}_2\text{O}$ .

The data of Table 1 indicate a loss of water and hydroxyl from the molecular ion, the loss of the elements of a propyl group, and a loss of a pentyloxy moiety. The last two losses, unprecedented in PTX or alloPTX alkaloids, are accommodated by the structural elements of one of the pyran substituents. A  $\text{C}_3\text{H}_7\text{O}_2$  loss to give the ion at  $m/z$  266 may involve the extrusion of carbons 7-9 with their hydroxyl groups by the fragmentation shown below (see Scheme 1). Fragmentations to give the  $m/z$  298 and 254 ions are also proposed. The peaks at  $m/z$  182 and 70 are typical of alloPTXs, while a major ion at  $m/z$  84 is typical of hydrogenated PTXs and alloPTXs in which the 6,10-double bond has been reduced.<sup>1,5</sup> Peaks at  $m/z$  114 and 112 are typical of alloPTXs.<sup>1,5</sup>



Scheme 1. Some MS spectral fragmentations proposed for alkaloid **341A** (**1**).

The FTIR spectrum of **1** (Figure 1) indicated two  $\nu_{OH}$  absorptions in an approximately 2:1 ratio at 3568 and *ca.* 3536  $\text{cm}^{-1}$ , the latter frequency typical of the internally hydrogen-bonded 8-hydroxyl group found in all pumiliotoxins with the exception of the small deoxy class. The 3650  $\text{cm}^{-1}$  absorption seen with the side-chain pumiliotoxin C-15 or C-16 hydroxyl absorptions or the 7 $\beta$ -hydroxyl stretching frequency of some allo-pumiliotoxins is now only a shoulder. The 3568  $\text{cm}^{-1}$  absorption may reflect intramolecular hydrogen bonding also. The strong Bohlmann bands at 2817  $\text{cm}^{-1}$  is consistent with an indolizidine structure, but is significantly stronger than the Bohlmann bands for alkylidene-containing PTXs. A broad and strong absorption at 1069  $\text{cm}^{-1}$ , the position of a C-O stretching vibration, supports one or more hydroxyl or ether groups. No carbonyl absorbance was detected.

The  $^1\text{H-NMR}$  spectrum obtained after HPLC purification (see Experimental) was taken first in  $\text{D}_2\text{O}$ , then  $\text{C}_6\text{D}_6$ . Both revealed four methyl signals, two singlets at approximately  $\delta$  1.15 and 1.34 in  $\text{D}_2\text{O}$  or  $\delta$  1.55 and 0.92 in  $\text{C}_6\text{D}_6$ . A doublet ( $J=6.6$  Hz) and triplet ( $J=7.2$  Hz) were seen at  $\delta$  0.96 and 0.87 in  $\text{D}_2\text{O}$  and  $\delta$  0.80 and 0.85 in  $\text{C}_6\text{D}_6$ , respectively. The most downfield methyl in  $\text{C}_6\text{D}_6$  is assigned to the 9- $\text{CH}_3$  and

the other singlet to a methyl attached to another hydroxyl-bearing carbon.




Figure 1. GC-FTIR spectrum of alkaloid **341A (1)**.

The most revealing structural details emerged from deuterobenzene spectra and the following discusses that data (see Table 2). The spectrum revealed clearly the distinctive  $^1\text{H-NMR}$  signals and multiplicities for indolizidine H- $3_{ax}$  and H- $5_{eq}$  protons at  $\delta$  2.04 (an apparent q,  $J=8.8$  Hz) and 2.63 (d,  $J=10.6$  Hz), respectively and confirms that ring system.<sup>6</sup> A doublet with a small long-range coupling (*ca.* 1 Hz) is seen at  $\delta$  3.73 and is assigned to H- $7_{eq}$  with the long range coupling identified as a "W"-type coupling with H- $5_{eq}$ . A number of other signals not seen in spectra of other pumiliotoxin class alkaloids, were seen in this spectrum that strongly implies a rigid side-chain moiety, *viz.* discrete dd or ddd signals. These are assigned as indicated in Table 2. In particular, the chemical shifts and multiplicities of signals assigned to the H- $10\alpha$ , H- $10\beta$ , H-11, H- $12\alpha$ , H- $12\beta$  and H-13 protons all indicate lack of free rotation. For example, the H-10 and H-12 signals show  $\Delta\delta$  separations of 0.5 and 1.05 ppm, respectively. The close correspondence of measured  $J$  values seen in the pyran signals with the dihedral angles calculated for a slightly distorted chair conformation are consistent with their being attached to this rigid six-membered ring. Included in Table 2 are selected dihedral angles calculated with an MM2 computer

program for **341A**. The skewed chair conformation resulted when one dihedral angle ( $\theta_{13-12\alpha}$ ) was fixed at  $40^\circ$  to agree with the  $J$  of 7 Hz observed for the coupling between H-13 and H-12 $\alpha$ . When this was done and the energy minimized, the other dihedral angles in the pyran ring agreed well with the observed coupling constants.

Table 2.  $^1\text{H-NMR}$  data on alkaloid **341A** (**1**) (500 MHz,  $\text{C}_6\text{D}_6$ )

H-Position	$\delta$ (ppm)	multiplicity, $J$ (Hz)	$\theta$ , Calc'd ( $^\circ$ ) <sup>a</sup>
1	1.75	m	
1 <sup>b</sup>	1.45	m	
2	1.75	m	
2'	1.45	m	
3ax.	2.04	q, 8.8	$\theta_{3ax-2\alpha}$ (141), $\theta_{3ax-2\beta}$ (21)
3eq.	2.72	td, 8.2, 1.8	$\theta_{3eq-2\beta}$ (99), $\theta_{3eq-2\alpha}$ (20)
5ax.	2.24	dd, 10.6, 1.5	
5eq.	2.63	br d; 10.6	
7eq.	3.73	d, ca. 1	
8a	2.58	dd, 9.7, 7.1	$\theta_{8a-2\alpha}$ (161), $\theta_{8a-2\beta}$ (40)
10 $\alpha$	1.35	overlaps H-16	$\theta_{10\alpha-11}$ (65)
10 $\beta$	1.85	dm, ca. 13, m	$\theta_{10\beta-11}$ (47)
11	1.60	m (14 lines)	$\theta_{11-12\alpha}$ (75), $\theta_{11-12\beta}$ (42)
12 $\alpha$	1.80	m	$\theta_{12\alpha-13}$ (40), $\theta_{12\alpha-11}$ (75)
12 $\beta$	0.75	ddd, 12.8, 10.3, 7.1	$\theta_{12\beta-13}$ (162), $\theta_{12\beta-11}$ (42)
13	3.28	dd, 9.9, 7.1	$\theta_{13-12\beta}$ (162), $\theta_{13-12\alpha}$ (40)
15, 15'	1.55	m	
16	1.30	m, overlaps H-10 $\alpha$	
16'	1.17	m	
CH <sub>3</sub> -9	1.55	s	
CH <sub>3</sub> -17	0.86	t, 7.2	
CH <sub>3</sub> -18	0.79	d, 6.6	
CH <sub>3</sub> -19	0.92	s	

<sup>a</sup> Indicated dihedral angles are calculated with the Chem3D program (Cambridge Scientific Computing) for the Macintosh computer.

<sup>b</sup> Primed hydrogens are upfield of unprimed and have no stereochemical significance.

The pyran chair conformation places the C-18 methyl group ( $\delta$  0.80; d, 6.6 Hz) in an axial orientation. Note that H-10 $\beta$  shows only one large, geminal, coupling and therefore cannot have any *trans* diaxial couplings. The equatorially disposed pentyloxy side-chain and H-12 $\beta$  are indicated as in *cis* orientation from the highly shielded environment and the couplings observed for H-12 $\beta$ , namely a ddd signal: one doublet,  $J=12.8$  Hz is merely a geminal coupling; the other doublets,  $J=10.3$  and 7.1 Hz, are assigned to couplings with H-13 and H-11, respectively. The former assignment is secured by noting that the 10.3

Hz coupling is one of the couplings observed in the dd signal of H-13 and is consistent with the calculated  $\theta_{12\beta-13}$  of  $162^\circ$ . The very unusual upfield position for H-12 $\beta$  ( $\delta$  0.75) is a consequence of shielding by the C-13 substituent; this significant  $\Delta\delta$  cannot be the effect alone of the ether-oxygen lone-pair anisotropy since there is no similarly significant shielding effect observed on H-10 $\beta$ .

The *R* configuration of C-11, is chosen as probable, since that configuration is present in all pumilio-toxins whose structures have been proven either by X-Ray analysis<sup>5</sup> or synthesis.<sup>7</sup> Unfortunately, the complex multiplicity of the H-11 signal prevents extracting the exact couplings between H-11 and the vicinal protons.

The 2D <sup>1</sup>H-NMR spectra (normal and relay) revealed the positions of all the hydrogens and established all connectivities (e.g. C-10-11-12-13; C-8a-1-2-3; C-15-16-17). Chemical shifts, and where possible, their couplings are indicated in Table 2. In 2D spectra (relay 0 and 1), a weak long-range coupling between H-10 $\alpha$  and H-5 $_{eq}$  and a much stronger one ("W"-type) between H-10 $\alpha$  and H-5 $_{ax}$ , are observed; a weak cross-peak between H-10 $\beta$  and H-5 $_{ax}$  is also observed. These also support a fixed orientation of the H-10 hydrogens relative to the indolizidine ring. Strong four-bond couplings are seen between CH<sub>3</sub>-18 and the H-10 $\alpha$ , 10 $\beta$  and H-12 $\alpha$  protons (H-12 $\beta$  and CH<sub>3</sub>-18 practically overlap, so a cross-peak is not seen). Weak long-range couplings are also seen between CH<sub>3</sub>-19 and H-13 and one or both H-15 protons. The relative stereochemistry at C-13 in **1** is shown as 13*S* (based on C-11 being in the *R*-configuration) with C-14 unassigned.

Certain microchemical procedures provided further relevant data. Hydrogenation did not affect **341A**. Acetylation in the presence of pyridine after 16 h at  $58^\circ$  gave a 90% conversion to a monoacetate with 10% recovered starting material. Little reaction was observed at  $65-70^\circ$  for 4.5 h, conditions that resulted in the complete acetylation of **251H** (**7**). The EI-MS spectrum indicated acetylation had occurred at the 7 $\beta$ -hydroxyl group since the *m/z* 254 ion is missing and is replaced by a fragment at *m/z* 296 (5%). A two-day reaction at  $70-75^\circ$  did give a mixture of mono and diacetates; the second acetyl group is most probably located at the 14-hydroxyl group. Interestingly, both **5** and **6** easily formed triacetates indicating the 7-hydroxyl group in those compounds reacts in either the  $\alpha$  or  $\beta$  configuration. Treatment of **341A** with butylboronic acid gave no reaction indicating that no *cis*-diol grouping was present. Under these conditions, synthetic **6**, gave a *bis*-butylboronate.<sup>8</sup>

Alkaloid **341A** occurs in frog skin extracts relatively rarely. Isolated from skin extracts from one population of the Ecuadoran *Epipedobates tricolor*, it has also been detected in certain Central American populations of *Dendrobates auratus*, *D. pumilio* and *D. granuliferus* and in certain populations of Colombian *D. lehmanni*, *Minyobates minutus* and *M. viridis*.<sup>1</sup> Alkaloid **341B**, detected along with **341A** in skin extracts from Colombian *D. lehmanni* and certain populations of Costa Rican *D. pumilio* (ref. 1

and unpublished results), appears to be a diastereomer of **341A**. The mass spectrum is somewhat different from that of **341A**.<sup>2</sup> Alkaloid **357**, detected in *Epipedobates tricolor*, is evidently a hydroxy analog of **341A**.

## EXPERIMENTAL

**Instrumentation:** A Hewlett-Packard model 5890 gas chromatograph having a 25 m  $\times$  0.32 mm i.d. HP-5 fused silica-bonded capillary column programmed from 100° to 280° at the rate of 10°/min, interfaced with a Hewlett-Packard model 5971 Mass Selective Detector and a Hewlett-Packard model 5965B IR instrument with a narrow band (4000-750 cm<sup>-1</sup>) detector and a Hewlett-Packard ChemStation (DOS based) were used to generate the chromatograms, EI-MS, and FTIR spectra of **341A** (instrument A). Modified EI-MS (ion trap) of GC peaks were generated with a Finnigan Model 800 Ion Trap Mass Detector interfaced with a Varian Model 3400 gas chromatograph fitted with a 30 m  $\times$  0.32 mm i.d. RTX-5 (Restek) fused silica-bonded column and using the same temperature program as above (instrument B). A Finnigan 4500 mass spectrometer with a 25 m  $\times$  0.25 mm i.d. OV-17 fused silica-bonded column (Supelco) with a 60°-280° program (10°/min or 5°/min) and an INCOS data system was also used for EI-MS (instrument C). HRMS were measured with a JEOL SX 102 instrument fitted with a 15 m  $\times$  0.20 mm i.d. HP-5 column. All HRMS measurements were within  $\pm$  5.6 ppm except that for the *m/z* 298 ion. Chemical ionization used the ion-trap instrument and NH<sub>3</sub> or ND<sub>3</sub> reagent gases. The Finnigan 4500 was used with a D<sub>2</sub>O bleed to obtain deuterium-exchanged EI-MS. The 1D- or 2D-(COSY) <sup>1</sup>H-NMR spectra in D<sub>2</sub>O or C<sub>6</sub>D<sub>6</sub> were measured with either a Varian XL-300 or a Varian VXR-500S spectrometer. Chemical shifts ( $\delta$ , ppm) in D<sub>2</sub>O are referred to HOD at 4.78; those in C<sub>6</sub>D<sub>6</sub> to TMS at 0.0 ppm.

**Isolation of 341A:** The usual acid/base partitioning procedure of methanolic extracts of skins of 750 *Epipedobates tricolor* frogs collected 16 km west of Santa Isabel, Azuay province in southwestern Ecuador<sup>9</sup> yielded 60 mg of an alkaloid mixture.<sup>5</sup> The alkaloid mixture was chromatographed on a Merck prepacked silica gel 60 (1.0  $\times$  24 cm) column with 500 mL of CHCl<sub>3</sub>:CH<sub>3</sub>OH:6 N NH<sub>3</sub> (800:10:0.1) followed by 1.0 L of 100:10:0.2 and collecting 5 mL fractions. Compound **341A** (ca. 9 mg) appeared in fractions 13-17, but these fractions also contained substantial amounts of pumiliotoxin **251D** and minor or trace amounts of other alkaloids, including quinolizidine **207I**, indolizidine **207A**, deoxypumiliotoxin **251H** (2 isomers), pumiliotoxins **265G**, **267C** (2 isomers) and **307A**, allopumiliotoxin **323B** and epibatidines **208/210** and **308/310**, as well as minor amounts of pumiliotoxin **357**, evidently a hydroxy congener of **341A** (EI-MS data below). The fractions containing **341A** were concentrated, dissolved in



methanol and purified using reverse phase HPLC with a C-18 column and a flow rate of 1 mL/min and the solvent system HOAc:CH<sub>3</sub>CN:H<sub>2</sub>O. After isocratic conditions (2:6:92) for the first 5 min, a gradient from 2:6:92 to 2:68:30 over 20 min was employed. Thirty fractions (1 mL each) were collected and analyzed by GC-MS after evaporation and redissolution in methanol. Additional purification of **341A**-containing fractions was performed with an Asahipack ODP-50 column (available from Hewlett-Packard; 4.6 mm x 25 cm) and the solvent system CH<sub>3</sub>CN:H<sub>2</sub>O with a gradient of 10:90 to 90:10 over 30 min and a flow rate of 0.5 mL/min. Thirty fractions (0.5 mL each) were collected with pure **341A** appearing in fractions 10 and 11 (ca. 1 mg total as an oil).

Characterization of **341A**:

EI-MS: Instrument A: 341(6), 323(10, probably a thermal loss of H<sub>2</sub>O), 306(2), 298(16), 266(9), 254(19), 236(3), 198(3), 184(7), 182(6), 180(5), 166(3), 164(3), 126(16), 125(15), 114(28), 112(100), 97(14), 96(11), 87(13), 84(61), 70(98), 55(18). Instrument B (Ion trap pseudo EI-MS): 342(14), 323(7), 306(4), 298(5), 254(7), 184(5), 182(3), 126(15), 125(15), 114(23), 112(40), 84(79), 70(100). Instrument C: 341(8), 323(8), 306(1), 298(10), 266(7), 254(13), 184(8), 182(8), 180(4), 126(12), 125(8), 114(16), 112(76), 97(8), 87(14), 84(50), 82(30), 70(100). The first EI-MS of **341A** was obtained with an LKB 300 spectrometer with a 1.5% OV-1 packed column (10). The EI-MS was as follows: 341(4), 324(3), 323(1), 306(1), 298(3), 266(4), 254(7), 114(10), 112(60), 84(42), 70(100).

IR: (see Figure 1) 3569(23), 3536(13), 2965(100), 2943(77), 2883(49), 2818(33), 1459(22), 1379(33), 1313(27), 1300(24), 1226(17), 1171(40), 1070(65), 821(12) cm<sup>-1</sup>.

Optical rotation was not measured.

Mono-O-acetate of **341A**; EI-MS: M<sup>+</sup> not detected, 340(3), 324(4), 306(3), 296(5), 280(4), 266(3), 236(3), 217(3), 166(3), 126(5), 102(7), 83(100), 70(34), 55(10).

Characterization of **357**:

EI-MS: 357(6), 339(6), 322(2), 314(8), 270(4), 184(6), 182(6), 128(12), 126(8), 125(8), 114(15), 110(74), 100(10), 96(8), 95(8), 87(16), 84(48), 83(28), 70(100), 55(16).

IR: 3684(5), 3657(5), 3575(28), 3536(13), 2972(100), 2942(83), 2885(50), 2816(42), 1458(25), 1380(42), 1307(35), 1225(28), 1173(52), 1071(87), 951(27), 824(13) cm<sup>-1</sup>.

## ACKNOWLEDGMENT

We thank Drs. L. K. Pannell and H. J. C. Yeh (NIH) for HRMS and the 2D <sup>1</sup>H-NMR data, respectively.

## REFERENCES AND NOTES

1. J. W. Daly and T. F. Spande, 'Alkaloids: Chemical and Biological Perspectives' Vol. 4, ed. by S. W. Pelletier, John Wiley and Sons, New York, 1986, p. 1.
2. J. W. Daly, H. M. Garraffo, and T. F. Spande, 'The Alkaloids', Vol. 43, ed. by G. A. Cordell, Acad. Press, San Diego, 1993, p. 185.
3. J. W. Daly, H. M. Garraffo, and T. F. Spande, 'Alkaloids: Chemical and Biological Perspectives' Vol. 13, ed. by S. W. Pelletier, Elsevier Science Ltd., Oxford, 1999, p. 1.
4. J. W. Daly, E. T. McNeal, L. E. Overman, and D. H. Ellison, *J. Med. Chem.*, 1985, **28**, 482; J. W. Daly, F. Gusovsky, E. T. McNeal S. Secunda, M. Bell, C. R. Creveling, Y. Nishizawa, L. E. Overman, M. J. Sharp, and D. P. Rossignol, *Biochem. Pharmacol.*, 1990, **40**, 315.
5. J. W. Daly, T. Tokuyama, T. Fujiwara, R. J. Highet, and I. L. Karle, *J. Am. Chem. Soc.*, 1980, **102**, 830.
6. P. Jain, H. M. Garraffo, T. F. Spande, H. J. C. Yeh, and J. W. Daly, *J. Nat. Prod.*, 1995, **58**, 100.
7. L. E. Overman and L. K. Bell, *J. Am. Chem. Soc.*, 1981, **103**, 1851; L. E. Overman, L. K. Bell, and F. Ito, *J. Am. Chem. Soc.*, 1984, **106**, 4192; L. E. Overman and S. W. Goldstein, *J. Am. Chem. Soc.*, 1984, **106**, 5360; L. E. Overman and N-H. Lin, *J. Org. Chem.*, 1985, **50**, 3669; A. S. Franklin and L. E. Overman, *Chem. Rev.*, 1996, **96**, 505.
8. B. M. Trost and T. S. Scanlan, *J. Am. Chem. Soc.*, 1989, **111**, 4988.
9. Voucher specimens at the American Museum of Natural History, New York City.
10. J. W. Daly, C. W. Myers, and N. Whittaker, *Toxicon*, 1987, **25**, 1023.

Received, 17th August, 1998