

REACTIONS OF PYRIDINE N-OXIDE WITH ENAMINES OF N-SUBSTITUTED
4-PIPERIDONES IN THE PRESENCE OF AN ACYLATING AGENT

Michio Nakanishi and Masahiro Yatabe

Research Laboratories, Yoshitomi Pharmaceutical Industries Ltd.,

Koiwai, Yoshitomi-cho, Chikujyo-gun, Fukuoka, Japan

Masatomo Hamana*

Faculty of Pharmaceutical Sciences, Kyushu University

Maidashi, Higashi-ku, Fukuoka, Japan

Pyridine N-oxide reacts with enamines of N-benzoyl-, N-ethoxycarbonyl- and N-acetyl-4-piperidones in the presence of benzoyl chloride to give N-substituted 3-(2-pyridyl)-4-piperidones in fair or good yields. Enamine of N-methyl- or N-benzyl-4-piperidone resists this reaction.

Pyridine N-oxide (I) readily reacts with enamines of cyclohexanone in the presence of benzoyl chloride to give 2-(2-pyridyl)-cyclohexanone on treatment of the reactants with 20% hydrochloric acid¹. While the reaction using 1(10)-dehydroquinolizidine as a heterocyclic enamine similarly proceeds², attempted reactions with enamines of N-substituted 4-piperidones such as morpholine enamine of N-benzoyl-4-piperidone

(A) were described to be unsuccessful³.

Recently we happened to find that a small amount of picolinic acid N-oxide was obtained from the reaction of I with A by oxidation of the residue from the 20% hydrochloric acid extract with 30% hydrogen peroxide and acetic acid. This fact stimulated to re-examine this reaction, and we succeeded in the isolation of N-benzoyl-3-(2-pyridyl)-4-piperidone (II) on treatment of the reaction mixture with conc. hydrochloric acid instead of the generally used 20% hydrochloric acid.


Thus, when benzoyl chloride (1.2 equiv) was added to an ice-cooled solution of I and A (3 equiv) in chloroform, an exothermic reaction occurred and the solution became dark red through yellow. The reaction mixture was kept at room temperature for 2 days, followed by extracting with conc. hydrochloric acid to give II, light yellow powders, mp 114-116° (isopropyl alcohol-isopropyl ether), as a main product in 26% yield.

Structure assignment of II is based on the satisfactory elemental analysis [C₁₇H₁₆O₂N₂], the ir spectrum [$\nu_{\text{max}}^{\text{KBr}}$: 2600 (a chelated hydrogen bond) and 1630 cm⁻¹ (an enol C=C bond)] and nmr spectrum [δ (CDCl₃): 2.48 (2H, t, J=6.0 Hz, C₅-H of piperidone ring), 3.68 (2H, t, J=6.0 Hz, C₆-H of piperidone ring), 4.33 (2H, s, C₂-H of piperidone ring), 8.30 (1H, m, α -H of pyridine ring) and 15.5 (1H, s, OH)]; apparently II exists chiefly as the enolic form (IIa) rather than the ketonic (IIb) and the enaminoic ones (IIc) in the same way with other 2-picolyl ketones⁴. Oxidation of II with 30% hydrogen peroxide-

acetic acid gave picolinic acid N-oxide (III).

The reaction can be explained by the addition-elimination process of the benzoyl-adduct of I as in the case reported earlier¹, and the reported failure³ in isolating the product II may be due to the sparing solubility of II in 20% hydrochloric acid.

It was further found that the use of morpholine enamine of N-ethoxycarbonyl-4-piperidone (B) instead of A gave the corre-



sponding product (IV), yellowish oil, bp 158-160° (0.3-0.4 mm Hg), in much better yield (85%).

Treatment of V with sodium borohydride in ethanol afforded two isomeric alcohols (V and VI); V was isolated as a crystalline hydrochloride, mp 181-184°, and VI as an oxalate, mp 140°.

Some detailed examinations of reactions with enamines of N-ethoxycarbonyl-4-piperidone revealed that there were no noticeable differences among reactions using 4-morpholino-,

4-piperidino- and 4-pyrrolidino-derivatives and the order in effectiveness of acylating agents were as follows : benzoyl chloride > tosyl chloride > acetyl chloride > acetic anhydride (see Table).

Much more remarkable is the large dependency of the ease with which the reaction occurs on the nature of the N-substituent of the piperidone; thus, the reaction of I with morpholine enamine of N-acetyl-4-piperidone (C) smoothly proceeded in the presence of benzoyl chloride to give the N-acetyl derivative (VII), bp 120-130° (bath temp.) (0.1-0.2 mm Hg), in 51.4% yield, but no definite product was obtained from the similar reaction with enamines of N-methyl- or N-benzyl-4-piperidone.

Table I

Exp. No.	(I) g	Amine of Enamine ^{a)} g (eq)	AX g (eq)	Et ₃ N g (eq)	React. time	Product (IV) g (%)
1	4.75	morpholine 16.8 (1.4)	PhCOCl 8.4 (1.2)	-	48hr	10.6 (85)
2	4.75	morpholine 16.8 (1.4)	TsCl 11.4 (1.2)	-	48hr	4.7 (37.9)
3	4.75	morpholine 16.8 (1.4)	AcCl 4.7 (1.2)	-	48hr	1.9 (15.3)
4	4.75	morpholine 16.8 (1.4)	Ac ₂ O 6.1 (1.2)	-	48hr	0.4 (3.3)
5	4.75	piperidine 16.7 (1.4)	PhCOCl 8.4 (1.2)	-	48hr	9.8 (79)
6	4.75	pyrrolidine 15.7 (1.4)	PhCOCl 8.4 (1.2)	-	48hr	9.9 (79.8)
7	4.75	morpholine 12.0 (1.0)	PhCOCl 7.0 (1.0)	-	48hr	7.7 (62)
8	4.75	morpholine 24.0 (2.0)	PhCOCl 7.0 (1.0)	-	48hr	10.1 (81.5)
9	4.75	morpholine 12.0 (1.0)	PhCOCl 7.0 (1.0)	5.06 (1.0)	48hr	5.7 (46)

a) Enamine of N-ethoxycarbonyl-4-piperidone

Enamine B is most reactive towards this type of reaction among enamines of N-substituted 4-piperidone so far examined and reacts with various derivatives of pyridine and other aromatic N-oxides. The details of this study will be published shortly.

REFERENCES

- 1 M. Hamana and H. Noda, Chem. and Pharm. Bull. (Japan), 1965, 13, 912.
- 2 S. Saeki, A. Yamashita, Y. Matsukura, and M. Hamana, Chem. and Pharm. Bull. (Japan), 1974, 22, 2341.
- 3 M. Hamana, "The Chemistry of Heterocycles", Vol.3, ed. by Y. Kitahara, T. Kametani, and T. Kato, Nankodo Co., Ltd., Tokyo and Kyoto, pp. 59-60, 1971.
- 4 R. F. Branch, A. H. Beckett, and D. B. Cowell, Tetrahedron, 1963, 19, 401; G. Klose, and E. Uhlemann, ibid., 1966, 22, 1373.

Received, 17th January, 1975