

STUDIES ON HOMOPIPERAZINE DERIVATIVES: CONFORMATIONAL
ANALYSIS OF 2-(4-ACYLHOMOPIPERAZIN-1-YL)-4-AMINO-6,7-
DIMETHOXYQUINAZOLINES BY NMR

Hachiro Sugimoto, Sachi-yuki Hamano, Tohru Takahashi, Takeshi Suzuki,
=====

Fukuko Kuriyagawa, and Kenya Sakaguchi
=====

Eisai Research Laboratories, Eisai Co. Ltd.


Koishikawa 4, Bunkyo-ku, Tokyo 112, Japan

During the course of synthetic studies of the title compounds, we observed unusual behaviors in the n.m.r. spectrum of (1), in which the signals of the terminal CH₃ protons at γ-position appeared as two sets of triplet and one of the aromatic protons showed two different chemical shifts.

Spectroscopic analysis suggested presence of intramolecular interaction in organic solvent such as d₆-DMSO and intermolecular interaction in D₂O. These intra- and intermolecular interactions might lead to stabilization of specific, at least two, conformations formed through some electrostatic forces between the lone pair electrons of nitrogen at position 1 and the positively charged carbon at the amide function.

The above mentioned unique n.m.r. informations may be explained by the presence of such conformers in the compound.

The conformational structures of (1) and its analogues are proposed.



(1) -R: -CH₂CH₂CH₃
 α β γ