

REACTION OF 4-AZIDO-5-(1-CYANOETHYLIDINE)-1, 3-DIMETHYL-2-PYR-
 ROLINONE WITH AMINES. A CONVENIENT SYNTHESIS OF 3-AMINOMETHYL-
 4-AMINO-5-(1-CYANOETHYLIDINE)-1-METHYL-2-PYRROLINONES

Annie Sing and Harold W. Moore *

Department of Chemistry
 University of California
 Irvine, California 92717


The reaction of 4-azido-5-(1-cyanoethylidene)-
 1, 3-dimethyl-2-pyrrolinone with amines (piperidine,
 morpholine, pyrrolidine, and aniline) is described.

This reaction results in the incorporation of an amino-
 methyl group at position -3 with concomitant reduction
 of the azide group to a primary amine, *i. e.*, the forma-
 tion of 3-aminomethyl-4-amino-5-(1-cyanoethylidene)-
 2-pyrrolinones, 6a-d.

In conjunction with a general synthetic objective of developing new routes
 to highly substituted 2-pyrrolinones, we report here a unique reaction of
 4-azido-5-(1-cyanoethylidene)-1, 3-dimethyl-2-pyrrolinone (3) with amines.
 This reaction results in the incorporation of an aminomethyl group at position
 -3 with concomitant reduction of the azide moiety to a primary amine, *i. e.*,
 the formation of 3-aminomethyl-4-amino-5-(1-cyanoethylidene)-1-methyl-2-

pyrrolinone (6a-d). This work was stimulated by the recent report of Yoshina and Kuo¹ who reported an analogous reaction of 7-azido-6-methyl-5,8-quinolindione with secondary amines.

The azidopyrrolinone, 3, was readily prepared from β -azido- γ -cyanoethylidene- α -methyl- $\Delta^{\alpha, \beta}$ -butenolide 1 upon treatment with methyl amine to give 4-azido-5-(1-cyanoethyl)-5-hydroxy-1,3-dimethyl-2-pyrrolinone (2): 79%; mp, 128-130 °C (decomp); ir (nujol, cm^{-1}) 3100, 2240, 2105, 1690; pmr (CDCl_3 , δ) 0.97 d(3) $J = 7$ Hz, 1.77 s(3), 2.67 s(3), 3.55 q (1) $J = 7$ Hz, 7.38 s(1)(acidic). Anal. Calcd. for $\text{C}_9\text{H}_{11}\text{N}_5\text{O}_2$: C, 48.87; H, 5.01; N, 31.66. Found: C, 48.91; H, 5.56; N, 30.97. Mass: M.W. = 221; observed $M^+ - 28(\text{N}_2) = 193$. Subsequent dehydration of 2 gave the desired azidopyrrolinone, 3, in 53% yield; mp, 100 - 102 °C (decomp); ir (nujol, cm^{-1}) 2200, 2090, 1700, 1605; pmr (CDCl_3 , δ) 2.12 s(3), 2.33 s(3), 3.42 s(3); Anal. Calcd for $\text{C}_9\text{H}_9\text{N}_5\text{O}$: C, 53.20; H, 4.46; N, 34.46; Found: C, 52.44; H, 4.45; N, 34.70. In view of the fact that azidobutenolides such as 1 are readily available from the acid catalyzed rearrangement of 2,5-diazido-1,4-benzoquinones^{2,3} and that they can be easily converted to azidopyrrolinones, as outlined here, one can envisage a variety of starting materials of structural type 3.



Conversion of 3 to the corresponding 3-aminomethyl derivatives, 6a-d, was accomplished by slowly adding the azide 3 to a hot chlorobenzene solution ($\sim 120^\circ\text{C}$) containing excess of the amine. The resulting Mannich base-type products, 6a-d, were then isolated in $\sim 50\%$ yield by recrystallization and/or chromatography on silica gel, and their structures were determined by the spectral properties as provided in Table I. Although the mechanism of this reaction has not been determined, a nitrene intermediate, 4, is proposed since 4-amino-5-(1-cyanoethylidene)-1,3-dimethyl-2-pyrrolinone was usually detected as a minor product of the reaction.









TABLE I

Physical and Spectral Properties

		Yield	m. p.	I.R. (nujol, cm ⁻¹)	pmr (δ ppm)	C ¹³ mr (δ ppm)	Analysis	Color
6a		49%	214° - 215.5°C	3375, 3305 (shoulder) 2190, 1715 1665, 1605	in CDCl ₃ 1.50 (6H, broad, piperidine) 1.67 (3H, s, CH ₃) 2.50 (4H, broad, piperidine) 3.30 (2H, s, CH ₂) 3.38 (3H, s, NCH ₃) 6.33 (2H, broad, NH ₂)	in CDCl ₃ 6.61, 24.39, 26.07 28.07, 53.32, 59.09 83.59, 100.05, 120.91, 150.73, 151.99, 172.31	Calcd: C, 64.59; H, 7.74; N, 21.52 Found: C, 64.69; H, 7.82; N, 21.39	bright yellow
6b		50%	198° - 199 °C	3380, 3330 2190, 1705 1665, 1595	in CDCl ₃ 1.75 (3H, s, CH ₃) 2.47-2.73 (4H, m, morphine) 3.41 (2H, s, CH ₂) 3.42 (3H, s, NCH ₃) 3.57-3.83 (4H, m, morphine) 6.02 (2H, broad, NH ₂)	in CDCl ₃ 6.21, 27.58, 52.02 58.28, 66.49, 81.87 100.38, 120.18, 149.70, 151.81, 171.61	Calcd: C, 59.53; H, 6.92; N, 21.36 Found: C, 59.66, H, 7.06; N, 21.30	bright yellow
6c		15%	201° - 202°C	3340 (shoulder), 3280 2195, 1690, 1655 1640, 1630, 1590	in CDCl ₃ 1.40-2.13 (4H, m, pyr- rolidine) 1.70 (3H, s, CH ₃) 2.20-3.00 (4H, m, pyr- rolidine) 3.38 (3H, s, NCH ₃) 3.47 (2H, s, CH ₂) 6.32 (2H, broad, NH ₂)	in CDCl ₃ 6.29, 23.56, 27.71 52.64, 55.72, 83.97 99.94, 120.45, 150.35, 151.37, 171.99	Calcd: C, 63.39; H, 7.37; N, 22.75 Found: C, 63.47; H, 7.43; N, 22.95	bright yellow
6d		59%	196.5° 198 °C	3330, 3120, 2195 1700, 1660, 1600	in acetone - d ₆ 1.68 (3H, s, CH ₃) 3.30 (3H, s, NCH ₃) 4.17 (2H, std, CH ₂) 5.52 (1H, broad, NH) 6.03 (2H, broad, NH ₂) 6.63 - 7.47 (5H, m, aromatic)	in DMSO - d ₆ 6.46, 27.65, 43.16 86.98, 98.74, 113.18 117.69, 119.28, 129.05, 147.50, 149.09, 150.28, 171.11	Calcd: C, 67.15; H, 6.01; N, 20.88 Found: C, 67.20; H, 6.08; N, 20.82	bright yellow

REFERENCES

1. S. Yoshina and H. S. Kuo, Heterocycles, 4, 1337 (1976).
2. H. W. Moore, P. Germeraad, and W. Weyler, Jr., J. Org. Chem., 38, 3865 (1973).
3. H. W. Moore, H. R. Shelden, D. W. Deters, and R. J. Wikholm, J. Am. Chem. Soc., 92, 1675 (1970).
4. The authors wish to thank the National Science Foundation (CHE-06932) for financial support of this work.

Received, 27th December, 1977