


THE TOTAL SYNTHESIS OF A MITOSENE

J. Rebek, Jr.* and S. H. Shaber


Department of Chemistry, University of Pittsburgh, Pittsburgh,
Pennsylvania 15260

ABSTRACT - The conversion of L-hydroxyproline to a mitosene (degradation product of a mitomycin) is described. The synthesis involves 19 steps and the key stereochemical feature arises from an unusual *cis* opening of an epoxide. The synthetic product, 1-methoxy-2,7-diamino-mitosene is shown to be identical with the *trans* isomer obtained from acid-catalyzed methanolysis of mitomycin C.


Mitosenes comprise a family of substances obtained from the chemical degradation of the mitomycin natural products.¹ The fragile structural elements of the mitomycins are absent in the mitosenes, yet some of the biological activity remains. Accordingly, the mitosenes, rather than the mitomycins, have been the objects of most synthetic schemes, yet of the dozen or so published approaches², none has yielded a mitosene obtainable from a natural product. Only Kishi's³ total synthesis of the mitomycins can be said to have increased the availability of the simpler mitosenes. Our own efforts have now produced mitosenes 1, and while it is unlikely that our methods can be adapted to the synthesis of the parent mitomycin, the prompt access described below may compensate for the modesty of the goal.




The tricyclic ketone **3** was prepared from the L-hydroxyproline derivative **2** by way of Huisgen pyrrole synthesis⁴ then Dieckmann cyclization as previously described.⁵ Saponification, mesylation then methoxide-induced elimination gave the racemic olefin **4**, mp 223-225°, from which the *trans* bromohydrin **5** was obtained as a mixture of diastereomers.⁶



The regiochemistry of the bromohydrin was determined to be as shown⁷; the intermediate 1,2-bromonium ion is therefore opened exclusively at C₁. Treatment of **5** with a number of nucleophiles under basic conditions gave products with regiochemistry consistent with the intermediacy of an elusive 1,2-epoxide which, like the bromonium ion, was attacked only at C₁. The regiochemistry here has its parallel in the hydrolysis of the parent mitomycins - only 2-amino mitosenes are isolated from the opening of the aziridine ring. Moreover, this latter reaction has frequently been observed to result in *cis* opening of aziridine in preference to *trans* opening.⁹ Whatever the cause¹⁰, this phenomenon has its happy counterpart in the stereochemistry of products from **5**. In dilute methanolic methoxide **5** gave a 5:1 mixture of isomers in which the *cis* ether **6** was the major component.



Mesylation of 6 as before gave 7 and, since the free carboxylic acid could not be dehydrogenated directly, 7 was protected as its benzyl ester ($\text{C}_7\text{H}_7\text{Br}/\text{K}_2\text{CO}_3$ -18-crown-6, Me_2CO reflux, 16 h) 8 before treatment with DDQ. The product phenol 9 , mp 171 - 172° reacted rapidly with excess bromine water to yield the yellow bromoquinone 10 , mp 169 - 171° .



While the requisite oxidation state of the 6-membered ring was achieved in a single operation from the phenol, subsequent reductions in other parts of the molecule were attended by reduction of the quinone as well. Fortunately, the quinones were easily regenerated through oxidative workup. Thus removal of the ester through cautious hydrogenolysis¹¹ followed by FeCl_3 treatment gave the acid 11 , mp 179 - 180° (dec.). The acid chloride was prepared (SOCl_2) and, without isolation, was reduced by careful treatment with NaBH_4 (THF , 0° - 25° , 3 h). Quenching with phosphate buffer and addition of Fremy's salt then produced the orange alcohol 13 , mp 185 - 187° (dec.).

Transformation of this bromoquinone mesylate 13 into the amino quinone 14, mp 210-212° (N₂ evolves), was accomplished in a single operation by heating with excess NaN₃ in aqueous DMF. Initially the halide is displaced (30° h, ½ h) then reduction of the orange azidoquinone to the purple amino quinone with azide occurs (60°, 2 h), followed slowly by displacement of the mesylate (90°, 16 h). These conditions were developed with model compounds; for the case at hand the intermediates described were characterized by spectroscopic means upon interruption of the reaction at the appropriate time and temperature.

Conversion of the alcohol 14 to the carbamate 16 followed the well-trodden path *via* the carbonate 15, mp 155-157°. The final reduction of the azide to the mitosene was accomplished selectively in pyridine with (C₆H₅)₃P and aqueous ammonia.¹² Material obtained in this way was identical with the *trans* isomer 1a obtained from treatment of mitomycin C with acidic methanol.⁹ Further confirmation¹³ of the identity was obtained through acetylation to 1b. The synthesis was accomplished in 19 steps with an average yield of 80%/step (1% overall).

Acknowledgements - We are grateful to the National Institutes of Health for financial support and the Bristol Laboratories for a sample of mitomycin C.

REFERENCES

1. J. S. Webb, D. B. Cosulich, J. H. Mowat, J. B. Patrick, R. W. Broschard, W. E. Meyer, R. P. Williams, C. F. Wolf, W. Fulmor, C. Pidacks, J. E. Lancaster, J. Am. Chem. Soc., 1962, 84, 3185, 3187.
2. For synthetic approaches before 1977 see the review: T. Kametani, K. Takahashi, Heterocycles, 1978, 9, 293 and through 1979 see: K. Takahashi, T. Kametani, ibid., 1980, 13, 411. Recent synthetic approaches to the mitosenes include the following: R. M. Coates, C. W. Hutchins, J. Org. Chem., 1979, 44, 4743. T. Kametani, Y. Kigawa, H. Nemoto, M. Ihara, K. Fukumoto, Heterocycles, 1980, 14, 799. S. N. Falling, H. Rapoport, J. Org. Chem., 1980, 45, 1260. F. E. Chou, N. R. Bachur, R. Kupper, C. J. Michesda, Abstract 21 ORGN Section American Chemical Society Meeting Las Vegas, Nevada, Aug. 1980. J. Rebeck, Jr., S. H. Shaber, Heterocycles, 1981, 15, 161.

REFERENCES Con't.

3. F. Nakatsubo, A. J. Cocuzza, D. E. Keeley, Y. Kishi, J. Am. Chem. Soc., 1977, 99, 4835. F. Nakatsubo, T. Fukuyama, A. J. Cocuzza, Y. Kishi, ibid., 1977, 99, 8115. T. Fukuyama, F. Nakatsubo, A. J. Cocuzza, Y. Kishi, Tetrahedron Lett., 1977, 4295.
4. R. Huisgen, H. Gotthard, H. O. Bayer, F. C. Schafer, Chemische Berichte, 1970, 103, 2611.
5. F. M. Hershenson, J. Org. Chem., 1975, 40, 1260; J. Rebek, Jr., J.-C. Gehret, Tetrahedron Lett., 1977, 3027; F. M. Hershenson, J. Het. Chem., 1979, 16, 1093.
6. All compounds gave satisfactory elemental analyses, high resolution mass spectra and showed the expected spectroscopic features.
7. Proton nmr spectra of the *cis* 1,2 diol (derived from 4 by osmium-catalyzed hydroxylation) and the *trans*, 1,2 dibromide (derived from 4 by dibromination with Br₂ in CCl₄) and a *cis* 1-hydroxy-2-bromo derivative⁸ were used to assign the regiochemistry and stereochemistry of 5.
8. W. A. Remers, R. H. Roth and M. Weiss, J. Org. Chem., 1965, 30, 2910.
9. a) W. G. Taylor and W. A. Remers, Tetrahedron Lett., 1974, 3483; b) W. G. Taylor and W. A. Remers, J. Med. Chem., 1975, 18, 307.
10. L. Cheng and W. A. Remers, ibid., 1977, 20, 767. R. W. Franck, Progress in Chemistry of Organic Natural Products, 1979, 38, 1. W. A. Remers, "Chemistry of Antitumor Antibiotics", J. Wiley & Sons, Inc., New York, 1979, 221.
11. Prolonged treatment results in hydrogenolysis of the aryl bromide; best results were obtained using an equal weight of 10% Pd/C and 10 with stirring under H₂ for 30 min.
12. W. S. Mungall, G. L. Greene, G. A. Heavner, R. L. Letsinger, J. Org. Chem., 1975, 40, 1659.
13. NMR spectra at 300 MHz of *trans* 1b in CDCl₃/(CD₃)₂CO showed C₁-H δ4.63 (s, 1H); C₂-H δ4.91-4.98 (m, 1H), C₃-H_α δ4.19-4.24 (d, J_{3a-3b} = 13.7); C₃-H_β δ4.43-4.49 (dd, J = 13.5, J₂₋₃ = 5.9); For a discussion of stereochemical assignments at C₁ and C₂ in mitosenes see: U. Hornemann, P. J. Keller, J. F. Kozlowski, J. Am. Chem. Soc., 1979, 101, 7121 and ref. 9b.

Received, 6th April, 1981