



STUDIES IN THE DIHYDROPYRIDINE SERIES. V.[†] SYNTHESIS OF PYRIDOCARBAZOLE
ALKALOIDS: OLIVACINE AND GUATAMBUINE.

James P. Kutney*, Masaki Noda, Norman G. Lewis, Beatrice Monteiro, Danuta
Mostowicz, and Brian R. Worth

Department of Chemistry, The University of British Columbia, 2036 Main Mall,
University Campus, Vancouver, B.C., V6T 1Y6, Canada

Abstract - The tricarbonylchromium(0) complex 8 effectively stabilized the
very reactive dihydropyridine system during one carbon insertion. Release
of the complex allowed cyclization to the pyridocarbazole product 13 which
was elaborated to olivacine 1 and guatambuine 2.

The pyridocarbazole alkaloids, including olivacine 1 and guatambuine 2, have received considerable
interest as a result of their anti-tumour activity²⁻⁴. Numerous syntheses of representatives of
this family of alkaloids have been reported⁵⁻²⁰, and several routes employed cyclization of
pyridyl-substituted indoles to generate the linear pyridocarbazole skeleton although the yields
have generally been low. Our interest in the stabilization and use of very reactive dihydro-
pyridine systems^{1,21} led us to investigate the utility of tricarbonylchromium(0) complexes (eg. 8)
in the synthesis of natural products. Here, the complex 8 was intended to provide a nucleophilic
centre at the indole-3- position and a masked enamine (dihydropyridine) for the introduction of
a one carbon unit and subsequent cyclization to the pyridocarbazole skeleton.




Condensation of 4, readily derived from indole 3, with 4-acetylpyridine gave 5 which was readily
hydrolysed to 6^{18,22}. Reaction with iodomethane produced the salt 7 in high yield. Reduction of
7 with sodium borohydride and protection of the resulting dihydropyridine, in the usual manner²¹,
gave the tricarbonylchromium(0) complex 8 (mp 64-65^o; ν_{\max} 3420, 1941, 1866, 1821; λ_{\max} 218,
398) in 55% yield (Scheme 1).

[†] For Part IV, see reference 1.

The complex 8 reacted with the Vilsmeier salt from DMF and POCl₃ to give an intermediate (*viz.* 9) which on treatment with pyridine gave a mixture of yellow products (*eg.* 11; *viz.* 9 → 10 → 11). Subsequent reaction with sodium borohydride gave 1-desmethylguatambuine 12 (mp 274–276°; ν_{\max}


scheme 1.



3220, 1613; λ_{\max} 232, 238, 248, 259, 287, 297, 326, 340) in 89% yield from 8. Alternatively, dehydrogenation of the crude mixture (*eg.* 11) using Pd/C at 300° gave a mixture of the salt 13 (60% from 8) and the pyridine derivative 14 (10% from 8). The former was easily demethylated to 14 using triphenylphosphine in DMF or HMPA (Scheme 2). Methyl lithium, in dry THF, reacted with 14 and subsequent oxidation of the dihydropyridine intermediate with iodine gave olivacine²³ 1 in 54% yield. Reaction of 1 with iodomethane and reduction with sodium borohydride gave (\pm) guatambuine 2.

Chromium tricarbonyl complexing provided a good method for stabilizing very reactive dihydropyridines. Reaction with pyridine released the dihydropyridine from the complex, allowing cyclization and subsequent elaboration of the pyridocarbazole skeleton in a synthesis of olivacine and (\pm) guatambuine.

Acknowledgement: Financial aid from the Natural Sciences and Engineering Research Council of Canada is gratefully acknowledged.



References

1. J. P. Kutney, T. C. W. Mak, D. Mostowicz, J. Trotter, and B. R. Worth, *Heterocycles*, 1979, 12, 1517.
2. A. Ahond, A. Cave, T. A. Connors, L. K. Dalton, N. Dat-Xuong, M. Hayat, F. le Goffic, A. Gouyette, M. M. Janot, C. Kan-Fan, G. Mathe, J. le Men, J. Miet, J. Poisson, P. Potier, and T. Sevenet, *Biomedicine*, 1974, 21, 101.
3. E. M. Acta, O. P. Crews, L. Goodman, and C. W. Mosher, *J. Medicin. Chem.*, 1966, 9, 237.
4. N. Dat-Xuong, C. Goss, C. Paoletti, J. B. le Pecq, *Compt. Rend.*, 1973, 277, 2289.
5. J. Schmutz, and H. Wittwer, *Helv. Chim. Acta*, 1960, 43, 793.
6. M. Sainsbury, *Synthesis*, 1977, 437.
7. D. S. Grierson and J. P. Kutney, *Heterocycles*, 1975, 3, 171.
8. T. Kametani, Y. Ichikawa, T. Suzuki, and K. Fukumoto, *Heterocycles*, 1975, 3, 401; *J. Chem. Soc. Perkin 1*, 1975, 2102.
9. R. Besselièvre and H. P. Husson, *Tetrahedron Letters*, 1976, 22, 1873.
10. F. A. Hochstein, G. A. Iacobucci, and R. B. Woodward, *J. Amer. Chem. Soc.*, 1959, 81, 4434.
11. P. A. Cranwell and J. E. Saxton, *J. Chem. Soc.*, 1966, 3482.
12. T. R. Govindachari, S. Rajappa, and V. Sudarsanam, *Indian J. Chem.*, 1963, 1, 247.
13. R. N. Stillwell, Ph. D. Thesis, Harvard University, 1964.
14. L. K. Dalton, S. Demerac, B. C. Elmes, J. W. Loder, J. M. Swan, and T. Teitei, *Austral. J. Chem.*, 1967, 20, 2715.
15. K. N. Kilminster and M. Sainsbury, *J. Chem. Soc. Perkin 1*, 1972, 2264.
16. A. Ahond, F. le Goffic, and A. Gouyette, *Tetrahedron*, 1973, 29, 3357.

17. N. M. Hasan and A. P. Kozikowski, J. Org. Chem., 1977, 42, 2039.
18. R. Besselièvre, C. Thal, H. P. Husson, and P. Potier, J. Chem. Soc. Chem. Comm., 1975, 90.
19. S. J. Martinez and J. A. Joule, J. Chem. Soc. Chem. Comm., 1976, 818.
20. T. Naito, N. Iida, and I. Ninomiya, J. Chem. Soc. Chem. Comm., 1981, 44.
21. J. P. Kutney, R. A. Badger, W. R. Cullen, R. Greenhouse, M. Noda, V. E. Ridaura-Sanz, Y.H. So, A. Zanarotti, and B. R. Worth, Can. J. Chem., 1979, 57, 300.
22. H. F. Russell and R. J. Sundberg, J. Org. Chem., 1973, 38, 3324.
23. Synthetic samples of 1 and 2 were identical with authentic samples of the respective natural products. All compounds gave satisfactory spectral and analytical data.

Received, 6th May; 1981