

NEW IRIDOID GLUCOSIDES, CAMPENOSIDE AND 5-HYDROXYCAMPENOSIDE,
FROM *CAMPESIS CHINENSIS* VOSS

Shigeru Kobayashi*, Yasuhiro Imakura, and Yoshinobu Yamahara
Faculty of Pharmaceutical Sciences, Tokushima University

Shomachi, Tokushima 770, Japan

Tetsuro Shingu









School of Pharmacy, Kobe Gakuin University

Ikawadani, Tarumi-ku, Kobe 673, Japan

Abstract — Two novel iridoid glucosides, campenoside and 5-hydroxycampenoside were isolated from the leaves of *Campsis chinensis* Voss and their structures were shown to be $\underline{2}$ and $\underline{3}$, respectively.

Recently, we have isolated an alkaloid, boschniakine($\underline{1}$)¹ from the roots of *Campsis chinensis* Voss (Bignoniaceae). This isolation prompted us to investigate the iridoid glucosides of *C. chinensis* Voss. Chromatographic examination of a methanolic extract of the leaves of this plant gave two novel iridoid glucosides, campenoside ($\underline{2}$) and 5-hydroxycampenoside ($\underline{3}$), having a formyl group at C-4 and a cinnamoyl ester group at C-7. This paper describes structural studies of these compounds.

Campenoside ($\underline{2}$) [mp 165-167°, $C_{25}H_{30}O_{10} \cdot 1/2H_2O$, $[\alpha]_D^{20} - 59.5^\circ(\text{MeOH})$] gave D-glucose on acidic hydrolysis. The spectroscopic properties of $\underline{2}$ revealed the presence of a conjugated carbonyl group [$\delta_H(\text{CD}_3\text{OD})$: 9.19(s, CHO) and 7.39(s, 3-H); ν_{max} (KBr) 1670 and 1630 cm^{-1}] characteristic of iridoids, a *trans*-cinnamoyl ester group [δ_H 7.66 and 6.48(AB-system, J=16 Hz), 7.6-7.3(5H, aromatic H); ν_{max} 1720 and 1640 cm^{-1}], a methyl group [δ_H 1.11(d, J=6 Hz)], and an anomeric proton [δ_H 4.69(d, J=7 Hz)] of the β -linked D-glucose. Acetylation of $\underline{2}$ with Ac_2O -pyridine gave a tetraacetate($\underline{4}$) [mp 179-181°, $C_{33}H_{38}O_{14}$, $[\alpha]_D^{20} - 73.7^\circ(\text{CHCl}_3)$]. From the fact that the signals of $\underline{2}$ and $\underline{4}$ correspond closely to those of loganin pentaacetate ($\underline{5}$)² and from the coupling pattern of hydrogen atoms at 6 position of $\underline{2}$ (Table 1), the structures of campenoside and its tetraacetate were considered to be $\underline{2}$ and $\underline{4}$, respectively, although there is still some uncertainty about their stereochemistry. The above assumption was confirmed first by comparison of the ^{13}C -NMR spectrum of $\underline{2}$ with that of $\underline{5}$ ³ and then by chemical



transformation of 4 into 5.

As shown in Table II, the spectrum of 2 is similar to that of 5, the differences being accounted due to the different functions at C-4 and C-7: a methoxycarbonyl group and an acetate function in 5 and a formyl group and a *trans*-cinnamoyl ester group in 2.

The acetate (4) was converted to 5 as follows: oxidation of 4 with Jones reagent gave an acid (6) [mp 139-143°, C₃₃H₃₈O₁₅ (Mass spectrum, M⁺: Calcd: 674.2211. Found: 674.2227)]. The acid (6) was hydrolyzed with 1N NaOH-MeOH at room temperature to give *trans*-cinnamic acid and an acid (7), which was subjected to acetylation with Ac₂O-pyridine followed by methylation with diazomethane. The ester (5) [mp 131-134°, [α]_D²⁰ - 70.0° (CHCl₃)] thus obtained was found to be identical with authentic loganin pentaacetate² by comparison of spectral data and by mixed melting point.

5-Hydroxycampenoside (3) showed mp 176-179°, C₂₅H₃₀O₁₁, [α]_D²⁰ - 73.7° (MeOH), and ν_{max} (KBr) cm⁻¹: 3500(OH), 1710(conjugated ester), 1660(conjugated carbonyl group), 1640(C=C).

The presence of D-glucose and *trans*-cinnamic acid in 3 was confirmed by hydrolytic experiments. The ¹H-NMR spectrum of 3 gave signals which were assigned as shown in Table I, and additional signals due to β-D-glucose (δ_H 4.63, d, J=7.5 Hz) and *trans*-cinnamoyl protons.

Table I. $^1\text{H-NMR}$ Spectral Data of Iridoids (δ)^{a)}

Compd. (solvent)	1-H	3-H	5-H	7-H	9-H	10-H	11-H
2^b) (CD_3OD)	5.46(d,3)	7.39(s)	3.13(m)	5.24(m)	2.14(m)	1.11(d,6)	9.19(s)
3^c) (CD_3OD)	5.84(d,2)	7.40(s)		5.08(m)	2.46(dd,12,2)	1.13(d,7)	9.25(s)
4 (CDCl_3)	5.34(d,2)	7.06(s)	3.06(m)	5.27(m)	ca. 2.30(m)	1.09(d,7)	9.23(s)
5 (CDCl_3)	5.19(d,2)	7.33(s)	2.96(m)	5.13(m)		1.02(d,7)	

a) Numerical values in parentheses are coupling constants in Hz. b) 6-H α at 1.85(m, $J_{6\alpha-6\beta}=15$, $J_{6\alpha-5}=6.5$, $J_{6\alpha-7}=6$ Hz) and 6-H β at 2.32(m, $J_{6\beta-6\alpha}=15$, $J_{6\beta-5}=8$, $J_{6\beta-7}<1$ Hz). c) 6-H α at 2.67(dd, $J_{6\alpha-6\beta}=16$, $J_{6\alpha-7}=6$ Hz) and 6-H β at 2.27(dd, $J_{6\beta-6\alpha}=16$, $J_{6\beta-7}=2$ Hz).

Acetylation of 3 with AcO_2 -pyridine gave a tetraacetate (3^a) [mp 191-192°, $\text{C}_{33}\text{H}_{38}\text{O}_{15}$, $[\alpha]_D^{20}$ -90.7°(CHCl_3)], in which one hydroxyl group [ν_{max} (KBr) 3560 cm^{-1}] remained unaffected indicating its tertiary nature. From these data, 3 was concluded to have the same functional groups as 2 at the same positions, and an additional β -hydroxyl group at C-5. This was further confirmed by comparison of the $^{13}\text{C-NMR}$ spectrum of 3 with those of related compounds, as shown in Table II. The relation of the chemical shifts of geniposidic acid (9)^{4,5} and theviridoside (10)^{4,5} is very similar to that of 2 and 3 : (i) the signals for C-5 of 3 (69.39 ppm) and 10 (76.23 ppm) are shifted ca. 40 ppm downfield from those of 2 (28.55 ppm) and 9 (36.25 ppm); (ii) the signals for C-9 and C-6 in 3 are 8.55 and 7.21 ppm, respectively, downfield from the corresponding signals of 2 . This is probably due to β -oxygenation at C-5, since similar deshielding effects were observed by comparing the signals at C-9(9.77 ppm) and C-6(7.36 ppm) of 10 with the corresponding signals of 9 . (iii) The signal of C-7 of 3 is shifted 1.41 ppm upfield from that of 2 due to the γ -effect of 5 β -hydroxyl group in 3 . Similarly the signal of the corresponding carbon of 10 is shifted 1.60 ppm upfield from that of 9 . This assignment for the hydroxyl group at C-5 in 3 was also supported by comparison of δ_{H} of C-9 of 3 with that of 2 : the former signal is shifted 0.3 ppm downfield from the latter one, since the H-9 signals in the 5 β -hydroxy-iridoides, yuheinoside (11) (δ_{H} 2.70 ppm)⁶ and ipolamide (12) (δ_{H} 2.48 ppm)⁷, are 0.2-0.1 ppm downfield from those of the corresponding deoxy-compounds, boschnaloside (13) (δ_{H} 2.35 ppm)⁶ and mussaenoside (14) (δ_{H} 2.36 ppm)⁸, respectively. Additional evidence regarding the stereochemistry at C-7, C-8, and C-1 of 3 was obtained from its NMR data: the abnormal high-field shift^{3,4} of C-10(12.52 ppm) in 3 , as well as that(12.61 ppm)(Table II) in 2 , suggests the same *cis*-relationship between the methyl group and the oxygen at C-7. In the NMR study on 3 , irradiation at δ 1.13(H-10) gave a 17% NOE increment in the signal of H-9 and a 12% increment in the signal of 1-H. Thus, the structure of 5-hydroxycampenoside was established as 3 .

Table II. ^{13}C -NMR Spectral Data of Iridoids^{a)}

Compd.	C-1	C-3	C-4	C-5	C-6	C-7	C-8	C-9	C-10	C-11
2 ^{b)}	96.35	160.65	123.21	28.55	37.25	76.35	38.45	45.16	12.61	190.17
3 ^{b)}	94.37	160.77	124.79	69.39	44.46	74.94	37.74	53.71	12.52	190.61
5 ^{c)}	94.8	149.1	113.7	29.9	38.9	77.1	38.9	45.6	12.5	167.1
9 ^{d)}	98.06	153.18	112.53	36.25	39.40	128.26	144.22	46.59	61.13	171.11
10 ^{d)}	96.85	153.92	114.14	76.23	46.79	126.66	141.63	56.36	60.78	168.09

a) Chemical shifts in ppm relative to internal $(\text{Me})_4\text{Si}$. Additional signals arising from glucose were omitted. Compounds 2 and 3 also showed signals from the cinnamoyl ester part.

b) In DMSO-d_6 . c) In CDCl_3 . d) In CD_3OD .

ACKNOWLEDGMENTS

We wish to express our thanks to Prof. H. Inouye, Kyoto University, for a generous gift of loganin and to Prof. F. Murai, Aichi Medical University, for information on the IR spectrum of boschniakine. Authors are indebted to Prof. A. Numata, Osaka College of Pharmacy, and Assistant Professor K. Yoshikawa, Tokushima University, for valuable discussion about ^{13}C -NMR spectral data, and Mr. K. Murakami, Tokushima University, for collection of plants.

REFERENCES AND NOTES

1. T. Sakan, F. Murai, Y. Hayashi, Y. Honda, T. Shono, M. Nakajima, and M. Kato, Tetrahedron, 1967, 23, 4635.
2. H. Inouye, T. Yoshida, S. Tobita, and M. Okigawa, Tetrahedron, 1970, 26, 3905.
3. S.R. Jensen, S.E. Lyse-Petersen, and B.J. Nielsen, Phytochemistry, 1979, 18, 273.
4. R.K. Chaudhuri, F.U. Afifi-Yazar, O. Sticher, and T. Winkler, Tetrahedron, 1980, 36, 2317.
5. The signals of 9 and 10 can be compared, since the difference between the signal at C-2 in propionic acid and that in methyl propionate is small (0.6 ppm) [cf. E. Lippmaa and T. Pehk, Eesti NSV Tead. Akad. Tomi. Keem. Geol., 1968, 17, 210].
6. In pyridine- d_5 . Y. Ozaki, S. Johne, and M. Hesse, Helv. Chim. Acta, 1979, 62, 2708.
7. In D_2O . A. Bianco, M. Guiso, C. Iavarone, R. Marini-Bettolo, and C. Trogolo, Gazz. Chim. Ital., 1976, 106, 947.
8. In D_2O . Y. Takeda, H. Nishimura, and H. Inouye, Phytochemistry, 1977, 16, 1401.

Received, 13th May, 1981