

SYNTHESIS OF NORRUFESCINE

Mary D. Menachery and Michael P. Cava*

Department of Chemistry, University of Pennsylvania
Philadelphia, PA 19104

and

Keith T. Buck and William J. Prinz


Fries and Fries Division

Mallinckrodt, Inc.


Cincinnati, OH 45216

Abstract - The phenolic azafluoranthene alkaloid norrufescine 1 has been synthesized. The key step in the synthesis involved the formation of the tetracyclic azafluoranthene nucleus by a novel aprotic thermolysis of a condensed triazine precursor.

The azafluoranthenes constitute a small group of isoquinoline-derived alkaloids consisting of the non-phenolic bases rufescine (2)¹, imeluteine (3)¹ and triclisine (4)², and the phenolic bases norrufescine (1)^{1,3} and telitoxine (5)⁴. The three non-phenolic bases have been synthesized^{2,5} using a classical Pschorr approach, although no synthesis of either phenolic base has yet appeared. We now report a total synthesis of norrufescine (1) by a route which involves, as the key step, an unusual aprotic Pschorr-type cyclization of a condensed benzotriazine intermediate.




1. R = H; R¹ = H
2. R = CH₃; R¹ = H
3. R = CH₃; R¹ = OCH₃



4. R = H
5. R = OH

Reaction of β -(2,3,4-trimethoxyphenyl)ethylamine⁶ with the acid chloride of 2-nitro-5-benzyloxybenzoic acid⁷ gave the corresponding crystalline amide 6, mp 121-122°C, which was converted by phosphorus oxychloride in acetonitrile into the dihydroisoquinoline 7, mp 128-130°C (58% from amine).

Reduction of 7 with hydrazine and 5% palladium/charcoal in ethanol gave the aminophenol 8 in 70% yield. Diazotization of 8 in 1N H₂SO₄, basification of the solution to pH 8 and extraction into chloroform gave the amorphous permanganate-colored benzotriazine dipole 9 in 90% yield; in accord with structure 9, this material showed no infrared diazo band in the 2200 cm⁻¹ region. Treatment of dipole 9 with a little conc. HCl, followed by crystallization from CHCl₃-ether, gave yellow-orange crystals of the triazinium chloride 10, mp 187-190°C dec (no band 2200 cm⁻¹). The nmr spectra of 10 (CDCl₃) showed three methoxyls at δ 3.98, 4.14 and 4.25 (3H each) and the B-ring protons of the isoquinoline system as triplets centered at δ 3.38 and δ 4.97. In the aromatic region, four protons appeared, a singlet at δ 7.63, doublet of a doublet at δ 7.97, doublet at δ 8.42 ($J=9$ Hz) and a doublet at δ 8.83 ($J=2$ Hz). The uv absorption spectrum showed maxima at 230 (log ϵ 4.26), 273(4.20), 326(3.86), 375(3.99), 440(3.60) and 498 nm (3.58).



Thermolysis of the benzotriazinium salt 10 in refluxing chlorobenzene gave dihydronorrufescine 11 (24%) and the deaminated product (17%). Dihydronorrufescine crystallized from methanol/ether as yellow crystals, mp 240-242°C; nmr spectrum (CDCl₃-CD₃OD) δ 2.81 and 4.13 (2H each, t), 3.89, 3.97 and 3.96 (3H each, s) 6.87 (1H, dd, $J=8$ Hz and 2Hz), 7.20 (1H, d, $J=2$ Hz), 7.56 (1H, d, $J=8$ Hz). Dehydrogenation of 11 by 10% palladium/charcoal in refluxing *p*-cymene afforded norrufescine (1), mp 234-236°C (46%), identical (ir, nmr) with the natural base.

REFERENCES

1. M. P. Cava, K. T. Buck, I. Noguchi, M. Srinivasan and M. G. Rao, Tetrahedron, 1975, 31, 1667.
2. R. Huls, J. Gaspers and R. Warin, Bull. Soc. R. Sci. Liege., 1976, 45, 40.
3. M. D. Klein, K. T. Buck, M. P. Cava and D. Voet, J. Am. Chem. Soc., 1978, 100, 662.
4. M. D. Menachery and M. P. Cava, J. Nat. Prods., 1981, 320.
5. M. P. Cava, K. T. Buck and A. I. DaRocha, J. Am. Chem. Soc., 1972, 94, 5931.
6. S. Kubota, T. Masui, E. Fujita and S. M. Kupchan, J. Org. Chem., 1966, 31, 516.
7. E. Giovannini and P. Portmann, Helv. Chim. Acta, 1948, 31, 1381.

Received, 5th August, 1982