

HOMOLYTICAL ALKYLATION OF HETEROAROMATIC BASES BY CYCLIC ACETALS


Semen Solomonovitch Zlotsky*, Vladimir Victorovitch Zorin, Yuri Borisovitch Zeletchonok, and Dilus Lutfullitch Rakhmankulov
 Ufa Oil Institute, Kosmonavtov Str.,1, Ufa, 450062, USSR

Abstract—— Homolytical reaction between protonated bases and 1,3-diheterocycloalkanes was described.


Alkoxylic radicals are known to react with 1,3-oxaheterocycloalkanes and attack preferably methylene or methenyl groups adjacent to two heteroatoms¹⁻³. In this connection we describe unexpected and surprising data of the work⁴, in which the main product of alkylation initiated by $(\text{CH}_3)_3\text{C-O-O-H}+\text{FeSO}_4$ system of protonated quinoxaline (I) with 1,3-dioxolane (II) is 2-(1,3-dioxolane-4-yl)quinoxaline (III) but not isomeric 2-(1,3-dioxolan-2-yl)quinoxaline(IV).



We repeated this experiment using a system of $\text{R-O-O-H}+\text{FeSO}_4$ ($\text{R}=(\text{CH}_3)_3\text{C}; \text{C}_6\text{H}_5(\text{CH}_3)_2\text{C}$) as initiator and found out that product (IV) is formed more selectively than isomeric substances (III)(Table I). Then we drew some 1,3-oxaheterocycloalkanes (II, VI-VIII) into homolytical alkylation of protonated 2-methylquinoline (V) using a system of $\text{R-O-O-H}+\text{FeSO}_4$ ($\text{R}=\text{H}; \text{C}_6\text{H}_5(\text{CH}_3)_2\text{C}$) as donor of alkoxylic radicals. In this case of compounds (II,VI,VII) we established parallel formation of 2- and 4-substituted 1,3-dioxacycloalkanes (IX-XIV), whereas oxathiocyclane (VIII) gave 2-substituted product (XV) only.
















We found out that selectivity of products (IX,XII) formation from 1,3-dioxolane (II) depends on initiator nature (Table I). Under the influence of hydroxyl radicals (system $H_2O_2+FeSO_4$) 2- and 4-substituted products (IX,XII) are formed in equal amount whereas 4-(1,3-dioxolan-2-yl)-quinaldine (IX) is formed six times as much as its isomer (XII) under the influence of $C_6H_5C(CH_3)_2\dot{O}$ radicals. At the same time the type of initiator does not influence the selectivity of formation of products from acetals VI and VII. We think that the difference in strength of C^2-H and C^4-H bonds in reaction with $\dot{O}H$ and $C_6H_5C(CH_3)_2\dot{O}$ radicals practically does not exist. Simultaneously with the joining to aromatic system cyclic dialkoxyalkyl radicals react with ions Fe^{+3} resulting in formation of the corresponding carbocations which form in aqueous medium at first unstable 2-oxy-1,3-dioxacyclanes and then monoformates of glycols XVI-XVIII. The rate of this reaction considerably depends on the cycle range in dialkoxyalkyl radicals; for example: oxidation of six- and seven-membered radicals proceeds much more rapidly than of 1,3-dioxolane-2-yl radicals⁵⁻⁶. This fact is supposed to explain the decrease of selectivity of formation of X and XI in comparison with IX and XIV.



During this reaction monoformates of the corresponding diols (XVI-XVIII) are formed as accessory products. Reaction of protonated base with acetals (II,VI-VIII) was carried out in glass reactor under the following conditions: concentrated aqueous solution of $FeSO_4$ (0.06mol) and 30% solution of hydrogen peroxide (H_2O_2) or cumyl hydroperoxide ($C_6H_5C(CH_3)_2OOH$)(0.06mol) were added for 30 min intensively stirring at 5-10°C to the reaction mixture and pH 4~5. The reaction mixture contained 0.04 mol of the base (I,V) protonated by sulfuric acid and 0.02 mol of 1,3-diheterocycloalkane (II,VI-VIII); both were dissolved in 50 ml of water. The pH of mixture was maintained at level 4-5 by concentrated Na_2CO_3 solution. After the synthesis was finished pH of the solution was increased to level 12-13 by means of addition of NH_4OH solution and was extracted by ether. Ethereal extracts were joined together and vaped; products were isolated by column liquid chromatography method (Al_2O_3 , eluent, mixture of petroleum ether and diethyl ether in ratio 3.5:1).

TABLE I

The Yield of Hydrogen Atom Substitution Products in Quinaldine and Benzpyrazine on 1,3-Oxaheterocycloalkyl Residue at 5-10°; pH=4-5; reaction time 30 min

Acetals	Products	Yield of the reacted base, %	
		H ₂ O ₂ +Fe ²⁺	C ₆ H ₅ C(CH ₃) ₂ OOH + Fe ²⁺
 II	 IX  XII	35	76
	 IV  III	-	71
		-	8
 VI	 X  XIII	8	7
		22	24
 VII	 XI  XIV	2	3
		31	30
 VIII	 XV	-	90





TABLE II

Reaction Products Spectral Data and Physico-Chemical Constants

Compounds	n_D^{20} t ^{melt.} °C	PMR				NMR			
		Acetals		Quinaldine		Acetals		Quinaldine	
		C ² -H	C ⁴ -H, C ⁵ -H, C ⁶ -H, C ⁷ -H	Me	Ar	C ²	C ⁴ , C ⁵ , C ⁶ , C ⁷	Me	Ar
IX 4-(1,3-Dioxolan-2-yl)-quinaldine	1.6058	6.15s	3.86s	2.58s	7.0-8.0s	100.7d	65.2t	25.0q	118.2-158.5m
XII 4-(1,3-Dioxolan-4-yl)-quinaldine	1.5875	5.03d	4.10-4.70m 5.33t	2.58s	7.20-8.03m	96.0t	71.3t 74.3d	25.9q	117.7-158.3m
X 4-(1,3-Dioxane-2-yl)-quinaldine	-	5.62s	1.57-1.80m 3.75-4.05m	2.55s	7.10-7.93m	99.4d	32.4t 67.7t	26.6q	119.3-158.3m
XIII 4-(1,3-Dioxan-4-yl)-quinaldine	-	4.84d	1.75-2.00m 3.63-4.22m 4.87d	2.55s	7.08-8.00m	94.4t	34.4t 66.9t 74.4d	26.0q	118.5-158.7m
XI 4-(1,3-Dioxypan-2-yl)-quinaldine	-	6.00s	1.28-1.90m 5.67-4.00m	2.60s	7.05-8.05m	98.9d	26.2t 66.4t	25.1q	123.0-158.7m
XIV 4-(1,3-Dioxypan-4-yl)-quinaldine	-	4.80d	1.30-1.91m 5.20d	2.60s	7.10-8.03m	93.7d	28.9t 36.7t 67.4t 75.2d	25.1q	118.1-158.3m
XV 4-(1,3-Oxathiolane-2-yl)-quinaldine	74	6.36s	2.81-3.17m 3.70-4.00m 4.23-4.55m	2.62s	7.06-8.05m	83.2d	34.4t 72.5t	25.8q	117.6-159.1m

REFERENCES

1. G.W.Hartgerink, L.C.Van der Laan, I.B.Engberts and Th.De Boer, Tetrahedron, 1971,27,4323.
2. M.I.Perkins, B.P.Roberts, J.Chem.Soc.Perkins Trans., 1975,1,77.
3. A.L.I.Beckwith and P.K.Tindal, Austral.J.Chem., 1971,24,2099.
4. W.Buratti, G.P.Gardini, F.Minisci, F. Bertini, R.Galli and M.Perchinunno, Tetrahedron, 1971,27,3655.
5. V.V.Zorin, S.S.Zlotsky, V.F.Shuvalov, A.P.Moravsky, D.L.Rakhmankulov and Ya.M.Paushkin, DAN SSSR, 1977,136,106.
6. V.N.Trifonova, V.V.Zorin, S.S.Zlotsky and D.L.Rakhmankulov, Zhurnal Organit-cheskoy Khimii, 1982,18,1050.

Received, 5th November, 1984