

1,3-CYCLOADDITIONS OF 2,2,4,4-TETRAMETHYLCYCLOBUTANE-1-ONE-3-THIONE S-METHYLIDE


Rolf Huisgen,* Grzegorz Mloston, and Claudia Fulka

Institut für Organische Chemie der Universität München,
Karlstr. 23, D-8000 München 2, Federal Republic of Germany

Abstract - The 1,3,4-thiadiazoline 2, prepared from 2,2,4,4-tetramethylcyclobutane-1-one-3-thione and diazomethane, loses N₂ at 40°C; the thiocarbonyl ylide 5 is a transient intermediate which *in situ* undergoes 1,3-dipolar cycloadditions to electron-deficient dipolarophiles with C≡C, C=C, C=S, C=O, and N=N bonds. Structures and mechanism are discussed.

Although simple aliphatic thioketones are known,^{1,2} only the sterically hindered representatives are storable in the *monomeric* state. Their merciful inertness toward the human olfactory system contributes to the ease of handling.

In 1970 Diebert³ reacted the red 2,2,4,4-tetramethylcyclobutane-1-one-3-thione (1)⁴ with diazomethane and isolated 67% of the spirothiirane 3; the structure of the spiro-1,3,4-thiadiazoline (2) was assigned to an intermediate which was observed at 0°C as a white solid.



We passed gaseous diazomethane into 3 mmoles of 1 in 10 ml ether at -78°C, evaporated at 0°C and dissolved the crystalline residue in 10 ml pentane. After 30 min at

-78°C, 88% of the colorless 2 were filtered, mp 40-42°C (dec.); ¹H NMR (CDCl₃): δ = 1.23 (s, 2 CH₃), 1.31 (s, 2 CH₃), 5.84 (s, 5-H₂). Correct CH, N, S analyses of 2 were obtained. The N₂ extrusion of 2 obeyed first-order kinetics with t_{1/2} = 25 min at +45°C in xylene, in agreement with Diebert's measurement (t_{1/2} = 22 min at 49°C in CCl₄).³ 2,2-Diphenyl-1,3,4-thiadiazoline (4) loses N₂ with t_{1/2} = 55 min in THF at -45°C⁵ whereas 2,2,5-triphenyl-1,3,4-thiadiazoline decomposes at -65°C with t_{1/2} ~ 30 sec.⁶ Thus, *alkylated* thiadiazolines are more stable than arylated ones.

The elimination of N₂ from 2 is a 1,3-dipolar cycloreversion and the thione *S*-methylide 5 the expected intermediate. The electrocyclization of the planar thiocarbonyl ylide system in 5 giving the thirane 3 is accompanied by two 90° rotations about the CS bonds. Such processes require substantial activation energy because most of the resonance energy of 5 is sacrificed in the transition state. Experience shows that *intermolecular* interception by active dipolarophiles is usually faster. The pioneer work of Buter, Wassenaar, and Kellogg⁷ demonstrated that thiocarbonyl ylides are involved in the N₂ extrusion from alkylated thiadiazolines. We have recently described 1,3-dipolar cycloadditions of adamantanethione *S*-methylide (8), prepared in analogy to 5, to electron-deficient dipolarophiles with CC double and triple bond as well as CO and CS double bond.^{8,9}


The thiocarbonyl ylide 5 is basic. When 2 was warmed in methanol with a drop of trifluoroacetic acid for 8 h at 40°C, ¹H NMR analysis indicated 80% of the *O,S*-dimethyl-acetal 7; mp near 20°C, ¹H NMR (CDCl₃): δ = 1.32 (s, 4 CH₃), 1.92 (s, SCH₃), 3.37 (s, OCH₃). The protonated species 6 is a logical intermediate and the CH₂ group of 5 is the more basic center among the two termini.

2,2,4,4-Tetramethylcyclobutane-1-one-3-thione *S*-methylide (5) is an active 1,3-dipole despite a steric encumbrance at one terminus which by far exceeds that of 8. After warming pure 2 in THF in the presence of 1.1 equiv of dipolarophile for 8 h at 40°C, the evolution of N₂ was complete. The less active dimethyl acetylenedicarboxylate was used in excess as solvent. Quantitative ¹H NMR analysis (CDCl₃) by comparison with a weighed standard (1,1,2,2-tetrachloroethane) afforded the yields listed in Table 1. The isolated adducts were characterized by analyses and spectra.

The NMR spectra of the cycloadducts are reliable probes of symmetry or chirality, respectively. Adducts 9 and 10 show singlets for 5-H₂ and pairwise equivalence of the methyl signals: δ 1.25 and 1.42 for 9, 1.45 and 1.95 for 10. The ¹³C NMR spec-

Table 1. Reactions of thiadiazoline 2 with dipolarophiles in THF (8 h 40°C); adduct yields based on ¹H NMR analysis (isolation)

Dipolarophile	% adduct	mp (°C)	Formula
Dimethyl acetylenedicarboxylate	87	76-78	<u>9</u>
Fumaronitrile	93	111-113	<u>11</u>
Maleonitrile	92	140-142	<u>12</u>
Maleic anhydride	(76)	207-209	<u>13</u>
N-Phenylmaleimide	(76)	194-196	<u>14</u>
Tetracyanoethylene	73	213-215	<u>10</u>
<i>trans</i> -β-Nitrostyrene	72	121-123	<u>15</u>
Thiobenzophenone	56	124-126	<u>17</u>
	+ 23	(not isol.)	<u>18</u>
Thioxanthione	(95)	174-177 ^o	<u>19/20</u>
Adamantanethione	88	139-141	<u>21</u>
2,2,4,4-Tetramethyl-cyclo-butane-1-one-3-thione	76	159-161	<u>22</u>
Chloral	95	65-67	<u>24</u>
Dimethyl azodicarboxylate	58	110-112	<u>25</u>




trum of 10 records δ 67.8 for C-2' + C-4', 22.9 and 22.7 for 4 CH₃ as well as 110.0 and 110.5 for 4 CN. Complex ABXY patterns appear in the ¹H NMR spectra of the chiral adducts 11-15, sometimes only the doublet of 3-H being resolved. All the methyl groups are different, *e.g.*, δ 1.30, 1.42, 1.55, 1.65 for 11. In the ¹³C NMR (CDCl₃) of 13 the 4 CH₃ were observed and three singlets are assigned to C-2', C-3', and C-4'.

trans- β -Nitrostyrene furnished only isomer 15 within the limits of the ^1H NMR spectrum. The doublet of 3-H at δ 5.45 - in contrast to $\delta(3\text{-H}) = 3.87$ for 11 and 12 - leaves no doubt that the deshielding nitro group is located in 3-position. Thus, the CH_2 terminus of 5 is the more nucleophilic center.

The cycloadditions to fumaronitrile and maleonitrile proceed stereospecifically; within the ^1H NMR analytical limits no mutual admixture of 11 and 12 could be recognized. $J_{3,4}$ amounts to 2.3 Hz for 11 and 4.5 Hz for 12, compared with 3.0 Hz for 15 and 7.0, 6.2 Hz for 13 and 14.

The base peak of most mass spectra corresponds to $\text{M}^+ - \text{C}_4\text{H}_6\text{O}$, *i.e.*, the elimination of dimethylketene provides radical cations 16. Also the peak of dimethylketene radical cation is present, and the ratio of the two peaks depends on the ionization potentials: 100:95 for 11, 100:90 for 12, 100:20 for 13, but 5:100 for the tetracyanoethylene adduct 10. The base peak of 15 is m/e 203 ($\text{M}^+ - \text{C}_4\text{H}_6\text{O} - \text{NO}_2$) and the occurrence of m/e 104 (styrene $^+$, 35%) is in accordance with the addition direction.




Both orientations were used by thiobenzophenone as a dipolarophile: 56% 17 and 23% 18. The minor product was enriched by thick-layer chromatography, but not obtained pure. The major product 17 likewise resulted from the reaction of thiobenzophenone *S*-methylide with the thione 1 (90%) at -45°C ; this convergence proves the presence of $-\text{S}-\text{CH}_2-\text{S}-$ in 17. Aromatic thione *S*-methylides always follow this course in their "Schönberg reactions" with dipolarophilic CS double bonds,^{10,11} *i.e.*, the *type A* of the preceding communication.⁹ Conceivably, it is steric hindrance which makes the *type B* addition - yielding 18 here - competitive. Structures 17 and 18 are confirmed by the ^{13}C NMR diagnostic criterion previously used.⁹ $\delta(\text{C}-2)$ of 17 emerges at 25.3

and $\delta(\text{C-5})$ of 18 at 50.0. The corresponding ^1H NMR singlets appear at δ 3.32 for 2-H₂ of 17 and at 3.75 for 5-H₂ of 18. A plane of symmetry reduces the number of ^1H and ^{13}C signals for 17 and 18.

Thioxanthione afforded a 5:1 mixture of 19 and 20 in high yield, but the separation did not succeed. The ^{13}C triplets of 2-CH₂ (19) and 5-CH₂ (20) were found at δ 37.4 and 45.0. In the ^1H NMR spectrum the CH₃ pairs of 19 emerge at δ 0.79 and 1.24 whereas those of 20 coincide at 1.31; the high-field absorption of 19 (0.79) reveals the shielding by the aromatic nuclei.

The addition of 5 to adamantanethione (88% 21) and thione 1 (76% 22) followed the *type B*⁹ pattern with its lower steric requirements. A diminished dipolarophilic activity of 1 is concluded from the modest yield of 22 and the presence of 10% thiirane 3. Diebert³ isolated 31% of 22 when he treated thione 1 with 0.8 equiv diazomethane. The analogy with Schönberg's dithiolane formation¹⁰ was recognized,³ but its mechanism was not yet understood at the time. The first 1,3-dithiolane of *type B* was described in 1965 as a 2:1 product from hexafluorothioacetone and diazomethane (-78°C).¹²

^{13}C triplets at δ 47.0 (21) and 43.4 (22) for C-5 mark both adducts as belonging to *type B*. According to ^1H and ^{13}C NMR evidence, 22 and 21, possesses *one* C₈ plane, *i.e.*, the two cyclobutanone systems in the doubly spiro linked 22 are non-equivalent. The mass spectra corroborate structures 21 and 22. Besides the elimination of dimethylketene (M⁺ - C₄H₆O), the fragmentation of 21 into C₈H₁₂S₂O⁺ (15%) - probably 23 - and C₁₁H₁₆⁺ (2-methyleneadamantane, 15%) is indicative of *type B*.⁹



The expectation of structure 24 for the chloral adduct (95% yield) is confirmed by the ABX spectrum of the 1,3-oxathiolane protons: dd of 5-H at δ 4.55. As foreseen, small signals, $m/e = 318$ and 316, occur as parent peaks due to ^{37}Cl and ^{35}C isotopes; $m/e = 248$ (85%) and 246 (85%) are the peaks for M⁺ - C₄H₆O whereas $m/e = 70$ (C₄H₆O⁺) constitutes the base peak. The interaction of 2 with dimethyl azodicarboxylate led to 25. Its heteroring rests in a frozen non-planar conformation, as the

AB spectrum of the 5-H₂ (δ 4.38 and 4.58, $J = 7.5$ Hz) and the nonequivalence of the methyl groups (δ 1.27 for 2 CH₃, 1.40, 1.65) testify.

The synthetic potential of the cycloadditions of thiocarbonyl ylides goes beyond the preparation of 5-membered S-containing heterocycles. The easy removal of sulfur by CS hydrogenolysis recommends the reaction sequence as a method of CC linking of three units (thione, diazoalkane, dipolarophile).

We presume that a thermodynamic factor contributes to the superiority of the CS over the CO double bond as dipolarophile. The π bond which is sacrificed in the cycloaddition amounts to 94 kcal mol⁻¹ for CO (ketones) and to 55-60 kcal mol⁻¹ for CS; a portion of the difference may become effective in the transition state of the concerted cycloaddition.

ACKNOWLEDGMENT

The Fonds der Chemischen Industrie kindly supported the work. G.M. is grateful to the Alexander von Humboldt Foundation for a fellowship.

REFERENCES

1. S. Bleich and R. Mayer, *Chem. Ber.*, 1966, 99, 1771.
2. D. Paquer and J. Vialle, *Bull. Soc. Chim. France*, 1972, 3138.
3. C.D. Diebert, *J. Org. Chem.*, 1970, 35, 1501.
4. E.U. Elam and H.E. Davis, *J. Org. Chem.*, 1967, 32, 1563.
5. I. Kalwisch, Li X., J. Gottstein, and R. Huisgen, *J. Am. Chem. Soc.*, 1981, 103, 7039.
6. R. Huisgen and Li X., *Heterocycles*, 1983, 20, 2363.
7. J. Buter, S. Wassenaar, and R.M. Kellogg, *J. Org. Chem.*, 1972, 37, 4045.
8. R. Huisgen and G. Mloston, *Tetrahedron Lett.*, 1985, 26, 1049.
9. G. Mloston and R. Huisgen, *Heterocycles*, preceding communication.
10. A. Schönberg, B. König, and E. Singer, *Chem. Ber.*, 1967, 100, 767.
11. Li X. and R. Huisgen, *Tetrahedron Lett.*, 1983, 24, 4181.
12. W.J. Middleton and W.H. Sharkey, *J. Org. Chem.*, 1965, 30, 1384.

Received, 21st May, 1985