

DEMETHYLATIONS IN THE CULARINE SERIES

Angel Rodriguez de Lera, Carmen Villaverde, and Luis Castedo*

Departamento de Química Orgánica de la Facultad de Química y Sección de Alcaloides del CSIC, Santiago de Compostela, Spain

Abstract - Selective demethylation in 7,4',5'-substituted cularine compounds can be achieved under two different sets of conditions (acidic and nucleophilic). Less satisfactory results are obtained in the 7,3',4'-substituted analogs.

INTRODUCTION

The methylation of phenols and the selective O-demethylation of aryl methyl ethers constitutes an important procedure for the interconversion of naturally occurring compounds.¹


The cularines are a group of isoquinoline alkaloids² whose number has recently been increased by the isolation of several new members, mainly of a phenolic nature.^{3,4} This fact makes the cularines an interesting choice for demethylation studies. For this purpose we selected two general reagents⁵ presenting opposite but in some instances complementary behaviour:

- A) 48% HBr/AcOH, an acidic reagent which causes cleavage by a combination of SN1 and/or a partially charged cyclic transition state.
- B) EtS⁻, a nucleophilic reagent in which the cleavage pattern is essentially SN2.

RESULTS AND DISCUSSION

Demethylation of 3',4'-substituted cularines.- The selective demethylation of cularine 1 under acidic conditions (48% HBr/AcOH, 135°C, 35 min) has been reported⁶ as giving 7,3'-didemethylcularine (breoganine) 6. However, in our hands a mixture of demethylated compounds 2-6 was obtained (Table I), breoganine 6 being the major component. By using milder conditions but longer times (65-70°C, 144 h) a mixture of the diphenolics celtisine 5 and breoganine 6 resulted (Table I). These results can be explained assuming that the methoxyl groups at positions 7 and 3' are more basic than that at 4' due to a mesomeric contribution of the oxepinic oxygen.


Under nucleophilic conditions⁷ the demethylation of cularine 1 occurred preferentially at position 4' (see Table I). When these conditions (NaSEt 5mmol, reflux DMF, 7 h) were used, the monophenolic cularines 3 and 4 produced only the diphenolics celtisine 5 (42%) and breoganine 6 (45%) respectively, together with a certain amount of the initial monophenolic material. The formation of a phenoxide anion in ring D now deactivates the methoxyl group at the ortho-position for a further nucleophilic attack. Treatment of monophenolic cularidine 2 with NaSEt (3mmol, reflux DMF, 2 h) gave the expected diphenols 5 (43%) and 6 (26%) in similar proportions as in the case of cularine 1.



Reaction conditions	48% HBr/AcOH(1:1)		NaSEt(3mmol) DMF/reflux	NaSEt(5mmol) DMF/reflux
	135°C	65-70°C		
Product	35 min	144 h	40 min	4 h
cularidine <u>2</u> (R ₁ =H, R ₂ =R ₃ =Me)	3	--	15	--
celtine <u>3</u> (R ₁ =R ₃ =Me, R ₂ =H)	3	--	33	--
3'-O-demethyl- cularine <u>4</u> (R ₁ =R ₂ =Me, R ₃ =H)	10.5	--	21	--
celtisine <u>5</u> (R ₁ =R ₂ =H, R ₃ =Me)	18	21	13	43
breoganine <u>6</u> (R ₁ =R ₃ =H, R ₂ =Me)	30.5	36	10	18

Table I.- Yields in % for demethylation products of cularine 1 (R₁=R₂=R₃=Me) under different conditions. All products have been identified by comparison with authentic samples.³ Separation of cularidine 2 and celtine 3 was accomplished by their transformation into their O-acetyl derivatives (see experimental).

Demethylation of 4',5'-substituted cularines.- The demethylation of sarcocapnine 7 under various conditions indicated in Table II occurred preferentially at its more hindered methoxyl group, that at 5' position. Under milder acidic conditions (60°C) demethylation at 5' was mainly observed, but a higher temperature (90°C) resulted in catechol 10, which has not been found in nature. Its identity was confirmed by conversion into the methylenedioxy derivative 12 (R₁=Me, R₂+R₃=-CH₂-) through methylenation with CH₂Br₂.⁸ However, a change in demethylation selectivity was observed when sarcocapnine 7 was treated with NaSEt, which gave clavicine 9. The initial loss of the methyl group at 5' position now ensures that the second demethylation takes place at 7 position.



Reaction conditions	48% HBr/AcOH(1:2)			NaSEt(3mmol) DMF/reflux	NaSEt(5mmol) DMF/reflux
	60°C	60°C	90°C		
Product	14 h	30 h	60 h	1 h	2 h
sarcocapnine <u>7</u> (R ₁ =R ₂ =R ₃ =Me)	31	--	--	28	--
sarcocapnidine <u>8</u> (R ₁ =R ₃ =Me, R ₂ =H)	56	58	--	6	--
clavicine <u>9</u> (R ₁ =R ₂ =H, R ₃ =Me)	--	--	--	22	77
<u>10</u> (R ₁ =Me, R ₂ =R ₃ =H)	--	10	47	--	--
<u>11</u> (R ₁ =R ₂ =R ₃ =H)	--	--	8.5	--	--

Table II.- Yields in % for demethylation products of sarcocapnine 7 (R₁=R₂=R₃=Me) under different conditions. All known products have been identified by direct comparison with authentic samples.³

EXPERIMENTAL

All melting points were measured in a Büchi apparatus and are uncorrected. NMR spectra were determined with a Bruker WM-250 spectrometer using TMS as internal reference. Mass spectra were run on a KRATOS MS-25 instrument operating at 70 eV. Microanalysis for C, H, N, were carried out on a Perkin-Elmer 240-B elemental analyzer.

1 mm and 0.2 mm layers of Merck 60 GF₂₅₄ silica gel were normally used for preparative and analytical thin layer chromatography, respectively; the solvent systems were benzene-ethyl acetate-diethyl amine (7:2:1) for initial separation of phenolic compounds and methylene chloride-ethyl alcohol (9.5:0.5) for further purification and for separation of non-phenolic compounds.

48% HBr was redistilled over a trace of 50% hypophosphorous acid. 80% NaH in oil dispersion was supplied by Merck.

General procedure

A) Demethylation in acidic conditions.- 0.58 mmol of substrate were dissolved in 3 ml of 48% HBr-AcOH (1:1 or 1:2) and heated under an inert atmosphere during the time indicated. The reaction mixture was diluted with water (10 ml), neutralized with NaHCO₃ and extracted with CH₂Cl₂ (5x15 ml). The extracts were dried and evaporated to give a residue which was subjected to preparative thin layer chromatography.

B) Demethylation using a nucleophilic reagent.- Ethanethiol (2.1 mmol) dissolved in dry DMF (1 ml) was added dropwise to a suspension of sodium hydride (2.1 mmol) in dry DMF (2 ml) under an atmosphere of nitrogen. The mixture was stirred for 5 min before addition of a solution of the cularidine compound (0.70 mmol) in dry DMF (2 ml) and then refluxed during the time indicated in the table, evaporated, water added, and extracted with CH₂Cl₂ (5x20 ml). The extracts were dried and evaporated to give a solid residue which was purified as above.

Separation of cularidine 2 and celtine 3.- The fraction containing cularidine 2 and celtine 3 was dissolved in dry pyridine (1 ml), 0.5 ml of acetic anhydride added and the solution refluxed during 1 h. Then the mixture was evaporated, water added and after extraction with CH₂Cl₂ afforded a mixture of acetates which was subjected to preparative tlc.

O-Acetylcularidine.- Crystallized from ethanol-ether as its hydrochloride mp 250-252°C (dec.). PMR (CDCl₃, 250MHz, δ): 2.39(s, 3H, OCOCH₃), 2.57(s, 3H, NMe), 2.70-3.30(m, 6H, 3x-CH₂), 3.79(s, 3H, OMe), 3.84(s, 3H, OMe), 4.35(dd, J_{AX}=11.7 Hz, J_{BX}=3.5 Hz, 1H, H-1), 6.50(s, 1H, Ar-H), 6.56(s, 1H, Ar-H), 6.88(d, J=8.3 Hz, 1H, Ar-H), 6.92(d, J=8.3 Hz, 1H, Ar-H). MS m/e(%): 369(M⁺, 57), 354(100), 312(91), 69(82). Anal. Calcd for C₂₁H₂₃NO₅ (free base): C, 68.29; H, 6.23; N, 3.79. Found: C, 68.01; H, 6.29; N, 4.07.

O-Acetylceltine.- Crystallized from ethanol mp 175-177°C. PMR(CDCl₃, 250 MHz, δ): 2.29(s, 3H, OCOCH₃), 2.57(s, 3H, NMe), 2.72-3.33(m, 6H, 3x-CH₂-), 3.74(s, 3H, OMe), 3.81(s, 3H, OMe), 4.44(dd, J_{AX}=12 Hz, J_{BX}=3.7 Hz, 1H, H-1), 6.60(s, 1H,

Ar-H), 6.74(d, J=8.4 Hz, 1H, H-6), 6.86(d, J=8.4 Hz, 1H, H-5), 7.00(s, 1H, Ar-H). MS m/e(%): 369(M⁺, 68), 354(44), 326(17), 312(100). Anal. Calcd for C₂₁H₂₃NO₅: C, 68.29; H, 6.23; N, 3.79. Found: C, 68.33; H, 6.34; N, 3.45.

Hydrolysis of O-acetyl compounds.- To a solution of 0.05 g (0.135 mmol) of O-acetyl compound in methanol (5 ml) was added in one portion 0.1 g of anhydrous Na₂CO₃. The mixture was stirred at room temperature during 1 h, diluted with water (25 ml), saturated with NH₄Cl and extracted with CH₂Cl₂ (3x15 ml) to afford the corresponding phenolic compound.

Diphenol 10.- Crystallized from ethanol-ether as its hydrochloride mp 248-250°C (dec.). PMR(CDCl₃), 250 MHz, δ): 2.59(s, 3H, NMe), 2.84-3.47(m, 6H, 3x-CH₂-), 3.87(s, 3H, OMe), 4.53(dd, J_{AX}=12.3 Hz, J_{BX}=4.2 Hz, 1H, H-1), 6.47, 6.59, 6.76 and 6.91(4d, J=8.4 Hz, 1H each, 4xAr-H). MS m/e(%): 313(M⁺, 100), 298(27), 296(40) 282(27), 175(19), 174(31). Anal. Calcd for C₁₈H₂₀NO₄Cl: C, 61.80; H, 5.72; N, 4.00. Found: C, 61.30; H, 5.96; N, 3.63.

Triphenol 11.- Highly hygroscopic material. Crystallized as its hydrochloride mp 206-208°C. PMR(CDCl₃+CD₃OD, 250 MHz, δ): 2.58(s, 3H, NMe), 2.60-3.38(m, 6H, 3x-CH₂-), 4.48(dd, J_{AX}=12.2 Hz, J_{BX}=4.0 Hz, 1H, H-1), 6.44(d, J=8.5 Hz, 1H, Ar-H), 6.56(d, J=8.5 Hz, 1H, Ar-H), 6.74(d, J=8.3 Hz, 1H, Ar-H), 6.80(d, J=8.3 Hz, 1H, Ar-H). MS m/e(%): 299(M⁺, 100), 272(82), 256(11), 255(11), 132(32). Anal. Calcd for C₁₇H₁₈NO₄Cl.H₂O: C, 57.70; H, 5.65; N, 3.96. Found: C, 57.38; H, 5.54; N, 3.70.

Methylenation of catechol 10.- Anhydrous KF (0.082 g, 1.58 mmol) was added to a solution of catechol 10 (0.1 g, 0.32 mmol) in anhydrous DMF (2 ml). Then dibromomethane (0.061 g, 0.352 mmol, 10% excess) was added to the solution and the mixture heated at 110°C under an argon atmosphere for 2 h. Evaporation to dryness followed by addition of water (15 ml), extraction with CH₂Cl₂ (5x15 ml), drying and evaporation afforded a residue which was separated by tlc to give 0.030 g of 10 and 0.035 g of 12.

Compound 12 crystallized from ethanol-ether as its hydrochloride mp 172-174°C (dec.). PMR(CDCl₃, 250 MHz, δ): 2.62(s, 3H, NMe), 2.80-3.38(m, 6H, 3x-CH₂-), 3.88(s, 3H, OMe), 4.42(dd, J_{AX}=11.8 Hz, J_{BX}=3.0 Hz, 1H, H-1), 5.98(d, J=1.4 Hz, 1H, OCH₂O), 6.09(d, J=1.4 Hz, 1H, OCH₂O), 6.52(d, J=8.1 Hz, 1H, Ar-H), 6.56(d, J=8.1 Hz, 1H, Ar-H), 6.80(d, J=8.4 Hz, 1H, Ar-H), 6.91(d, J=8.4 Hz, 1H, Ar-H). MS m/e(%): 325(M⁺, 100), 310(31), 308(52), 294(27), 174(50). Anal. Calcd for C₁₉H₂₀NO₄Cl: C, 63.07; H, 5.53; N, 3.87. Found: C, 62.81; H, 5.42; N, 4.21.

ACKNOWLEDGEMENT

We thank the Comisión Asesora (Spain) for its financial support.

REFERENCES

1. L. Castedo, A. Rodriguez de Lera, J. M. Saá, R. Suau, and C. Villaverde, Heterocycles, 1980, 14, 1135.
2. a) L. Castedo, "The Chemistry and Pharmacology of Cularine Alkaloids", in The Chemistry and Biology of Isoquinoline Alkaloids, ed. by Phillipson et al., Springer-Verlag Berlin Heidelberg, 1985, pp. 102-125.
b) B. Gözler and M. Shamma, J. Nat. Prod., 1984, 47, 753.
3. a) J. M. Boente, L. Castedo, R. Cuadros, A. Rodriguez de Lera, J. M. Saá, R. Suau, and M. C. Vidal, Tetrahedron Lett., 1983, 24, 2303.
b) J. M. Boente, L. Castedo, A. Rodriguez de Lera, J. M. Saá, R. Suau, and M. C. Vidal, Tetrahedron Lett., 1984, 25, 1829.
4. a) D. P. Allais and H. Guinadeau, J. Nat. Prod., 1983, 46, 881.
b) H. Guinadeau and D. P. Allais, Heterocycles, 1984, 22, 107.
5. M. V. Bhatt and S. V. Kulkarni, Synthesis, 1983, 249.
6. T. Kametani, S. Shibuya, and S. Sasaki, Yakugaku Zasshi, 1967, 87, 198.
7. G. I. Feutrill and R. N. Mirrington, Tetrahedron Lett., 1970, 1327; Aust. J. Chem., 1972, 25, 1719, 1731.
8. J. H. Clark, H. L. Holland, and J. M. Miller, Tetrahedron Lett., 1976, 38, 3361.

Received, 9th August, 1985