

**AN IMPROVED AND DIVERGENT INTRODUCTION OF THE STREPTONIGRIN AND LAVENDAMYCIN
QUINOLINE-5,8-QUINONE AB RING SYSTEMS**


Dale L. Boger^{*1} and Masami Yasuda

Department of Chemistry, Purdue University


West Lafayette, IN 47907, U.S.A.

Abstract — An improved and divergent approach to the introduction of the streptonigrin and lavendamycin quinoline-5,8-quinone AB ring systems is detailed and is based on the selective and controlled nucleophilic substitution of common 7-bromoquinoline-5,8-quinone intermediates derived from the Friedländer condensation products of 2-amino-3-benzyloxy-4-bromobenzaldehyde.


Streptonigrin (1),² lavendamycin (2),³ and streptonigrone (3),⁴ three structurally and biosynthetically related natural products isolated from *Streptomyces flocculus* and *Streptomyces lavendulae*, each possess a characteristic and functionalized quinoline-5,8-quinone AB ring system.⁴ Early efforts on streptonigrin^{5,6} and more recent efforts on lavendamycin^{5,7} have confirmed that the minimal structural component of the naturally occurring antitumor-antibiotics necessary for observed *in vitro* cytotoxic and antimicrobial activity consists of this substituted quinoline-5,8-quinone AB ring system. In conjunction with synthetic efforts on the total synthesis of streptonigrin,⁸ streptonigrone⁹ and lavendamycin¹⁰ and in efforts designed to delineate the structural features of the naturally-occurring antitumor-antibiotics which potentiate the antitumor and antimicrobial activity of the quinoline-5,8-quinones we have devised an improved, divergent approach to the introduction of the required quinoline-5,8-quinone systems.¹¹



The approach which has been implemented in efforts required for the introduction of the lavendamycin 7-aminoquinoline-5,8-quinone AB ring system^{10b} has been extended as detailed herein to provide an improved method for the introduction of the streptonigrin 7-amino-6-methoxyquinoline-5,8-quinone AB ring system and is based on the selective and controlled nucleophilic substitution of common 7-bromoquinoline-5,8-quinone intermediates derived from the Friedländer condensation products of 2-amino-3-benzyloxy-4-bromobenzaldehyde,^{10b} equation 1.¹² Consequently the approach permits the introduction of the streptonigrin 7-amino-6-methoxyquinoline-5,8-quinone or the lavendamycin 7-aminoquinoline-5,8-quinone AB ring systems from common intermediates.



As detailed in earlier work,¹⁰ direct azide displacement at C-7 of 7-bromoquinoline-5,8-quinones under carefully controlled conditions followed by subsequent triphenylphosphine mediated azide reduction, which proceeds without competing quinone to hydroquinone reduction and with the generation of a stable and isolable triphenylphosphine imine,^{10b} permits the introduction of the 7-aminoquinoline-5,8-quinone system, equation 2.¹³ In a complementary sequence, treatment of 7-bromoquinoline-5,8-quinones with cerium(III) methoxide in methanol leads to C-6 methoxy substitution in which cerium(III) cation coordination with the quinoline substrate serves to direct nucleophilic attack to C-6 and which proceeds with the apparent subsequent stabilization of the initial addition product, equation 2 - 3. This stabilization of the hydroquinone addition product prevents subsequent elimination - addition reactions observed when conventional conditions (sodium methoxide - methanol) are employed for the reaction. Table I summarizes a representative series of experimental results of this C-6 directed nucleophilic substitution of 7-bromoquinoline-5,8-quinones. The nucleophilic substitution reaction of methoxide does proceed at a decelerated and controlled rate in the presence of cerium(III) and the cerium(III) cation coordination does not, in these instances,¹⁴ accelerate the apparent rate of nucleophilic addition.



Eq 2

a R = H





Table I. C-6 Nucleophilic Substitution of 7-Bromoquinoline-5,8-quinones.

Substrate	Conditions		Product	% Yield
	equiv of NaOMe (equiv of CeCl ₃)	temp, °C (time, h)		
5a ^{10b}	2-3	0 (0.1-0.3)	8a	0
	1.1	0 (1 min)		25
	2 (1, Ce(NO ₃) ₃ ·6H ₂ O)	0 (2)		33
	3 (1, CeCl ₃ ·7H ₂ O)	0 (2), 25 (1)		40
	2 LiOCH ₃ (1, CuCl ₂)	25 (0.5)		21
	3 (1)	0 (0.5)		20
	2 (1)	0 (0.5), 25 (1)		45
5b ^{10b}	2 (1)	0 (1), 25 (5.5)	8b	39
	2 (1)	0 (1), 25 (10)		60



Eq 3

Implementation of this work in the preparation of the naturally occurring antitumor-antibiotics and related, synthetic quinoline-5,8-quinones is in progress.

EXPERIMENTAL

General Procedure for the Preparation of 7-Bromo-6-methoxyquinoline-5,8-quinones. Table I: A stirred solution of anhydrous cerous chloride and sodium methoxide in methanol was treated with substrate (5a/5b, 0.1 mmole) at 0 °C under a N₂ atmosphere and the mixture was stirred at 25 °C. The solution was diluted with H₂O (20 ml) and extracted with CH₂Cl₂ (3 x 10 ml). The combined extracts were washed with saturated aqueous NaCl (1 x 20 ml), dried (Na₂SO₄), and concentrated in vacuo. Chromatography (SiO₂, 1 x 8 cm, 40% EtOAc-hexane eluant) afforded the 7-bromo-6-methoxyquinoline-5,8-quinones **8a/8b** as yellow solids. For 7-bromo-6-methoxyquinoline-5,8-quinone (**8a**): mp 183-185 °C (lit.^{13a} mp 185-187 °C dec.): ¹H NMR (CDCl₃) δ 9.04 (1H, dd, J = 4.7 Hz, 1.7 Hz, C-2 H), 8.43 (1H, dd, J = 7.9 Hz, 1.7 Hz, C-4 H), 7.69 (1H, dd, J = 7.9 Hz, 4.7 Hz, C-3 H), 4.35 (3H, s, OCH₃).

For 7-Bromo-6-methoxy-2-(2'-pyridyl)quinoline-5,8-quinone (**8b**): mp 178-179 °C dec.; ¹H NMR (CDCl₃) δ 8.84 (1H, d, J = 8.3 Hz, C-4 H) 8.59-8.78 (2H, m, C-3' H and C-6' H), 8.50 (1H, d, J = 8.3 Hz, C-3 H), 7.90 (1H, dt, J = 7.9 Hz, 1.6 Hz, C-4' H), 7.29-7.48 (1H, m, C-5' H), 4.36 (3H, s, OCH₃); IR(KBr) ν_{max} 3080, 2930, 1674, 1586, 1563, 1452, 1435, 1334, 1307, 1250, 1143, 1078 cm⁻¹; CIMS m/z (rel. intensity) 345/347 (M + 1, 1/1, base), 333(2.5), 267(4); HRMS m/z for C₁₅H₉N₂O₃Br requires 343.9797; found 343.9789.

7-Amino-6-methoxyquinoline-5,8-quinone (6a): A stirred suspension of 7-bromo-6-methoxyquinoline-5,8-quinone (**8a**, 33 mg, 0.123 mmol) in 0.6 ml of THF was treated with a solution of sodium azide (8.8 mg, 0.135 mmol, 1.1 equiv) in 0.08 ml of H₂O at 25 °C under a N₂ atmosphere and the mixture was stirred at 25 °C for 15 h with protection from light. The suspension was poured onto 20 ml of H₂O and extracted with CH₂Cl₂ (3 x 15 ml). The combined extracts were dried (Na₂SO₄) and concentrated in vacuo. Chromatography (SiO₂, 1 x 20 cm, 50% EtOAc-hexane eluant) afforded 23.0 mg (28.3 mg theor., 81%) of 7-azido-6-methoxyquinoline-5,8-quinone as an orange-yellow solid: mp 132-133 °C dec. (lit.^{13a} mp 131.5-132 °C dec.); ¹H NMR (CDCl₃) δ 8.99 (1H, dd, J = 4.7 Hz, 1.6 Hz, C-2 H), 8.40 (1H, dd, J = 7.8 Hz, 1.6 Hz, C-4 H), 7.66 (1H, dd, J = 7.8 Hz, 4.7 Hz, C-3 H), 4.24 (3H, s, OCH₃).

A stirred suspension of the azodiquinoline-5,8-quinone (7.0 mg, 0.0304 mmol) in 0.15 ml of THF was treated with a solution of sodium hydrosulfite (18.5 mg, 0.106 mmol, 3.5 equiv) in 0.08 ml of H₂O at 25 °C under a N₂ atmosphere and the mixture was stirred at 25 °C for 2 h. The mixture was poured onto H₂O (15 ml) and extracted with CH₂Cl₂ (3 x 10 ml). The combined extracts were dried (Na₂SO₄) and concentrated in vacuo. Chromatography (SiO₂, 1 x 12 cm, 60% EtOAc-hexane eluant)

afforded 3.6 mg (6.2 mg theor., 58%) of **6a** as a purple solid: mp 201-202 °C dec. (lit.^{13a} mp 202-203 °C dec.); ¹H NMR (CDCl₃) δ 8.88 (1H, dd, J = 4.7 Hz, 1.7 Hz, C-2 H), 8.37 (1H, dd, J = 7.8 Hz, 1.7 Hz, C-4 H), 7.60 (1H, dd, J = 7.8 Hz, 4.8 Hz, C-3 H), 5.21 (2H, br s, NH₂), 4.07 (3H, s, OCH₃).

7-Amino-6-methoxy-2-(2'-pyridyl)quinoline-5,8-quinone (6b). A stirred suspension of 7-bromo-6-methoxy-2-(2'-pyridyl)quinoline-5,8-quinone (**8b**, 11.5 mg, 0.0333 mmol) in 0.5 ml of DMF and 0.5 ml of methanol was treated with powdered sodium azide (2.4 mg, 0.0366 mmol, 1.1 equiv) at 25 °C under a N₂ atmosphere and the mixture was stirred at 25 °C for 16 h with protection from light. The crystals were collected, washed with 50% methanol, dried *in vacuo* to give the azidoquinoline-5,8-quinone as an orange solid (6.2 mg, 10.2 mg theor., 61%). The filtrate was extracted with CH₂Cl₂ (1 x 15 ml) and the extract was dried (Na₂SO₄) and concentrated *in vacuo*. Chromatography (SiO₂, 1 x 9.5 cm, 50% EtOAc-hexane eluant) afforded an additional 3.9 mg (10.1 mg total, 99%) of the azidoquinoline-5,8-quinone as an orange solid: mp 136-138 °C (lit.^{15a} mp 137-139 °C); ¹H NMR (CDCl₃) δ 8.82 (1H, d, J = 8.3 Hz, C-4 H), 8.60-8.83 (2H, m, C-3' H and C-6' H), 8.48 (1H, d, J = 8.3 Hz, C-3 H), 7.89 (1H, dt, J = 8.0 Hz, 1.6 Hz, C-4' H), 7.32-8.00 (1H, m, C-5' H), 4.26 (3H, s, OCH₃).

A stirred suspension of the azidoquinoline-5,8-quinone (7.0 mg, 0.0228 mmol) in 50% aqueous methanol (0.8 ml) was treated with sodium hydrosulfite (13.9 mg, 0.0798 mmol, 3.5 equiv) under a N₂ atmosphere at 25 °C for 21 h. The mixture was diluted with H₂O (20 ml) and extracted with EtOAc (3 x 15 ml). The combined extracts were dried (Na₂SO₄) and concentrated *in vacuo*. Chromatography (SiO₂, 1 x 8.5 cm, 80% EtOAc-hexane eluant) afforded 4.2 mg (6.4 mg theor., 65%) of **6b** as a purple solid: mp 172-173.5 °C (lit.^{15b} mp 172-174 °C); ¹H NMR (CDCl₃) δ 8.75 (1H, d, J = 8.2 Hz, C-4 H), 8.57-8.76 (2H, m, C-3' H and C-6' H), 8.46 (1H, d, J = 8.2 Hz, C-3 H), 7.87 (1H, dt, J = 7.7 Hz, 1.8 Hz, C-4' H), 7.34-7.97 (1H, m, C-5' H), 5.20 (2H, br s, NH₂), 4.10 (3H, s, OCH₃).

ACKNOWLEDGMENTS

This work was assisted financially by the National Institutes of Health (CA 00898/01134, 33668/42056) and Searle Scholars Fund.

REFERENCES AND NOTES

1. National Institutes of Health Research Career Development Award Recipient 1983-88, (CA 00898/01134). Searle Scholar Recipient, 1981-85. Alfred P. Sloan research fellow, 1985-89.
2. K. V. Rao and W. P. Cullen, *Antibiot. Annu.*, **1959-1960**, 950; K. V. Rao, K. Biemann, and R. B. Woodward, *J. Am. Chem. Soc.*, **1963**, 65, 2532; Y.-Y. H. Chiu and W. N. Lipscomb, *J. Am. Chem. Soc.*, **1975**, 97, 2525.


3. T. W. Doyle, D. M. Balitz, R. E. Grulich, D. E. Nettleton, S. J. Gould, C. Tann, and A. E. Moews, Tetrahedron Lett., **1981**, 22, 4595; D. M. Balitz, J. A. Bush, W. T. Bradner, T. W. Doyle, F. A. O'Herron, and D. E. Nettleton, J. Antibiotics, **1982**, 35, 259.
4. A. J. Herlt, R. W. Rickards, and J. P.-Wu, J. Antibiotics, **1985**, 38, 516.
5. For a recent review, see: S. J. Gould and S. M. Weinreb, Fortschr. Chem. Org. Naturst., **1982**, 41, 77. For recent biosynthetic studies, see: S. J. Gould and D. E. Cane, J. Am. Chem. Soc., **1982**, 104, 343; W. R. Erickson and S. J. Gould, J. Am. Chem. Soc., **1985**, 107, 5831; S. J. Gould, C. C. Chang, D. S. Darling, J. D. Roberts, and M. Squillacote, J. Am. Chem. Soc., **1980**, 102, 1707.
6. K. V. Rao, Cancer Chemother. Rep., **1974**, 4, 11; N. I. Nissen, T. Pajak, I. Glidewell, H. Blom, M. Flaherty, D. Hayes, R. McIntyre, and J. F. Hollond, Cancer Treatment Rep., **1977**, 61, 1097.
7. D. L. Boger, M. Yasuda, J. S. Panek, and P. A. Kitos, unpublished observations. For instance, in vitro cytotoxic activity L1210 and 9PS: ID₅₀ streptonigin = 0.61 and 0.045 μ g/mL respectively; ID₅₀ **7a** = 0.34 and 0.19 μ g/mL respectively; ID₅₀ **7b** = 0.42 and 0.0023 μ g/mL respectively; ID₅₀ **6a** = 1.0 and 0.4 μ g/mL respectively; ID₅₀ **6b** = nt and 0.034 μ g/mL respectively.
8. D. L. Boger and J. S. Panek, J. Am. Chem. Soc., **1985**, 107, 5745; D. L. Boger and J. S. Panek, J. Org. Chem., **1983**, 48, 621; S. M. Weinreb, F. Z. Basha, S. Hibino, N. A. Khatri, D. Kim, W. E. Pye, and T.-T. Wu, J. Am. Chem. Soc., **1982**, 104, 536; A. S. Kende, D. P. Lorah, and R. J. Boatman, J. Am. Chem. Soc., **1981**, 103, 1271.
9. D. L. Boger and M. Yasuda, unpublished observations.
10. (a) A. S. Kende and F. H. Ebetino, Tetrahedron Lett., **1984**, 25, 923; S. Hibino, M. Okazaki, M. Ichikawa, K. Sato, and T. Ishizu, Heterocycles, **1985**, 23, 261. (b) D. L. Boger, S. R. Duff, J. S. Panek, and M. Yasuda, J. Org. Chem., **1986**, 51, 0000.
11. For studies on the introduction of the streptonigrin 7-amino-6-methoxyquinoline-5,8-quinone AB ring system, see: reference 5, 8 and 13, and references cited therein. For studies detailing the introduction of the lavendamycin 7-aminoquinoline-5,8-quinone AB ring system, see: reference 10, 13 and references cited therein.
12. Less successful direct and indirect approaches for the conversion of 7-bromoquinoline-5,8-quinones to the streptonigrin 7-amino-6-methoxyquinoline-5,8-quinone AB ring system are summarized below in Scheme I.
13. For the initial observation that azide will directly displace C-7 halides of 7-haloquinoline-5,8-quinones, see: (a) T. K. Liao, P. J. Wittek, and C. C. Cheng, J. Heterocycl. Chem., **1976**, 13, 1283; T. K. Liao, W. H. Nyberg, and C. C. Cheng, J. Heterocycl. Chem., **1976**, 13, 1063; Idem Angew. Chem. Int. Ed. Engl., **1967**, 6, 82. For further studies on this observation and its

implementation in the preparation of 7-azidoquinoline-5,8-quinones, see: (b) H. S. Kuo and S. Yoshina, *Yakugaku Zasshi*, **1977**, 97, 827.

14. For initial studies on the utility of cerium(III) promoted nucleophilic substitution and addition reactions of haloquinones, see: Y. T. Pratt, *J. Org. Chem.*, **1962**, 27, 3905 and reference 13b.

15. (a) S. Hibino and S. M. Weinreb, *J. Org. Chem.*, **1977**, 42, 232. (b) K. V. Rao, *J. Heterocyclic Chem.*, **1975**, 12, 725.

Scheme I



Received, 2nd December, 1985